



Global, regional, and national disability-adjusted life-years (DALYs) for 333 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016



GBD 2016 DALYs and HALE Collaborators*

Summary

Lancet 2017; 390: 1260–344

*Collaborators listed at the end of the Article

Correspondence to:

Prof Simon Iain Hay, Institute for Health Metrics and Evaluation, Seattle, WA 98121, USA
sihay@uw.edu

Background Measurement of changes in health across locations is useful to compare and contrast changing epidemiological patterns against health system performance and identify specific needs for resource allocation in research, policy development, and programme decision making. Using the Global Burden of Diseases, Injuries, and Risk Factors Study 2016, we drew from two widely used summary measures to monitor such changes in population health: disability-adjusted life-years (DALYs) and healthy life expectancy (HALE). We used these measures to track trends and benchmark progress compared with expected trends on the basis of the Socio-demographic Index (SDI).

Methods We used results from the Global Burden of Diseases, Injuries, and Risk Factors Study 2016 for all-cause mortality, cause-specific mortality, and non-fatal disease burden to derive HALE and DALYs by sex for 195 countries and territories from 1990 to 2016. We calculated DALYs by summing years of life lost and years of life lived with disability for each location, age group, sex, and year. We estimated HALE using age-specific death rates and years of life lived with disability per capita. We explored how DALYs and HALE differed from expected trends when compared with the SDI: the geometric mean of income per person, educational attainment in the population older than age 15 years, and total fertility rate.

Findings The highest globally observed HALE at birth for both women and men was in Singapore, at 75·2 years (95% uncertainty interval 71·9–78·6) for females and 72·0 years (68·8–75·1) for males. The lowest for females was in the Central African Republic (45·6 years [42·0–49·5]) and for males was in Lesotho (41·5 years [39·0–44·0]). From 1990 to 2016, global HALE increased by an average of 6·24 years (5·97–6·48) for both sexes combined. Global HALE increased by 6·04 years (5·74–6·27) for males and 6·49 years (6·08–6·77) for females, whereas HALE at age 65 years increased by 1·78 years (1·61–1·93) for males and 1·96 years (1·69–2·13) for females. Total global DALYs remained largely unchanged from 1990 to 2016 (–2·3% [–5·9 to 0·9]), with decreases in communicable, maternal, neonatal, and nutritional (CMNN) disease DALYs offset by increased DALYs due to non-communicable diseases (NCDs). The exemplars, calculated as the five lowest ratios of observed to expected age-standardised DALY rates in 2016, were Nicaragua, Costa Rica, the Maldives, Peru, and Israel. The leading three causes of DALYs globally were ischaemic heart disease, cerebrovascular disease, and lower respiratory infections, comprising 16·1% of all DALYs. Total DALYs and age-standardised DALY rates due to most CMNN causes decreased from 1990 to 2016. Conversely, the total DALY burden rose for most NCDs; however, age-standardised DALY rates due to NCDs declined globally.

Interpretation At a global level, DALYs and HALE continue to show improvements. At the same time, we observe that many populations are facing growing functional health loss. Rising SDI was associated with increases in cumulative years of life lived with disability and decreases in CMNN DALYs offset by increased NCD DALYs. Relative compression of morbidity highlights the importance of continued health interventions, which has changed in most locations in pace with the gross domestic product per person, education, and family planning. The analysis of DALYs and HALE and their relationship to SDI represents a robust framework with which to benchmark location-specific health performance. Country-specific drivers of disease burden, particularly for causes with higher-than-expected DALYs, should inform health policies, health system improvement initiatives, targeted prevention efforts, and development assistance for health, including financial and research investments for all countries, regardless of their level of sociodemographic development. The presence of countries that substantially outperform others suggests the need for increased scrutiny for proven examples of best practices, which can help to extend gains, whereas the presence of underperforming countries suggests the need for devotion of extra attention to health systems that need more robust support.

Funding Bill & Melinda Gates Foundation.

Copyright © The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

Introduction

Objective measurement of population health is a fundamental requirement of good governance that allows international, regional, national, and local actors to frame evidence-based policy informed by past trends and current performance of health systems.¹⁻⁴ Summary measures of population health include techniques that measure the overall burden of health loss due to fatal and non-fatal diseases, as well as measures of expected fatal and non-fatal disease burden based on Socio-demographic Index (SDI).⁵ The disability-adjusted life-year (DALY) measures health loss due to both fatal and non-fatal disease burden. DALYs are the sum of the years of life lost (YLLs) due to premature mortality and years of life lived with disability (YLDs).⁶ The YLL is based on remaining life expectancy when compared with a reference standard life table at age

of death,⁷ and the YLD is calculated by multiplying the prevalence of a disease or injury and its main disabling outcomes by its weighted level of severity.^{6,8} One DALY represents 1 year of healthy life lost. Examination of levels and trends of DALYs facilitates quick comparison between different diseases and injuries. Conversely, healthy life expectancy (HALE), a metric based on methods by Sullivan,⁹ provides a single summary measure of population health across all causes combined by weighting years lived with a measure of functional health loss before death and is the most comprehensive among competing expectancy metrics.¹⁻⁴ Together, DALYs and HALE enable comparisons of the magnitude of functional health loss across societies due to diseases, injuries, and risk factors, against which provisioning and performance of health systems can be calibrated.⁴

Research in context

Evidence before this study

The Global Burden of Diseases, Injuries, and Risk Factors Study 2015 (GBD 2015) provided disability-adjusted life-year (DALY) estimates for 315 diseases and injuries for 195 countries and territories, including subnational assessments for 11 countries and, thus, for a total of 519 locations, from 1990 to 2015. GBD 2015 also introduced analyses of DALYs and healthy life expectancy (HALE) in relation to the Socio-demographic Index (SDI). Only the WHO Global Health Estimates has published updated estimates of DALYs, and these estimates were heavily reliant on GBD 2015 results.

Added value of this study

This study, the Global Burden of Diseases, Injuries, and Risk Factors Study 2016 (GBD 2016), updates and improves the first of the annual Global Burden of Disease iterations, GBD 2015. GBD 2016 is, to our knowledge, the only peer-reviewed, Guidelines for Accurate and Transparent Health Estimates Reporting-compliant, comprehensive, and annual assessment of DALYs and HALE by age group, sex, cause, and location, analysed consistently from 1990 to 2016. The improved approaches to the analysis and refinements in data (gap fills, updates, and revisions), as well as the widening of scope by cause, location, age, and time are all relevant to this study. The summary population health metrics of DALYs and HALE synthesise the cumulative effect of all of these improvements, the most notable of which are as follows. First, we added substantial location-years of cause-specific mortality data and non-fatal data for GBD 2016. The added data progressively fill gaps in the period of estimation, most substantially for India. Second, many analytical methods have been improved, such as improvement of mortality to incidence ratios for cancers to better reflect lower survival than in GBD 2015 and for non-fatal tuberculosis to better reflect higher incidence in low-income and middle-income countries based on SDI. Third, we included new subnational assessments

for Indonesia at the provincial level and further disaggregated subnational estimation in England to the local government area level. Fourth, we refined our estimation of age-specific outcomes for ages 80 years and older into 5 year groups extending to age 95 years and older to better account for disease burden in elderly populations than in GBD 2015. Fifth, we estimated DALYs for several additional causes for the first time. Sixth, we improved our analysis of the epidemiological transition as a function of SDI, which allowed for a more nuanced interpretation of global health trends against the sociodemographic development spectrum than in GBD 2015. Finally, we used these analyses to identify the exemplar countries that exceeded population health summary metric expectations relative to their SDI position alone. The GBD 2016 iteration supersedes all previous GBD studies of DALYs and HALE and re-estimates these measures for the complete time series from 1990 to 2016. We focus on new methods and approaches since GBD 2015 and highlight nations that overperformed or underperformed on the basis of what would be expected on the basis of their SDI.

Implications of all the available evidence

The epidemiological transition continues apace globally, with a shift from DALYs attributable to communicable, maternal, neonatal, and nutritional diseases to those attributable to non-communicable diseases. This progression is concomitant with improvements in SDI and thus improvements in education, fertility rates, and economic status. A more detailed analysis than in this study of the epidemiological changes that have occurred in countries that have consistently exceeded expectations could provide improved insights into good practice in public health policy, which might be emulated elsewhere. A similarly detailed appraisal of countries that are lagging in DALYs and HALE relative to expectations on the level of SDI alone will help identify countries in most need of domestic and international attention across the development continuum.

As the second in a series^{7,8,10,11} of now annual updates, the Global Burden of Diseases, Injuries, and Risk Factors Study 2016 (GBD 2016) is the most comprehensive and current source of summary health metrics. The Global Burden of Disease (GBD) is based on development of the largest available database of health outcomes, risk factor exposure, intervention coverage, and sociodemographic factors related to health. We applied analytical techniques to reduce data biases and support comparability, propagated the uncertainty in these estimates, and provided insights at the highest temporal and spatial resolution afforded by the data.

The purpose of this study is to present the results of GBD 2016 for DALYs and HALE, building on updated estimates of mortality, causes of death, and non-fatal health loss^{7,10} to identify nations with notable variation in health performance from that expected on the basis of SDI. Approaches to the analysis have been previously described.^{2-4,12} GBD 2016 improvements include addition of newly available retrospective data, refined analytical methods (such as improvement to mortality to incidence ratios [MIRs] for cancers to better reflect lower survival in low-income and middle-income countries based on SDI), new subnational estimation for England and Indonesia, disaggregation of certain cause groupings to capture greater detail, and expansion of older age groups to enhance relevance for a wider range of health policy decisions.⁶

Methods

Overview

We used the results of GBD 2016 to evaluate trends in epidemiological patterns and health performance on a global, regional, national, and subnational scale using DALYs and HALE as summary measures of changes in health states. Greater detail than presented in this section for methods used to estimate DALYs and HALE, including analytic approaches for assessment of relative morbidity and mortality from individual diseases and injuries, is provided in related publications in this series^{8,10} and the appendix.

This analysis follows the Guidelines for Accurate and Transparent Health Estimates Reporting,^{13,14} which include recommendations on documentation of data sources, estimation methods, and statistical analysis. We did analyses using Python version 2.7.12 and 2.7.3, Stata version 13.1, and R version 3.2.2. For more information on Guidelines for Accurate and Transparent Health Estimates Reporting compliance, please refer to the appendix (pp 13–15). Additionally, interactive online tools are available to explore GBD 2016 data sources in detail. Cause-specific estimation for GBD 2016 covers the years 1990–2016. For a subset of analyses, we focus on the last decade, from 2006 to 2016, to address current policy priorities. The GBD 2016 results for all years and by location can be explored further with dynamic data visualisations.

Cause and location hierarchies

In the GBD 2016 study, causes of mortality and morbidity are structured with use of a four-level classification hierarchy to produce levels that are mutually exclusive and collectively exhaustive. GBD 2016 estimates 333 causes of DALYs, 68 of which are a source of disability but not a cause of death (such as trachoma, hookworm, and low back and neck pain) and five of which are causes of death but not sources of morbidity (sudden infant death syndrome, aortic aneurysm, late maternal deaths, indirect maternal deaths, and maternal deaths aggravated by HIV/AIDS). Within each level of the hierarchy, the number of collectively exhaustive and mutually exclusive fatal and non-fatal causes for which the GBD study estimates is three at Level 1, 21 at Level 2, 168 at Level 3, and 276 at Level 4. The full GBD cause hierarchy, including corresponding International Classification of Diseases (ICD)-9 and ICD-10 codes, is detailed in GBD 2016 publications on cause-specific mortality¹⁰ and non-fatal health outcomes,⁸ with cause-specific methods detailed in the corresponding appendices.

The GBD study is organised by a geographical hierarchy of seven super-regions containing 21 regions, with 195 countries and territories nested within those regions.¹² GBD 2016 included new subnational assessments for Indonesia by province and for England by local government area. In this study, we present subnational data for the five countries with a population greater than 200 million people in 2016: Brazil, China, India, Indonesia, and the USA.

Estimation of mortality and non-fatal health loss

To estimate all-cause and cause-specific mortality, the GBD study first systematically addressed known data challenges—such as variation in coding of causes or age group reporting, misclassification of deaths from HIV/AIDS, or methods for incorporation of population-based cancer registry data—using standardised methods described in detail in the GBD 2016 mortality⁷ and causes of death¹⁰ publications. As noted in other GBD publications, each death is attributed to a single underlying cause in accordance with the ICD. We take steps to standardise cause of death data to address the small fraction of deaths that are not assigned an age or sex, deaths assigned to broad age groups that are not 5 year age groups, and various revisions and national variants of the ICD. Additionally, we identify and redistribute deaths assigned to ICD codes that cannot be underlying causes of death, are intermediate causes of death rather than the underlying causes, or lack specificity in coding.¹⁰ We estimated cause-specific mortality using standardised modelling processes—most commonly, the Cause of Death Ensemble model, which uses covariate selection and out-of-sample validity analyses and generates estimates for each location-year, age, and sex.¹⁰ Additional detail, including model specifications and data availability for each cause-specific model, can be found in the

See Online for appendix

For the online tools see
<http://ghdx.healthdata.org>

For the data visualisations see
<https://vizhub.healthdata.org/gbd-compare>

appendix of the GBD 2016 mortality⁷ and causes of death¹⁰ publications. We used the all-cause mortality estimates to establish a reference life table from the lowest death rates for each age group among locations with total populations greater than 5 million.⁷ From this reference life table, we multiplied life expectancy at the age of death by cause-specific deaths to calculate cause-specific YLLs. We then used the GBD world population age standard to calculate age-standardised rates for deaths and YLLs.⁷ The GBD world population age standard and the standard life expectancies are available in the appendix of the GBD 2016 mortality publication.⁷

Changes implemented since the Global Burden of Diseases, Injuries, and Risk Factors Study 2015 (GBD 2015) for cause-specific mortality include incorporation of substantial sources of new mortality data; important model improvements for HIV, malaria, tuberculosis, injuries, diabetes, and cancers; disaggregation of specific causes into subgroupings to provide additional detail (the following were all estimated separately for the first time: alcoholic cardiomyopathy; urogenital, musculoskeletal, and digestive congenital anomalies; Zika virus disease; Guinea worm disease; self-harm by firearm; sexual violence; myocarditis; and the following types of tuberculosis: extensively drug-resistant tuberculosis, multidrug-resistant tuberculosis without extensive drug resistance, drug-susceptible tuberculosis, extensively drug-resistant HIV/AIDS-tuberculosis, multidrug-resistant HIV/AIDS-tuberculosis without extensive drug resistance, and drug-susceptible HIV/AIDS-tuberculosis); modelling of antiretroviral therapy (ART) coverage for each location-year by CD4-positive cell count at initiation; breakdown of terminal age groups from 80 years and older to 80–84 years, 85–89 years, 90–94 years, and 95 years and older; expansion of the GBD location hierarchy; and changes in the calculation of SDI.¹⁰ The database for GBD 2016 now includes data for the 333 causes estimated for DALYs and new subnational units for Indonesia (n=34) and England (n=150). For GBD 2016, we included substantial amounts of additional data sources from new studies and our network of collaborators; details of the types of data added can be found in the GBD 2016 cause of death¹⁰ and non-fatal⁸ publications. Additionally, research teams did systematic reviews to incorporate literature data into fatal and non-fatal models. Further details on search strings are available in the GBD 2016 non-fatal⁸ and cause of death¹⁰ publication appendices. The Registrar General of India provided improved verbal autopsy data collected through their Sample Registration System, enabling a more detailed and thorough analysis of subnational data for India than in GBD 2015. The methods for constructing the SDI, initially developed for GBD 2015,¹⁵ were revised for GBD 2016 to account for expansion in the number of subnational estimates and the effect of a growing time period of estimation given fixed limits for index components.¹⁰ The components of SDI—total fertility rate (TFR), educational attainment in the population aged older

than 15 years, and lag-distributed income (LDI)—are based on new systematic assessments of educational attainment, LDI, and fertility, and each component is scaled relative to maximum effect on health outcomes.¹⁰

In most cases, we estimated non-fatal health loss using the Bayesian meta-regression tool DisMod-MR 2.1 to synthesise variable data sources to produce internally consistent estimates of incidence, prevalence, remission, and excess mortality.¹⁶ Cause-specific data availability and epidemiological characteristics required additional analytical techniques in some cases (details are available in the appendix of the GBD 2016 non-fatal publication⁸); these causes include many neglected tropical diseases (NTDs) such as dengue, as well as injuries, malaria, and HIV/AIDS.^{17,18}

We estimated each non-fatal sequela separately and assessed the occurrence of comorbidity in each age group, sex, location, and year separately using a microsimulation framework. We distributed disability estimated for comorbid conditions to each contributing cause during the comorbidity estimation process. Although the distribution of sequelae—and therefore the severity and cumulative disability per case of a condition—can be different by age, sex, location, and year, previous studies have found that disability weights do not substantially vary across locations, income, or levels of educational attainment.^{19,20} In the GBD study, disability weights were based on population surveys with 60 890 respondents and held invariant between locations and over time.²⁰ Additional details, including model specifications and data availability for each cause-specific model and development of disability weights by cause and their use in the estimation of non-fatal health loss, are available in the appendix of the GBD 2016 non-fatal publication.⁸

For non-fatal estimation, several methodological changes were made for GBD 2016. New data for the main causes of YLDs were identified through our collaboration with the Indian Council of Medical Research and the Public Health Foundation of India. For particular risk factors and diseases, the volume of available data increased substantially, such as child growth failure (stunting, wasting, or underweight), anaemia, congenital anomalies, schistosomiasis, intestinal helminths, and lymphatic filariasis. We have improved our analysis of total admissions per person by country, year, age, and sex, which facilitated incorporation of additional hospital data sources that were previously excluded because of incomplete knowledge of catchment population size. We extended our analyses of linked USA medical claims data to impute age-specific and sex-specific ratios for multiple admissions per illness episode, ICD code appearance in the non-primary position, and inpatient versus outpatient use.⁸ We applied each of the three ratios sequentially to non-linked hospital inpatient data from elsewhere that only had a single ICD code per visit to adjust prevalence and incidence data. We have incorporated more predictive covariates into our non-fatal disease models to

better predict variation in disease levels rather than measurement error as the source of variation, and we improved our analysis of the MIRs for cancers, resulting in considerably higher ratios in lower SDI quintiles and thus substantially lower YLD estimates for cancer.

For the statistical code see <https://github.com/ihmeuw/ihme-modeling>

Estimation of DALYs, HALE, and corresponding uncertainty

We calculated DALYs as the sum of YLLs and YLDs for each cause, location, age group, sex, and year.^{8,10} The same estimates of YLDs per person for each location, age, sex, and year from 1990 to 2016 are used to establish HALE by age group within abridged multiple-decrement life tables with use of methods developed by Sullivan.⁹

For all results, we report 95% uncertainty intervals (UIs) derived from 1000 draws from the posterior distribution of each step in the estimation process. Unlike confidence intervals, UIs capture uncertainty from multiple modelling steps, as well as from sources such as model estimation and model specification, rather than from sampling error alone. Uncertainty associated with estimation of mortality and YLLs reflects sample sizes of data sources, adjustment and standardisation methods applied to data, parameter uncertainty in model estimation, and uncertainty within all-cause and cause-specific mortality models. For estimation of prevalence, incidence, and YLDs, UIs incorporated variability from sample sizes within data sources, adjustments to data to account for non-reference definitions, parameter uncertainty in model estimation, and uncertainty associated with establishment of disability weights. Because direct information about the correlation between uncertainty in YLLs and YLDs was scarce, we assumed that uncertainty in age-specific YLDs was independent of age-specific YLLs or death rates.

Epidemiological transition and relationship between DALYs, HALE, and SDI

For GBD 2016, the composite indicator of SDI was again based on the geometric mean of three measures—LDI per person, average years of schooling among populations aged 15 years or older, and TFR—but the analysis was strengthened in three important ways.¹⁰ First, we substantially revised estimates of education, adding new data and improved methods for subnational locations. Second, instead of using estimates of TFR from the UN Population Division, we systematically reviewed, extracted, and analysed fertility data from all available locations to derive a time series of TFR for each national and subnational GBD location.⁷ Third, rather than rescaling SDI on the basis of the full range of observed values within the time series, we developed a fixed scale for GBD 2016; details on development of this fixed scale are available in the GBD 2016 mortality publication.⁷ We examined the average relationship between DALYs, HALE, and SDI using a Gaussian process regression model; we used these regressions to estimate expected values of these summary measures

at each level of SDI. Additional detail on SDI calculation and location-specific SDI values are available in the appendix of the GBD 2016 mortality publication.⁷

Data sharing

The statistical code used in the entire process is available through an online repository.

Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Results

Global levels of and trends for DALYs and HALE

The total number of all-age DALYs in 2016 was 2.39 billion (95% UI 2.18 billion to 2.63 billion). Total all-age DALY counts for CMNN causes fell by 40.1% (37.4–42.7) from 1.11 billion (1.07 billion to 1.16 billion) in 1990 to 668 million (632 million to 708 million) in 2016, whereas total all-age DALY counts from NCDs increased by 36.6% from 1.07 billion (958 million to 1.20 billion) in 1990 to 1.47 billion (1.30 billion to 1.66 billion) in 2016 (table 1). Total DALYs from injuries decreased by 1.6% (–3.8 to 6.2) from 260 million (243 million to 277 million) in 1990 to 255 million (236 million to 281 million) in 2016. Age groups older than 80 years had 149 million (139 million to 159 million) all-age DALYs in 2016 compared with 75.1 million (71.1 million to 79.5 million) in 1990, with increases across all SDI quintiles. Of these, 87.8% were due to NCDs in 2016 compared with 86.8% in 1990.

From 1990 to 2016, global HALE at birth increased from 56.9 years to 63.1 years, with 160 of 195 locations registering significant improvements. Global HALE increased by an average of 6.24 years (95% UI 5.97–6.48) for both sexes combined. Globally, HALE at birth increased from 55.38 years (53.27–57.31) in 1990 to 61.42 years (59.01–63.58) in 2016 for males and from 58.42 years (55.80–60.77) to 64.91 years (61.88–67.54) for females, rising 6.04 years (5.74–6.27) for males and 6.49 years (6.08–6.77) for females (tables 2 and 3). The total number of years of functional health lost (life expectancy minus HALE) increased from 1990 to 2016, from 8.22 years to 9.34 years. The gap between life expectancy at birth and HALE, which represents years of functional health lost, grew between 1990 and 2016 from 7.32 years (life expectancy 62.70 [62.42–62.99] vs HALE 55.38 [53.27–57.31]) to 8.37 years (69.79 [69.29–70.22] vs 61.42 [59.01–63.58]) for males and from 9.15 years (67.57 [67.33–67.77] vs 58.42 [55.80–60.77]) to 10.42 years (75.33 [74.95–75.64] vs 64.91 [61.88–67.54]) for females. Globally, in 2016, life expectancy at age 65 years was 18.57 years (18.37–18.72) for females and 15.72 years (15.61–15.83) for males,

whereas HALE was 13.88 years (12.57–15.02) for females and 11.87 years (10.83–12.80) for males. HALE increased by 1.96 years (1.69–2.13) from 11.92 (10.88–12.89) in 1990 for females and by 1.78 years (1.61–1.93) from 10.09 years (9.22–10.87) for males.

Global trends for all-age DALYs and age-standardised DALY rates for Level 1 causes by SDI quintile are shown in figure 1. Trends in total DALYs, which show the absolute burden at each SDI quintile, are shown in figure 1A. The figure highlights the large burden and subsequent declines in low-middle-SDI (decreased by 44.5% [95% UI 41.3–47.6]) and middle-SDI (decreased by 56.2% [53.6–59.0]) locations for CMNN DALYs from 1990 to 2016, offset by large increases for NCD DALYs over the same time period in low-middle SDI (increased by 54.4% [47.9–60.8]) and middle-SDI (increased by 36.5% [39.9–32.8]) locations. Progress has been made in low-SDI countries for CMNN DALYs, which decreased by 20.7% (16.8–24.7) since 2006. At all levels of SDI, total NCD DALYs have increased since 1990. Trends in age-standardised rates—which account for both population size and age structure—emphasise the reduction in the contribution of CMNN causes to DALYs, both over time and with increasing SDI (figure 1B). These rapid decreases in age-standardised rates were fastest at low SDI, where age-standardised DALY rates from CMNN causes decreased by 51.5% (49.0–54.1) between 1990 and 2016 to be on par with age-standardised DALY rates for NCDs in 2016. At low-middle SDI, age-standardised CMNN DALY rates were more than double those for NCDs in 1990 (31.7 thousand [30.3 thousand to 33.3 thousand] per 100 000), but decreased to 14.1 thousand (13.2 thousand to 15.1 thousand) per 100 000 in 2016. At all other levels of SDI, age-standardised DALY rates for CMNN causes were lower than those for NCDs and lower than those of injuries in the high-SDI quintile. Reductions in age-standardised rates for NCDs occurred across all levels of SDI between 1990 and 2016, a trend that was also evident, albeit less strongly, for age-standardised DALY rates from injuries.

Global causes of DALYs

Age-standardised DALY rates for all causes decreased by 30.5% (95% UI 28.6–32.6) between 1990 and 2016 (appendix pp 49–62). In 2016, CMNN causes accounted for 28.0% (26.4–29.7) of global DALYs, NCDs contributed 61.4% (59.4–63.2), and injuries contributed 10.7% (10.1–11.3; appendix pp 35–48). From 2006 to 2016, CMNN causes decreased by 31.9% (29.7–34.2), with 48 Level 4 CMNN causes experiencing decreases in age-standardised DALY rates of greater than 20% (table 1). Decreases were greater than 70% for three infectious diseases: Guinea worm disease (decreased by 99.6% [99.5–99.7]), human African trypanosomiasis (decreased by 78.2% [68.8–84.6]), and measles (decreased by 73.6% [68.8–77.8]). By contrast with the overall trend of decreasing DALYs, a subset of Level 4 CMNN causes had increases in age-standardised

DALY rates, including dengue (50.5% [24.7–97.7]) and cutaneous and mucocutaneous leishmaniasis (12.5% [1.7–26.1]). Overall, total all-age DALYs attributable to maternal disorders decreased by 23.9% (17.4–29.3) between 2006 and 2016 and by 30.4% (24.4–35.4) in terms of age-standardised DALY rates. As a cause group, neonatal disorders decreased by 22.8% (18.9–26.8) in all-age DALYs and 23.1% (19.2–27.0) in terms of age-standardised DALY rates over the same time period; however, this decrease was not significant for neonatal sepsis. Total DALYs from the London Declaration NTDs was 9.0 million (5.3 million to 14.5 million) in 2016.

In 2016, the leading Level 3 causes of total DALYs among NCDs included ischaemic heart disease (175 million [95% UI 170 million to 180 million] DALYs), cerebrovascular disease (116 million [111 million to 121 million]), and low back and neck pain (87 million [61 million to 114 million]), comprising 16.1% (13.99–17.67) of all DALYs (table 1). Among chronic respiratory diseases, all causes, with the exception of interstitial lung disease, pulmonary sarcoidosis, and other chronic respiratory diseases, decreased in age-standardised DALY rates between 2006 and 2016, whereas total all-age DALY counts increased from 2006 to 2016 for all chronic respiratory diseases, with the exception of silicosis. Cirrhosis and other chronic liver diseases had a mean change of 7.6% (2.5–13.7) from 2006 to 2016 in total all-age DALY counts, but had a mean decrease in age-standardised DALY rates of 12.0% (7.2–16.1) over the same period. Age-standardised DALY rates of digestive diseases decreased from 2006 to 2016, with a mean percentage decrease of 13.6% (10.8–16.4); however, all-age DALY counts for digestive diseases increased by 4.1% (0.4–7.7) over the same period. Total DALYs associated with most neurological disorders increased from 2006 to 2016, with Alzheimer's disease and other dementias (increase of 37.5% [35.3–39.7]) and Parkinson's disease (increase of 35.6% [32.9–38.2]) increasing by more than 30% each. Between 2006 and 2016, various NCDs significantly increased in terms of both total burden and age-standardised DALY rates. Several mental and substance use disorders followed this pattern, including eating disorders (age-standardised DALY rates increased by 8.9% [7.6–10.1]) and bipolar disorder (increased by 0.8% [0.2–1.4]). Diabetes (all-age DALY count increased by 24.4% [22.7–26.2]) and chronic kidney disease (increased by 20.0% [17.4–22.7]) both also increased in all-age DALY counts, as did musculoskeletal disorders (increased by 19.6% [18.5–20.8]).

Percentage change in age-standardised DALY rates of unintentional injuries (decreased by 15.1% [95% UI 12.2–18.0]), road injuries (decreased by 14.7% [12.8–16.8]), and transport injuries (decreased by 14.3% [12.3–16.4]) each decreased substantially between 2006 and 2016 (table 1). Among unintentional injuries, drowning had the largest reduction in both all-age DALY burden (26.6% [20.1–30.1]) and age-standardised DALY rates (32.1%

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|---|--|--|--|-----------------------------------|-----------------------------------|---|---|---|-----------------------------------|-----------------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 |
| All causes | 2 448 430.5 (2 305 218.2 to 2 608 339.5) | 2 490 698.9 (2 308 527.1 to 2 689 861.1) | 2 391 258.0 (2 184 254.1 to 2 631 699.0) | -2.3 (-5.9 to 0.9) | -4.0 (-6.0 to -2.1)* | 48 407.8 (45 385.4 to 51 762.0) | 40 485.1 (37 556.0 to 43 679.3) | 33 641.0 (30 808.7 to 36 924.3) | -30.5 (-32.6 to -28.6)* | -16.9 (-18.6 to -15.3)* |
| Communicable, maternal, neonatal, and nutritional diseases | 1 114 176.6 (1 073 948.8 to 1 156 050.2) | 918 804.8 (885 242.1 to 959 452.3) | 667 823.7 (632 212.4 to 708 405.1) | -40.1 (-42.7 to -37.4)* | -27.3 (-29.8 to -24.9)* | 18 071.6 (17 386.0 to 18 790.5) | 13 801.1 (13 286.8 to 14 406.5) | 9 396.8 (8 894.5 to 9 956.2) | -48.0 (-50.1 to -45.8)* | -31.9 (-34.2 to -29.7)* |
| HIV/AIDS and tuberculosis | 84 184.5 (79 728.6 to 89 558.0) | 159 063.9 (152 851.5 to 165 139.0) | 101 133.3 (97 487.1 to 105 092.9) | 20.1 (13.0 to 26.7)* | -36.4 (-37.9 to -34.7)* | 17 888.1 (16 901.1 to 19 188.3) | 24 399.5 (23 458.8 to 25 315.5) | 13 559.2 (13 066.2 to 14 075.5) | -24.2 (-29.7 to -20.1)* | -44.5 (-45.7 to -43.0)* |
| Tuberculosis | 68 029.7 (64 153.4 to 73 066.3) | 56 881.5 (54 312.6 to 59 442.6) | 43 557.9 (41 529.0 to 45 716.5) | -36.0 (-41.3 to -32.4)* | -23.4 (-26.1 to -20.6)* | 14 801.1 (13 904.4 to 16 136.6) | 9 169.5 (8 747.7 to 9 567.7) | 5 931.1 (5 655.5 to 6 217.7) | -59.9 (-63.7 to -57.0)* | -35.3 (-37.6 to -32.9)* |
| Drug-susceptible tuberculosis | 67 560.5 (63 730.5 to 72 611.6) | 51 760.2 (49 194.0 to 54 289.2) | 39 869.8 (38 054.8 to 41 916.2) | -41.0 (-45.6 to -37.5)* | -23.0 (-25.7 to -20.0)* | 14 699.5 (13 807.7 to 16 036.6) | 8 341.1 (7 916.6 to 8 736.6) | 5 430.0 (5 178.8 to 5 711.1) | -63.0 (-66.4 to -60.8)* | -34.9 (-37.2 to -32.5)* |
| Multidrug-resistant tuberculosis without extensive drug resistance | 469.2 (378.3 to 578.8) | 4 886.9 (4 122.2 to 5 829.2) | 3 319.4 (2 787.6 to 3 910.3) | 607.5 (511.8 to 717.0)* | -32.1 (-38.5 to -24.8)* | 10.6 (8.5 to 13.1) | 78.7 (66.3 to 93.9) | 45.1 (37.9 to 53.2) | 327.6 (267.7 to 394.1)* | -42.7 (-48.1 to -36.5)* |
| Extensively drug-resistant tuberculosis | .. | 234.5 (194.6 to 279.1) | 368.8 (301.1 to 444.5) | .. | 57.3 (36.1 to 82.1)* | .. | 3.8 (3.1 to 4.5) | 5.0 (4.1 to 6.0) | .. | 32.5 (14.8 to 53.2)* |
| Latent tuberculosis infection | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| HIV/AIDS | 16 154.8 (14 497.1 to 18 106.5) | 102 182.3 (96 751.1 to 107 544.2) | 57 575.4 (54 618.5 to 60 967.9) | 256.4 (220.1 to 293.5)* | -43.6 (-45.4 to -41.6)* | 3 081.1 (2 762.2 to 3 456.6) | 15 229.9 (14 436.6 to 16 016.3) | 7 621.1 (7 236.6 to 8 062.2) | 147.4 (121.5 to 173.2)* | -50.0 (-51.5 to -48.2)* |
| Drug-susceptible HIV/AIDS-Tuberculosis | 4 668.5 (3 624.4 to 5 760.2) | 24 070.5 (16 708.0 to 31 379.1) | 11 724.0 (8 154.4 to 15 522.4) | 151.1 (116.8 to 191.3)* | -51.3 (-54.0 to -48.6)* | 88.2 (68.0 to 109.1) | 3 599.7 (2 499.9 to 4 688.7) | 1 555.5 (1 088.2 to 2 059.9) | 76.2 (52.6 to 103.8)* | -56.8 (-59.2 to -54.4)* |
| Multidrug-resistant HIV/AIDS-Tuberculosis without extensive drug resistance | 25.9 (16.0 to 40.6) | 2 051.8 (1 282.8 to 3 070.2) | 979.2 (597.7 to 1 481.6) | 3 673.4 (2 732.2 to 4 952.8)* | -52.3 (-61.7 to -41.2)* | 0.5 (0.3 to 0.8) | 30.7 (19.2 to 45.8) | 13.0 (7.9 to 19.7) | 2 486.4 (1 853.0 to 3 358.6)* | -57.6 (-66.1 to -47.8)* |
| Extensively drug-resistant HIV/AIDS-Tuberculosis | .. | 39.9 (24.8 to 61.1) | 57.3 (34.5 to 89.4) | .. | 43.5 (25.5 to 65.4)* | .. | 0.6 (0.4 to 0.9) | 0.8 (0.5 to 1.2) | .. | 26.8 (10.8 to 46.4)* |
| HIV/AIDS resulting in other diseases | 11 460.3 (9 938.9 to 13 435.6) | 76 020.0 (67 021.8 to 86 026.2) | 44 814.9 (39 932.9 to 50 112.4) | 291.1 (245.9 to 337.5)* | -41.0 (-43.6 to -38.2)* | 2 199.3 (1 899.8 to 2 574.4) | 11 319.9 (9 999.9 to 12 809.9) | 5 929.9 (5 286.6 to 6 630.0) | 170.3 (138.2 to 203.1)* | -47.6 (-49.9 to -45.1)* |
| Diarrhoea, lower respiratory, and other common infectious diseases | 557 388.0 (522 551.7 to 600 325.4) | 337 062.8 (317 957.5 to 359 176.2) | 229 961.4 (213 682.3 to 247 975.2) | -58.7 (-61.9 to -55.2)* | -31.8 (-35.3 to -27.8)* | 89 512.2 (83 780.0 to 96 017.7) | 51 527.7 (48 588.8 to 55 149.9) | 32 756.6 (30 517.7 to 35 318.8) | -63.4 (-66.0 to -60.7)* | -36.4 (-39.7 to -33.0)* |
| Diarrhoeal diseases | 175 168.6 (150 592.6 to 201 351.3) | 113 944.8 (99 183.9 to 135 659.8) | 74 414.6 (63 402.0 to 93 414.9) | -57.5 (-62.8 to -50.1)* | -34.7 (-41.0 to -28.1)* | 29 142.2 (24 829.9 to 34 692.2) | 17 685.5 (15 266.0 to 21 367.0) | 10 631.1 (9 075.5 to 13 323.3) | -63.5 (-67.6 to -58.3)* | -39.9 (-45.2 to -34.3)* |
| Intestinal infectious diseases | 15 662.6 (8 797.4 to 25 360.4) | 12 822.7 (7 207.6 to 20 879.4) | 10 601.7 (6 041.1 to 17 309.3) | -32.3 (-43.5 to -21.8)* | -17.3 (-25.1 to -10.8)* | 2 499.7 (1 406.6 to 4 048.8) | 1 846.6 (1 038.8 to 3 004.4) | 1 443.3 (823.3 to 2 352.2) | -42.2 (-51.5 to -33.7)* | -21.9 (-29.4 to -15.6)* |
| Typhoid fever | 13 362.8 (7 235.9 to 22 248.3) | 10 793.8 (5 876.4 to 17 717.0) | 8 843.0 (4 901.5 to 14 436.1) | -33.8 (-44.0 to -23.6)* | -18.1 (-25.7 to -12.1)* | 2 125.5 (1 150.0 to 3 539.9) | 1 553.3 (848.8 to 2 540.0) | 1 204.4 (666.6 to 1 968.8) | -43.3 (-51.9 to -35.1)* | -22.4 (-29.7 to -16.6)* |
| Paratyphoid fever | 1 867.3 (850.2 to 3 711.1) | 1 773.6 (826.8 to 3 439.6) | 1 607.0 (759.0 to 3 109.8) | -13.9 (-27.1 to -1.5)* | -9.4 (-17.9 to -1.8)* | 306.6 (139.9 to 604.4) | 256.6 (119.9 to 496.6) | 217.7 (102.0 to 420.0) | -29.0 (-39.4 to -19.2)* | -15.1 (-22.9 to -8.0)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|--|--|--|---|---------------------------------------|---|--|--|---|---|--|
| | 1990 | 2006 | 2016 | Percentage change, 1990-2016 | Percentage change, 2006-16 | 1990 | 2006 | 2016 | Percentage change, 1990-2016 | Percentage change, 2006-16 |
| (Continued from previous page) | | | | | | | | | | |
| Other intestinal infectious diseases | 432.5 (107.2 to 1290.8) | 255.4 (58.5 to 753.4) | 151.7 (42.1 to 412.1) | -64.9 (-90.6 to 42.1) | -40.6 (-84.7 to 126.3) | 6.7 (1.7 to 19.5) | 3.8 (0.9 to 11.2) | 2.2 (0.6 to 5.9) | -67.5 (-91.3 to 31.5) | -42.9 (-85.2 to 114.5) |
| Lower respiratory infections | 202365.5 (182794.4 to 220607.6) | 131015.4 (121489.8 to 139228.6) | 91844.6 (84674.4 to 98252.6) | -54.6 (-58.7 to -49.4)* | -29.9 (-34.4 to -25.3)* | 3237.1 (2942.1 to 3515.4) | 2022.6 (1879.0 to 2147.6) | 1326.7 (1221.8 to 1419.7) | -59.0 (-62.5 to -54.6)* | -34.4 (-38.6 to -30.2)* |
| Upper respiratory infections | 4868.6 (3012.8 to 7444.9) | 5551.2 (3380.7 to 8488.2) | 5991.2 (3621.1 to 9193.8) | 23.1 (19.0 to 26.3)* | 7.9 (6.2 to 9.4)* | 88.2 (54.9 to 134.0) | 83.0 (50.8 to 126.9) | 81.0 (49.0 to 124.0) | -8.3 (-10.5 to -6.9)* | -2.5 (-3.8 to -1.5)* |
| Otitis media | 3111.7 (2057.2 to 4485.0) | 3171.4 (2005.0 to 4675.3) | 3187.5 (1993.2 to 4716.5) | 2.4 (-3.8 to 7.0) | 0.5 (-2.1 to 3.0) | 53.4 (35.3 to 76.9) | 46.7 (29.6 to 68.9) | 43.3 (27.1 to 64.2) | -18.9 (-23.7 to -15.1)* | -7.3 (-9.8 to -4.9)* |
| Meningitis | 30239.3 (23939.3 to 34552.6) | 24957.4 (21655.0 to 28764.2) | 21865.9 (18204.6 to 28280.5) | -27.7 (-41.7 to 3.1) | -12.4 (-23.9 to 7.8) | 481.8 (385.6 to 549.0) | 369.9 (321.3 to 426.1) | 306.1 (254.0 to 398.0) | -36.5 (-48.5 to -9.7)* | -17.2 (-28.1 to 2.1) |
| Pneumococcal meningitis | 2187.5 (1808.2 to 2576.0) | 1940.0 (1649.9 to 2287.4) | 1902.8 (1569.5 to 2382.2) | -13.0 (-26.3 to 10.0) | -1.9 (-11.5 to 12.0) | 37.2 (31.0 to 43.4) | 29.1 (24.8 to 34.2) | 26.2 (21.6 to 32.7) | -29.6 (-39.6 to -11.3)* | -9.9 (-18.7 to 3.2) |
| Haemophilus influenzae type B meningitis | 3330.3 (2606.2 to 3982.2) | 2725.7 (2276.3 to 3165.4) | 2426.0 (1967.2 to 3212.2) | -27.1 (-41.5 to 3.7) | -11.0 (-23.9 to 9.3) | 52.4 (41.5 to 62.0) | 40.3 (33.7 to 46.8) | 34.1 (27.6 to 45.1) | -34.8 (-47.4 to -7.4)* | -15.3 (-27.6 to 3.9) |
| Meningococcal meningitis | 14191.0 (11094.1 to 16492.1) | 11548.6 (9913.5 to 13418.2) | 8327.1 (6806.4 to 10911.9) | -41.3 (-53.1 to -16.3)* | -27.9 (-37.7 to -12.2)* | 224.2 (177.0 to 259.3) | 170.8 (146.7 to 199.1) | 116.6 (95.2 to 152.9) | -48.0 (-58.4 to -26.0)* | -31.7 (-41.1 to -16.8)* |
| Other meningitis | 10530.5 (8030.6 to 12434.2) | 8743.1 (7350.6 to 10148.8) | 9210.0 (7559.7 to 12250.5) | -12.5 (-30.3 to 28.9) | 5.3 (-9.4 to 33.0) | 168.1 (129.1 to 197.5) | 129.7 (109.2 to 150.4) | 129.2 (105.5 to 173.3) | -23.1 (-38.2 to 12.4) | -0.4 (-14.4 to 26.1) |
| Encephalitis | 7918.4 (5206.9 to 10751.2) | 7380.9 (6422.5 to 9033.9) | 6704.1 (5469.3 to 8574.2) | -15.3 (-44.0 to 40.6) | -9.2 (-24.4 to 10.8) | 135.5 (91.8 to 180.2) | 111.5 (97.1 to 136.5) | 92.7 (75.7 to 118.4) | -31.6 (-53.6 to 10.0) | -16.9 (-30.8 to 1.2) |
| Diphtheria | 842.7 (611.3 to 1167.2) | 263.8 (183.1 to 374.7) | 86.9 (62.5 to 123.4) | -89.7 (-93.2 to -84.0)* | -67.0 (-78.6 to -47.7)* | 12.9 (9.4 to 17.9) | 3.9 (2.7 to 5.6) | 1.2 (0.9 to 1.8) | -90.5 (-93.7 to -85.3)* | -68.6 (-79.8 to -49.7)* |
| Whooping cough | 14651.2 (6598.0 to 28290.2) | 9778.0 (4727.9 to 17764.8) | 6249.9 (3360.7 to 10754.7) | -57.3 (-77.1 to -19.2)* | -36.1 (-63.1 to 17.1) | 219.5 (98.9 to 424.0) | 144.6 (69.9 to 262.6) | 89.4 (48.1 to 153.9) | -59.3 (-78.1 to -22.8)* | -38.1 (-64.3 to 13.3) |
| Tetanus | 24893.6 (14235.3 to 33445.8) | 6340.9 (3695.4 to 7940.5) | 2366.6 (1446.0 to 3062.9) | -90.5 (-92.7 to -87.7)* | -62.7 (-68.8 to -55.4)* | 385.3 (222.8 to 516.6) | 93.3 (54.4 to 116.8) | 33.6 (20.3 to 43.4) | -91.3 (-93.2 to -88.9)* | -64.0 (-70.0 to -57.1)* |
| Measles | 76350.8 (31267.6 to 147358.9) | 20794.3 (8237.6 to 43871.3) | 5724.8 (2148.6 to 12257.6) | -92.5 (-94.4 to -90.5)* | -72.5 (-76.9 to -67.7)* | 1150.7 (471.2 to 2220.4) | 307.9 (122.0 to 649.2) | 81.3 (30.5 to 174.2) | -92.9 (-94.7 to -91.0)* | -73.6 (-77.8 to -69.0)* |
| Varicella and herpes zoster | 1314.9 (1138.7 to 1509.6) | 1042.2 (909.6 to 1205.0) | 923.5 (779.5 to 1098.7) | -29.8 (-40.3 to -18.8)* | -11.4 (-20.6 to -2.8)* | 22.8 (19.8 to 25.9) | 16.2 (14.1 to 18.9) | 13.0 (11.0 to 15.4) | -42.8 (-50.3 to -35.4)* | -19.7 (-27.6 to -12.2)* |
| Neglected tropical diseases and malaria | 87294.8 (71756.4 to 103455.7) | 99229.2 (85820.3 to 113978.1) | 74995.1 (63114.8 to 86650.7) | -14.1 (-31.0 to 6.2) | -24.4 (-37.6 to -8.6)* | 1423.8 (1183.4 to 1676.5) | 1478.5 (1280.4 to 1696.3) | 1050.5 (882.7 to 1217.9) | -26.2 (-40.4 to -9.3)* | -28.9 (-41.5 to -14.0)* |
| Malaria | 60389.3 (46548.2 to 74912.5) | 77253.7 (64810.3 to 91256.8) | 56201.2 (45785.6 to 67880.8) | -6.9 (-30.5 to 26.5) | -27.2 (-43.3 to -6.7)* | 931.0 (722.1 to 1150.5) | 1147.0 (963.0 to 1354.6) | 794.7 (646.5 to 962.2) | -14.6 (-36.1 to 15.2) | -30.7 (-46.2 to -11.1)* |
| Chagas disease | 309.8 (286.3 to 334.8) | 226.1 (204.9 to 251.3) | 219.0 (194.6 to 250.7) | -29.3 (-34.4 to -23.7)* | -3.1 (-8.2 to 2.7) | 7.7 (7.1 to 8.3) | 4.0 (3.7 to 4.5) | 3.1 (2.8 to 3.6) | -59.3 (-62.2 to -56.0)* | -22.6 (-26.7 to -17.8)* |
| Leishmaniasis | 2531.5 (1470.2 to 4203.0) | 1897.2 (1151.9 to 3064.4) | 981.0 (658.3 to 1480.6) | -61.2 (-67.2 to -52.0)* | -48.3 (-54.0 to -39.6)* | 45.7 (27.0 to 75.4) | 28.4 (17.2 to 45.7) | 13.4 (9.0 to 20.3) | -70.6 (-74.9 to -64.1)* | -52.6 (-57.7 to -44.9)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|---|--------------------------------|------------------------------|------------------------------|------------------------------|----------------------------|--|------------------------|------------------------|------------------------------|----------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 |
| (Continued from previous page) | | | | | | | | | | |
| Visceral leishmaniasis | 2406.1 (1350.5 to 4080.8) | 1684.8 (943.7 to 2878.0) | 707.9 (400.1 to 1206.2) | -70.6 (-74.4 to -66.4)* | -58.0 (-61.9 to -53.9)* | 43.1 (24.4 to 72.6) | 25.1 (14.1 to 42.9) | 9.8 (5.5 to 16.6) | -77.4 (-79.9 to -74.6)* | -61.1 (-64.7 to -57.2)* |
| Cutaneous and mucocutaneous leishmaniasis | 125.3 (67.7 to 217.2) | 212.4 (131.6 to 329.4) | 273.1 (177.2 to 398.9) | 117.9 (67.5 to 215.8)* | 28.6 (16.8 to 42.9)* | 2.6 (1.4 to 4.4) | 3.3 (2.0 to 5.0) | 3.7 (2.4 to 5.4) | 43.5 (11.8 to 103.5)* | 12.5 (1.7 to 26.1)* |
| African trypanosomiasis | 1046.8 (559.1 to 1711.5) | 539.0 (288.0 to 876.8) | 128.4 (64.7 to 215.0) | -87.7 (-91.3 to -82.2)* | -76.2 (-83.2 to -65.9)* | 19.2 (10.3 to 31.6) | 7.8 (4.2 to 12.8) | 1.7 (0.9 to 2.9) | -91.1 (-93.6 to -87.2)* | -78.2 (-84.6 to -68.8)* |
| Schistosomiasis | 2096.8 (1340.2 to 3410.1) | 2464.8 (1447.7 to 4194.7) | 1863.6 (1122.0 to 3175.2) | -11.1 (-17.5 to -6.7)* | -24.4 (-26.2 to -21.9)* | 42.8 (27.5 to 69.2) | 37.5 (22.2 to 63.5) | 24.9 (15.0 to 42.3) | -41.9 (-46.3 to -38.9)* | -33.7 (-35.4 to -31.6)* |
| Cysticercosis | 489.0 (363.4 to 621.4) | 500.8 (359.2 to 657.1) | 468.1 (322.8 to 625.8) | -4.3 (-13.5 to 3.9) | -6.5 (-12.4 to -0.9)* | 10.5 (7.8 to 13.4) | 8.0 (5.7 to 10.5) | 6.3 (4.4 to 8.4) | -40.0 (-45.2 to -35.0)* | -21.0 (-26.0 to -16.5)* |
| Cystic echinococcosis | 326.8 (237.4 to 449.1) | 226.2 (161.3 to 313.8) | 136.5 (95.3 to 193.7) | -58.2 (-68.4 to -44.3)* | -39.6 (-53.9 to -16.3)* | 6.3 (4.5 to 8.7) | 3.5 (2.5 to 4.8) | 1.8 (1.3 to 2.6) | -70.6 (-77.7 to -61.1)* | -46.6 (-59.1 to -26.5)* |
| Lymphatic filariasis | 1595.7 (733.4 to 2983.5) | 1897.7 (873.9 to 3542.8) | 1189.0 (587.7 to 2114.9) | -25.5 (-41.4 to -9.3)* | -37.4 (-52.4 to -26.0)* | 32.5 (14.9 to 60.7) | 28.9 (13.3 to 54.0) | 15.8 (7.8 to 28.1) | -51.5 (-61.8 to -41.2)* | -45.3 (-58.4 to -35.5)* |
| Onchocerciasis | 1420.4 (777.0 to 2254.9) | 1266.4 (705.0 to 2003.3) | 962.5 (452.3 to 1672.1) | -32.2 (-47.6 to -16.8)* | -24.0 (-41.6 to -6.5)* | 28.4 (16.0 to 45.4) | 19.1 (11.0 to 30.1) | 12.9 (6.1 to 22.4) | -54.6 (-65.7 to -43.4)* | -32.6 (-48.5 to -17.5)* |
| Trachoma | 231.1 (156.9 to 324.3) | 246.5 (166.2 to 348.4) | 245.2 (162.4 to 353.6) | 6.1 (-2.6 to 14.6) | -0.5 (-6.5 to 5.2) | 6.7 (4.5 to 9.4) | 4.9 (3.3 to 6.9) | 3.7 (2.5 to 5.3) | -44.3 (-49.0 to -39.6)* | -23.8 (-28.5 to -19.3)* |
| Dengue | 822.8 (308.1 to 1364.0) | 1798.2 (789.6 to 2494.8) | 2956.9 (1359.2 to 4146.9) | 259.4 (104.2 to 683.3)* | 64.4 (36.2 to 115.9)* | 13.2 (5.2 to 23.2) | 26.7 (11.7 to 37.1) | 40.2 (18.6 to 56.3) | 189.0 (65.5 to 523.4)* | 50.5 (24.7 to 97.7)* |
| Yellow fever | 784.5 (170.6 to 2314.6) | 424.5 (89.8 to 1247.2) | 374.0 (80.8 to 1075.1) | -52.3 (-61.3 to -41.2)* | -11.9 (-26.9 to 7.4) | 13.2 (2.9 to 39.0) | 6.1 (1.3 to 18.0) | 5.0 (1.1 to 14.5) | -61.8 (-68.5 to -52.9)* | -17.1 (-31.3 to 1.3) |
| Rabies | 2979.4 (1867.4 to 4076.4) | 1451.6 (867.1 to 1868.0) | 744.2 (383.8 to 1106.3) | -75.0 (-82.7 to -61.4)* | -48.7 (-58.9 to -34.1)* | 51.3 (32.6 to 71.3) | 21.5 (12.9 to 27.5) | 10.1 (5.2 to 15.1) | -80.3 (-86.5 to -69.9)* | -52.9 (-62.4 to -39.5)* |
| Intestinal nematode infections | 7460.9 (4726.6 to 11 584.0) | 4083.3 (2617.1 to 6154.1) | 3331.2 (2076.2 to 5158.6) | -55.4 (-57.9 to -52.6)* | -18.4 (-22.9 to -14.3)* | 132.8 (83.2 to 206.9) | 60.8 (39.0 to 91.6) | 45.0 (28.2 to 69.6) | -66.1 (-68.0 to -64.0)* | -25.9 (-30.0 to -22.2)* |
| Ascariasis | 4634.7 (2996.9 to 7119.9) | 1902.0 (1325.4 to 2758.9) | 1308.8 (883.2 to 1942.4) | -71.8 (-74.6 to -68.4)* | -31.2 (-37.6 to -24.8)* | 80.3 (51.4 to 124.0) | 28.3 (19.8 to 41.1) | 17.9 (12.1 to 26.4) | -77.8 (-79.9 to -74.9)* | -37.0 (-42.8 to -31.0)* |
| Trichuriasis | 671.4 (364.9 to 1142.9) | 421.5 (233.0 to 717.2) | 337.0 (186.2 to 573.6) | -49.8 (-55.2 to -44.3)* | -20.0 (-27.9 to -11.6)* | 12.5 (6.8 to 21.2) | 6.3 (3.5 to 10.7) | 4.5 (2.5 to 7.7) | -63.7 (-67.8 to -59.7)* | -27.8 (-34.9 to -20.2)* |
| Hookworm disease | 2154.8 (1278.7 to 3371.1) | 1759.8 (1058.9 to 2739.1) | 1685.4 (1001.5 to 2648.9) | -21.8 (-26.7 to -16.6)* | -4.2 (-9.5 to 1.3) | 40.0 (23.7 to 62.8) | 26.2 (15.7 to 40.7) | 22.6 (13.5 to 35.5) | -43.4 (-47.2 to -39.6)* | -13.5 (-18.2 to -8.5)* |
| Food-borne trematodiasis | 1425.0 (591.7 to 2937.7) | 1659.6 (832.5 to 3083.5) | 1771.2 (923.9 to 3158.4) | 24.3 (-3.9 to 71.7) | 6.7 (1.3 to 15.7)* | 27.9 (12.0 to 56.7) | 25.4 (12.9 to 46.8) | 23.7 (12.2 to 42.0) | -15.3 (-34.0 to 13.1) | -7.0 (-11.4 to 0.5) |
| Leprosy | 23.0 (15.5 to 32.3) | 31.3 (21.3 to 44.0) | 31.6 (21.4 to 44.0) | 37.5 (34.0 to 40.9)* | 1.1 (-1.3 to 3.6) | 0.6 (0.4 to 0.8) | 0.5 (0.4 to 0.8) | 0.4 (0.3 to 0.6) | -20.7 (-22.7 to -18.8)* | -18.1 (-20.0 to -16.2)* |
| Ebola virus disease | .. | .. | 0.3 (0.2 to 1.1) | .. | .. | .. | .. | .. | .. | .. |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|---|---|---|---|------------------------------------|------------------------------------|--|--------------------------------------|--------------------------------------|------------------------------------|------------------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 |
| (Continued from previous page) | | | | | | | | | | |
| Zika virus disease | .. | .. | 5.1 (3.4–8.0) | .. | .. | .. | .. | 0.1 (0.0–0.1) | .. | .. |
| Guinea worm disease | 50.7 (35.3 to 69.2) | 0.2 (0.1 to 0.3) | .. | -100.0 (-100.0 to -100.0)* | -99.5 (-99.7 to -99.4)* | 1.1 (0.7 to 1.4) | .. | .. | -100.0 (-100.0 to -100.0)* | -99.6 (-99.7 to -99.5)* |
| Other neglected tropical diseases | 3311.2 (2409.7 to 4421.6) | 3262.2 (2470.4 to 4130.1) | 3386.0 (2569.6 to 4260.7) | 2.3 (-22.0 to 31.6) | 3.8 (-14.9 to 23.9) | 52.2 (38.2 to 69.0) | 48.3 (36.6 to 61.2) | 47.4 (35.9 to 59.7) | -9.1 (-29.8 to 15.8) | -1.9 (-19.5 to 17.3) |
| Maternal disorders | 21 597.1 (20 063.6 to 22 834.1) | 18 093.0 (16 785.8 to 19 171.8) | 13 763.0 (12 668.6 to 15 064.0) | -36.3 (-41.3 to -31.0)* | -23.9 (-29.3 to -17.4)* | 388.6 (361.4 to 411.5) | 257.2 (238.8 to 272.5) | 179.0 (164.7 to 195.8) | -53.9 (-57.6 to -50.2)* | -30.4 (-35.4 to -24.4)* |
| Maternal haemorrhage | 6945.7 (5770.8 to 8257.7) | 5416.4 (4627.6 to 6301.7) | 4078.3 (3311.7 to 5035.2) | -41.3 (-47.4 to -34.9)* | -24.7 (-32.5 to -16.0)* | 124.4 (103.1 to 148.1) | 77.0 (65.8 to 89.7) | 53.0 (43.1 to 65.4) | -57.4 (-61.7 to -53.0)* | -31.2 (-38.2 to -23.3)* |
| Maternal sepsis and other maternal infections | 2102.5 (1614.0 to 2721.4) | 1562.3 (1223.7 to 1984.3) | 1139.4 (833.1 to 1525.2) | -45.8 (-51.9 to -39.7)* | -27.1 (-35.7 to -18.1)* | 37.6 (29.0 to 48.2) | 22.1 (17.3 to 28.0) | 14.8 (10.9 to 19.8) | -60.6 (-64.8 to -56.1)* | -33.0 (-40.9 to -24.8)* |
| Maternal hypertensive disorders | 2469.0 (1945.4 to 3125.5) | 2478.1 (1983.1 to 3052.0) | 1996.8 (1569.9 to 2483.1) | -19.1 (-26.0 to -11.2)* | -19.4 (-27.3 to -11.1)* | 44.0 (34.8 to 55.2) | 35.1 (28.2 to 43.2) | 26.0 (20.5 to 32.3) | -40.9 (-45.8 to -35.4)* | -25.8 (-33.1 to -17.9)* |
| Maternal obstructed labour and uterine rupture | 1440.8 (1086.2 to 1865.8) | 1240.4 (944.6 to 1582.8) | 969.0 (716.6 to 1270.1) | -32.8 (-37.4 to -27.7)* | -21.9 (-27.4 to -15.9)* | 27.0 (20.4 to 35.0) | 17.9 (13.6 to 22.9) | 12.6 (9.3 to 16.5) | -53.3 (-56.5 to -50.0)* | -29.5 (-34.5 to -24.1)* |
| Maternal abortion, miscarriage, and ectopic pregnancy | 1743.4 (1338.9 to 2235.5) | 1469.0 (1150.0 to 1845.5) | 1145.1 (855.9 to 1541.6) | -34.3 (-40.9 to -26.8)* | -22.1 (-30.6 to -12.6)* | 31.9 (24.6 to 40.6) | 20.9 (16.4 to 26.3) | 14.9 (11.1 to 19.9) | -53.3 (-57.8 to -48.0)* | -28.8 (-36.6 to -20.4)* |
| Indirect maternal deaths | 2611.6 (1943.4 to 3385.8) | 2577.2 (1949.1 to 3277.1) | 1987.9 (1463.8 to 2619.8) | -23.9 (-30.7 to -16.4)* | -22.9 (-30.1 to -14.8)* | 47.0 (35.2 to 60.4) | 36.7 (27.7 to 46.6) | 25.8 (19.1 to 34.0) | -45.0 (-49.7 to -39.9)* | -29.6 (-36.2 to -22.1)* |
| Late maternal deaths | 374.2 (244.4 to 552.4) | 298.2 (178.6 to 480.8) | 228.5 (134.6 to 370.9) | -39.0 (-50.1 to -28.8)* | -23.4 (-29.9 to -16.2)* | 6.7 (4.4 to 9.8) | 4.2 (2.5 to 6.8) | 3.0 (1.8 to 4.8) | -55.8 (-63.7 to -49.2)* | -29.9 (-35.5 to -23.5)* |
| Maternal deaths aggravated by HIV/AIDS | 36.9 (21.5 to 52.5) | 128.5 (81.8 to 169.0) | 105.4 (66.7 to 142.9) | 185.9 (142.7 to 243.7)* | -17.9 (-27.8 to -6.0)* | 0.7 (0.4 to 0.9) | 1.9 (1.2 to 2.4) | 1.4 (0.9 to 1.9) | 104.5 (73.6 to 146.1)* | -26.1 (-35.0 to -15.3)* |
| Other maternal disorders | 3872.9 (3015.3 to 4777.9) | 2923.1 (2365.7 to 3539.5) | 2112.5 (1628.9 to 2673.2) | -45.5 (-50.6 to -39.9)* | -27.7 (-34.7 to -19.7)* | 69.3 (54.4 to 84.9) | 41.5 (33.6 to 50.0) | 27.5 (21.2 to 34.8) | -60.4 (-64.0 to -56.5)* | -33.7 (-40.1 to -26.4)* |
| Neonatal disorders | 261 357.2 (248 875.2 to 282 758.2) | 211 984.8 (203 477.1 to 221 317.2) | 163 569.7 (154 643.2 to 172 756.7) | -37.4 (-42.7 to -32.7)* | -22.8 (-26.8 to -18.9)* | 3818.4 (3635.3 to 4130.2) | 3073.5 (2949.7 to 3208.5) | 2364.2 (2237.5 to 2493.4) | -38.1 (-43.2 to -33.5)* | -23.1 (-27.0 to -19.2)* |
| Neonatal preterm birth complications | 112 767.2 (105 488.9 to 124 122.5) | 81 159.7 (76 378.0 to 89 409.0) | 62 031.6 (57 062.9 to 67 530.0) | -45.0 (-50.3 to -39.5)* | -23.6 (-29.9 to -17.4)* | 1652.8 (1546.2 to 1819.4) | 1176.6 (1107.3 to 1296.0) | 892.7 (822.1 to 970.7) | -46.0 (-51.2 to -40.7)* | -24.1 (-30.4 to -18.1)* |
| Neonatal encephalopathy due to birth asphyxia, and trauma | 68 251.9 (61 749.3 to 76 860.5) | 60 334.8 (55 822.8 to 65 138.9) | 47 031.7 (41 794.2 to 51 919.1) | -31.1 (-41.3 to -20.8)* | -22.1 (-29.3 to -14.6)* | 993.0 (897.9 to 1117.7) | 874.2 (808.8 to 943.9) | 682.2 (606.6 to 751.3) | -31.3 (-41.4 to -21.4)* | -22.0 (-29.1 to -14.6)* |
| Neonatal sepsis and other neonatal infections | 24 573.0 (18 972.5 to 31 156.6) | 25 874.0 (21 266.6 to 32 360.3) | 23 675.8 (20 056.0 to 30 684.5) | -3.6 (-22.7 to 20.9) | -8.5 (-19.1 to 4.0) | 360.8 (278.4 to 460.0) | 375.7 (308.9 to 469.6) | 341.7 (291.0 to 444.5) | -5.3 (-23.7 to 18.6) | -9.0 (-19.5 to 3.2) |
| Haemolytic disease and other neonatal jaundice | 12 277.3 (10 225.3 to 15 113.1) | 7996.0 (7149.5 to 9006.9) | 4912.8 (4310.5 to 5605.8) | -60.0 (-69.1 to -50.4)* | -38.6 (-46.2 to -30.3)* | 179.9 (149.9 to 221.4) | 116.1 (103.8 to 130.8) | 70.7 (62.0 to 80.7) | -60.7 (-69.6 to -51.4)* | -39.1 (-46.6 to -30.8)* |
| Other neonatal disorders | 43 487.8 (37 402.4 to 50 951.9) | 36 620.3 (33 314.2 to 40 250.5) | 25 917.7 (23 440.4 to 28 405.4) | -40.4 (-50.6 to -28.5)* | -29.2 (-36.1 to -21.0)* | 631.9 (543.4 to 740.9) | 530.9 (483.0 to 583.6) | 376.8 (340.8 to 413.1) | -40.4 (-50.6 to -28.4)* | -29.0 (-35.9 to -20.7)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|---|---|---|---|-----------------------------------|-----------------------------------|--|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 |
| (Continued from previous page) | | | | | | | | | | |
| Nutritional deficiencies | 69 823.3 (57 049.4 to 85 126.9) | 64 648.9 (51 489.2 to 80 980.6) | 60 936.1 (46 656.8 to 79 062.8) | -12.7 (-23.8 to -1.2)* | -5.7 (-12.3 to 0.1) | 1167.8 (949.2 to 1434.1) | 971.1 (775.1 to 1213.5) | 844.3 (649.4 to 1090.3) | -27.7 (-35.9 to -19.1)* | -13.1 (-18.9 to -7.7)* |
| Protein-energy malnutrition | 35 843.1 (29 589.9 to 41 569.5) | 26 417.6 (23 896.5 to 29 406.0) | 20 718.9 (18 009.6 to 24 194.8) | -42.2 (-51.8 to -26.7)* | -21.6 (-32.3 to -8.8)* | 561.0 (467.3 to 647.0) | 398.8 (361.7 to 443.4) | 296.7 (258.1 to 346.4) | -47.1 (-55.8 to -33.4)* | -25.6 (-35.7 to -13.6)* |
| Iodine deficiency | 4355.1 (2983.7 to 6048.6) | 3453.9 (2363.8 to 4790.9) | 3240.6 (2213.3 to 4488.7) | -25.6 (-28.2 to -22.9)* | -6.2 (-8.7 to -3.5)* | 84.5 (58.0 to 117.3) | 52.4 (35.9 to 72.6) | 43.5 (29.8 to 60.3) | -48.5 (-50.2 to -46.7)* | -16.9 (-19.2 to -14.5)* |
| Vitamin A deficiency | 188.5 (116.6 to 294.4) | 225.3 (139.7 to 348.3) | 252.4 (158.8 to 388.3) | 33.9 (27.4 to 41.0)* | 12.0 (8.7 to 15.5)* | 3.2 (2.0 to 4.9) | 3.4 (2.1 to 5.2) | 3.4 (2.2 to 5.3) | 7.0 (2.2 to 11.4)* | 2.6 (-0.3 to 5.6) |
| Iron-deficiency anaemia | 27 097.5 (18 018.7 to 38 749.1) | 32 422.9 (21 535.8 to 46 668.3) | 34 841.8 (23 085.2 to 49 693.9) | 28.6 (26.5 to 30.4)* | 7.5 (6.2 to 8.9)* | 475.7 (317.4 to 679.7) | 482.7 (320.9 to 693.0) | 474.1 (314.1 to 676.0) | -0.3 (-1.5 to 0.7) | -1.8 (-3.0 to -0.5)* |
| Other nutritional deficiencies | 2339.1 (1908.0 to 2766.9) | 2129.2 (1791.2 to 2366.5) | 1882.4 (1592.2 to 2158.7) | -19.5 (-32.9 to 5.7) | -11.6 (-20.4 to 0.7) | 43.3 (36.7 to 50.2) | 33.8 (28.4 to 37.4) | 26.4 (22.4 to 30.3) | -38.9 (-48.1 to -24.3)* | -21.7 (-29.1 to -11.6)* |
| Other communicable, maternal, neonatal, and nutritional diseases | 32 531.7 (25 112.3 to 42 021.4) | 28 722.2 (22 674.6 to 36 833.4) | 23 465.1 (18 762.9 to 29 098.8) | -27.9 (-36.2 to -18.6)* | -18.3 (-25.2 to -10.6)* | 533.7 (424.3 to 675.0) | 428.7 (340.1 to 547.4) | 328.0 (261.3 to 409.8) | -38.5 (-44.8 to -31.3)* | -23.5 (-29.8 to -16.4)* |
| Sexually transmitted diseases excluding HIV | 16 447.6 (9960.1 to 25 590.4) | 15 145.5 (9416.5 to 22 818.4) | 12 016.0 (7764.9 to 17 118.7) | -26.9 (-37.6 to -13.9)* | -20.7 (-30.2 to -9.4)* | 254.0 (157.8 to 389.9) | 222.6 (138.7 to 335.0) | 169.4 (108.1 to 242.8) | -33.3 (-42.1 to -22.6)* | -23.9 (-32.8 to -13.4)* |
| Syphilis | 14 613.8 (8275.0 to 23 755.0) | 12 825.3 (7321.4 to 20 493.5) | 9415.8 (5467.7 to 14 602.8) | -35.6 (-45.5 to -23.1)* | -26.6 (-35.8 to -14.9)* | 217.9 (124.4 to 352.0) | 188.3 (107.7 to 300.6) | 135.5 (78.6 to 210.2) | -37.8 (-47.4 to -25.8)* | -28.1 (-37.1 to -16.8)* |
| Chlamydial infection | 425.2 (287.5 to 639.0) | 519.6 (341.4 to 782.1) | 562.4 (370.3 to 851.1) | 32.3 (25.6 to 39.6)* | 8.2 (5.9 to 10.4)* | 8.0 (5.4 to 11.8) | 7.5 (5.0 to 11.3) | 7.3 (4.8 to 11.0) | -8.6 (-14.2 to -3.4)* | -3.3 (-5.6 to -1.4)* |
| Gonococcal infection | 465.0 (334.8 to 630.0) | 582.3 (412.5 to 824.5) | 675.2 (467.7 to 974.9) | 45.2 (26.9 to 64.5)* | 16.0 (-30.2 to 21.8)* | 9.2 (6.8 to 12.4) | 8.6 (6.1 to 12.1) | 8.8 (6.1 to 12.7) | -4.3 (-17.1 to 9.2) | 2.7 (-2.7 to 7.8) |
| Trichomoniasis | 125.7 (48.3 to 265.4) | 170.9 (65.2 to 362.0) | 198.2 (75.9 to 420.8) | 57.7 (55.1 to 60.5)* | 15.9 (14.8 to 17.1)* | 2.5 (0.9 to 5.2) | 2.5 (1.0 to 5.4) | 2.6 (1.0 to 5.5) | 4.7 (3.6 to 5.8)* | 1.8 (0.9 to 2.7)* |
| Genital herpes | 132.8 (43.0 to 302.2) | 187.9 (61.0 to 428.1) | 221.4 (71.2 to 507.1) | 66.8 (61.5 to 70.0)* | 17.8 (15.5 to 19.7)* | 2.9 (0.9 to 6.6) | 3.0 (1.0 to 6.8) | 3.0 (1.0 to 6.8) | 2.9 (1.3 to 4.8)* | -0.2 (-1.6 to 1.5) |
| Other sexually transmitted diseases | 685.1 (474.3 to 964.9) | 859.5 (589.4 to 1221.9) | 943.0 (643.7 to 1349.4) | 37.6 (30.6 to 44.7)* | 9.7 (7.4 to 12.2)* | 13.5 (9.4 to 18.9) | 12.6 (8.7 to 17.9) | 12.3 (8.4 to 17.6) | -8.7 (-13.7 to -4.0)* | -2.6 (-4.8 to -0.4)* |
| Hepatitis | 9017.2 (8255.1 to 9723.9) | 7718.6 (7259.2 to 8169.0) | 5777.8 (5492.2 to 6078.9) | -35.9 (-41.5 to -30.1)* | -25.1 (-29.2 to -21.0)* | 163.7 (151.1 to 175.3) | 117.8 (111.0 to 124.6) | 78.6 (74.8 to 82.7) | -52.0 (-55.9 to -48.3)* | -33.3 (-36.8 to -29.6)* |
| Acute hepatitis A | 1271.9 (1017.1 to 1540.7) | 849.3 (677.0 to 1036.3) | 450.7 (364.0 to 544.8) | -64.6 (-72.1 to -55.0)* | -46.9 (-58.9 to -32.0)* | 175.5 (15.7 to 23.5) | 12.6 (10.1 to 15.3) | 6.4 (5.2 to 7.8) | -67.1 (-74.0 to -58.4)* | -49.0 (-60.6 to -34.6)* |
| Hepatitis B | 4656.5 (4209.1 to 5096.9) | 4373.2 (4013.0 to 4741.4) | 3823.8 (3543.7 to 4119.0) | -17.9 (-24.6 to -10.7)* | -12.6 (-17.9 to -7.0)* | 94.2 (86.2 to 102.2) | 68.7 (63.5 to 74.3) | 51.6 (47.9 to 55.6) | -45.2 (-49.2 to -40.9)* | -24.9 (-29.2 to -20.4)* |
| Hepatitis C | 88.4 (72.0 to 107.8) | 90.9 (73.8 to 112.2) | 83.7 (66.3 to 104.2) | -5.3 (-17.1 to 7.5) | -7.9 (-17.6 to 3.0) | 1.8 (1.5 to 2.3) | 1.5 (1.2 to 1.9) | 1.1 (0.9 to 1.4) | -37.1 (-44.2 to -29.1)* | -22.3 (-30.4 to -13.0)* |
| Acute hepatitis E | 3000.4 (2479.7 to 3449.1) | 2405.2 (2090.1 to 2702.9) | 1419.6 (1230.3 to 1610.1) | -52.7 (-60.4 to -44.5)* | -41.0 (-47.5 to -33.9)* | 48.1 (40.0 to 55.0) | 35.1 (30.5 to 39.4) | 19.4 (16.9 to 22.0) | -59.6 (-65.9 to -52.9)* | -44.6 (-50.6 to -38.0)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|--------------------------------------|---|---|---|---------------------------------|---------------------------------|--|--|--|------------------------------------|-----------------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 |
| (Continued from previous page) | | | | | | | | | | |
| Other infectious diseases | 7066.9 (4328.0 to 9252.2) | 5858.1 (4069.7 to 7316.8) | 5671.3 (3883.9 to 7280.7) | -19.8 (-39.1 to 6.8) | -3.2 (-20.2 to 20.7) | 116.1 (74.8 to 148.6) | 88.2 (61.7 to 109.9) | 80.0 (54.2 to 103.5) | -31.1 (-46.9 to -10.6)* | -9.4 (-25.1 to 13.1) |
| Non-communicable diseases | 1 074 539.5 (957 634.8 to 1 197 008.3) | 1 312 102.0 (1 163 063.8 to 1 472 494.4) | 1 468 000.0 (1 296 535.4 to 1 658 537.7) | 36.6 (32.9 to 39.9)* | 11.9 (10.3 to 13.2)* | 25 460.8 (22 961.5 to 28 096.0) | 22 707.2 (20 288.0 to 25 308.0) | 20 786.9 (18 438.7 to 23 410.0) | -18.4 (-20.6 to -16.4)* | -8.5 (-9.9 to -7.3)* |
| Neoplasms | 1 515 506.6 (1 484 429.7 to 1 551 144.0) | 1 890 944.4 (1 859 939.0 to 1 920 035.5) | 2 132 221.0 (2 084 458.6 to 2 175 844.4) | 40.7 (36.8 to 44.9)* | 12.8 (10.8 to 15.0)* | 3 812.1 (3 739.6 to 3 895.3) | 3 377.2 (3 321.1 to 3 427.3) | 3 024.9 (2 956.4 to 3 086.9) | -20.6 (-22.7 to -18.4)* | -10.4 (-12.0 to -8.7)* |
| Lip and oral cavity cancer | 2583.8 (2488.9 to 2719.1) | 3668.0 (3546.8 to 3781.2) | 4634.7 (4429.7 to 4823.4) | 79.4 (64.8 to 88.9)* | 26.4 (20.8 to 31.5)* | 65.9 (63.5 to 69.2) | 64.6 (62.5 to 66.6) | 64.4 (61.6 to 67.0) | -2.2 (-10.0 to 2.9) | -0.2 (-4.5 to 3.8) |
| Nasopharynx cancer | 1590.5 (1506.1 to 1686.4) | 1785.6 (1722.8 to 1842.5) | 1906.0 (1809.3 to 2011.6) | 19.8 (13.1 to 28.0)* | 6.7 (0.2 to 13.1)* | 37.7 (35.7 to 40.0) | 30.1 (29.1 to 31.0) | 26.0 (24.7 to 27.4) | -31.1 (-34.9 to -26.4)* | -13.5 (-18.7 to -8.5)* |
| Other pharynx cancer | 1794.9 (1683.0 to 1987.4) | 2530.4 (2412.5 to 2620.7) | 3209.6 (2949.9 to 3393.4) | 78.8 (51.1 to 97.1)* | 26.8 (15.8 to 34.9)* | 45.5 (42.7 to 50.3) | 44.5 (42.4 to 46.1) | 44.4 (40.8 to 46.9) | -2.3 (-17.3 to 7.5) | -0.2 (-8.7 to 6.2) |
| Oesophageal cancer | 7809.6 (7554.6 to 8065.4) | 9206.9 (9018.2 to 9394.6) | 9276.1 (9019.5 to 9564.2) | 18.8 (13.7 to 24.4)* | 0.8 (-2.2 to 4.3) | 207.4 (200.7 to 214.2) | 170.0 (166.6 to 173.5) | 132.7 (129.1 to 136.8) | -36.0 (-38.8 to -33.1)* | -22.0 (-24.2 to -19.3)* |
| Stomach cancer | 19 027.0 (18 463.7 to 19 584.4) | 19 048.8 (18 712.9 to 19 405.6) | 18 345.5 (17 888.0 to 18 860.9) | -3.6 (-6.8 to 0.4) | -3.7 (-6.2 to -1.2)* | 499.4 (485.0 to 513.6) | 418.5 (342.0 to 354.6) | 262.9 (256.2 to 270.1) | -47.4 (-49.1 to -45.3)* | -24.5 (-26.5 to -22.6)* |
| Colon and rectum cancer | 11 021.7 (10 741.7 to 11 482.7) | 14 637.5 (14 356.4 to 14 918.1) | 17 197.2 (16 480.2 to 17 851.9) | 56.0 (44.5 to 63.5)* | 17.5 (11.6 to 22.3)* | 297.6 (290.6 to 309.1) | 272.4 (267.2 to 277.4) | 249.0 (238.6 to 258.4) | -16.4 (-22.3 to -12.5)* | -8.6 (-13.0 to -4.8)* |
| Liver cancer | 13 311.9 (12 450.1 to 14 010.2) | 18 332.3 (17 719.9 to 18 813.2) | 21 143.8 (20 268.1 to 21 985.9) | 58.8 (49.2 to 70.4)* | 15.3 (11.5 to 19.9)* | 331.7 (311.0 to 348.5) | 321.7 (310.9 to 329.8) | 295.2 (283.0 to 306.7) | -11.0 (-16.4 to -4.6)* | -8.2 (-11.3 to -4.7)* |
| Due to hepatitis B | 6537.7 (5718.9 to 7253.4) | 8838.7 (7823.1 to 9718.0) | 9802.3 (8568.3 to 10 960.1) | 49.9 (39.7 to 63.4)* | 10.9 (6.3 to 16.3)* | 158.5 (138.5 to 176.2) | 151.2 (133.7 to 167.0) | 134.7 (118.2 to 150.6) | -15.0 (-20.6 to -7.6)* | -10.9 (-14.4 to -6.6)* |
| Due to hepatitis C | 1850.2 (1639.8 to 2057.7) | 2735.7 (2447.6 to 3011.7) | 3310.5 (2930.7 to 3673.3) | 78.9 (69.0 to 86.6)* | 21.0 (16.9 to 24.9)* | 49.6 (44.1 to 55.3) | 51.2 (45.9 to 56.4) | 47.9 (43.5 to 53.0) | -3.6 (-8.7 to 0.3) | -6.5 (-9.8 to -3.7)* |
| Due to alcohol use | 1686.3 (1384.8 to 2013.9) | 2282.2 (1912.1 to 2710.9) | 2925.7 (2463.2 to 3400.9) | 73.5 (55.4 to 102.0)* | 28.2 (20.4 to 38.5)* | 45.0 (36.9 to 53.3) | 42.1 (35.2 to 49.9) | 41.8 (35.4 to 48.5) | -7.0 (-16.5 to 7.8) | -0.6 (-6.7 to 7.2) |
| Due to other causes | 3237.7 (2880.7 to 3688.9) | 4475.8 (4004.8 to 5058.9) | 5105.3 (4527.7 to 5757.7) | 57.7 (46.9 to 68.8)* | 14.1 (8.5 to 19.1)* | 78.6 (70.0 to 89.7) | 77.2 (69.0 to 87.0) | 70.8 (62.9 to 79.7) | -9.9 (-15.9 to -3.8)* | -8.3 (-12.4 to -4.6)* |
| Gallbladder and biliary tract cancer | 2311.8 (2201.3 to 2555.0) | 2885.3 (2655.7 to 3078.8) | 3311.1 (3002.9 to 3530.9) | 43.2 (26.0 to 53.5)* | 14.8 (9.7 to 19.8)* | 63.2 (60.3 to 69.5) | 53.9 (49.7 to 57.4) | 47.8 (43.5 to 50.9) | -24.3 (-33.2 to -19.0)* | -11.2 (-15.0 to -7.5)* |
| Pancreatic cancer | 4505.6 (4423.5 to 4576.2) | 6492.4 (6404.8 to 6602.1) | 8230.7 (8008.2 to 8445.8) | 82.7 (78.3 to 87.1)* | 26.8 (22.7 to 30.4)* | 122.8 (120.5 to 124.6) | 122.1 (120.5 to 124.1) | 119.5 (116.2 to 122.6) | -2.6 (-5.0 to -0.3)* | -2.1 (-5.2 to 0.7) |
| Larynx cancer | 2290.5 (2207.2 to 2369.6) | 2505.6 (2431.0 to 2569.8) | 2749.8 (2661.2 to 2845.7) | 20.1 (14.9 to 25.6)* | 9.8 (6.0 to 13.6)* | 59.6 (57.5 to 61.6) | 45.3 (44.0 to 46.5) | 38.7 (37.4 to 40.0) | -35.1 (-37.9 to -32.1)* | -14.6 (-17.5 to -11.7)* |
| Tracheal, bronchus, and lung cancer | 24 411.4 (23 862.3 to 25 086.9) | 32 059.3 (31 518.1 to 32 631.0) | 36 441.0 (35 401.2 to 37 462.8) | 49.3 (43.0 to 54.5)* | 13.7 (10.1 to 16.9)* | 650.6 (636.2 to 667.8) | 596.0 (586.5 to 606.2) | 526.1 (511.2 to 540.8) | -19.1 (-22.5 to -16.4)* | -11.7 (-14.4 to -9.2)* |
| Malignant skin melanoma | 947.1 (853.4 to 1081.6) | 1296.7 (1166.4 to 1434.0) | 1550.5 (1378.4 to 1714.1) | 63.7 (46.0 to 74.9)* | 19.6 (13.6 to 25.4)* | 23.6 (21.3 to 27.2) | 22.8 (20.5 to 25.1) | 21.8 (19.4 to 24.0) | -7.4 (-17.8 to -1.3)* | -4.3 (-9.0 to 0.3) |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|--|----------------------------------|------------------------------------|------------------------------------|------------------------------|----------------------------|--|---------------------------|---------------------------|------------------------------|----------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 |
| (Continued from previous page) | | | | | | | | | | |
| Non-melanoma skin cancer | 635.7 (611.9 to 658.6) | 864.4 (841.4 to 887.8) | 1022.6 (981.1 to 1068.7) | 60.9 (54.6 to 70.1)* | 18.3 (14.0 to 23.6)* | 17.0 (16.4 to 17.7) | 16.1 (15.7 to 16.6) | 14.9 (14.3 to 15.6) | -12.3 (-15.6 to -7.5)* | -7.6 (-10.8 to -3.5)* |
| Non-melanoma skin cancer (squamous-cell carcinoma) | 635.2 (611.4 to 658.1) | 863.5 (839.9 to 886.8) | 1021.5 (980.2 to 1067.6) | 60.8 (54.6 to 70.1)* | 18.3 (14.0 to 23.6)* | 17.0 (16.4 to 17.7) | 16.1 (15.7 to 16.6) | 14.9 (14.3 to 15.6) | -12.3 (-15.6 to -7.5)* | -7.6 (-10.8 to -3.5)* |
| Non-melanoma skin cancer (basal-cell carcinoma) | 0.5 (0.2 to 1.0) | 0.9 (0.4 to 1.8) | 1.1 (0.4 to 2.2) | 122.9 (109.6 to 138.3) | 23.8 (16.1 to 31.7) | .. | .. | .. | 15.8 (10.3 to 21.7) | -5.6 (-11.5 to 0.4) |
| Breast cancer | 9804.9 (9186.6 to 10 790.8) | 13 209.0 (12 668.7 to 13 875.4) | 15 107.8 (14 264.6 to 16 175.8) | 54.1 (34.8 to 70.0)* | 14.4 (6.3 to 22.2)* | 248.6 (233.7 to 272.3) | 229.4 (220.2 to 240.7) | 208.5 (196.9 to 223.1) | -16.2 (-26.2 to -8.2)* | -9.1 (-15.4 to -3.2)* |
| Cervical cancer | 6047.7 (5226.5 to 7655.0) | 7040.2 (5878.2 to 7516.1) | 7390.0 (6019.9 to 7868.6) | 22.2 (2.0 to 41.2)* | 5.0 (-1.3 to 13.2) | 149.0 (129.1 to 188.6) | 119.3 (99.8 to 127.4) | 100.6 (82.0 to 107.1) | -32.5 (-43.5 to -22.0)* | -15.7 (-20.8 to -9.2)* |
| Uterine cancer | 1573.9 (1480.6 to 1651.3) | 1955.1 (1868.8 to 2035.2) | 2122.8 (2010.4 to 2229.7) | 34.9 (26.3 to 47.1)* | 8.6 (2.7 to 16.4)* | 41.3 (39.0 to 43.3) | 35.6 (34.0 to 37.0) | 30.1 (28.5 to 31.7) | -27.1 (-31.6 to -20.6)* | -15.3 (-19.9 to -9.3)* |
| Ovarian cancer | 2661.5 (2531.2 to 2741.5) | 3523.0 (3393.4 to 3660.7) | 4258.1 (4035.8 to 4459.3) | 60.0 (51.8 to 68.2)* | 20.9 (13.9 to 27.0)* | 68.3 (65.3 to 70.2) | 62.5 (60.2 to 64.8) | 59.3 (56.2 to 62.1) | -13.2 (-17.5 to -8.8)* | -5.0 (-10.4 to -0.2)* |
| Prostate cancer | 3226.7 (2808.5 to 3505.0) | 4765.1 (4006.7 to 5107.8) | 6073.7 (4992.7 to 6582.3) | 88.2 (67.8 to 100.9)* | 27.5 (20.3 to 33.4)* | 97.5 (85.4 to 105.9) | 97.2 (82.2 to 104.2) | 93.9 (77.2 to 101.8) | -3.7 (-13.6 to 2.7) | -3.4 (-8.7 to 1.3) |
| Testicular cancer | 400.0 (380.0 to 423.9) | 393.1 (378.2 to 412.0) | 391.8 (372.4 to 412.0) | -2.0 (-8.2 to 3.7) | -0.3 (-4.4 to 4.0) | 7.9 (7.5 to 8.3) | 5.9 (5.7 to 6.2) | 5.2 (4.9 to 5.4) | -34.3 (-38.5 to -30.5)* | -12.3 (-15.8 to -8.6)* |
| Kidney cancer | 1753.1 (1707.2 to 1806.3) | 2476.1 (2418.4 to 2528.9) | 3023.6 (2911.4 to 3135.9) | 72.5 (64.5 to 80.9)* | 22.1 (17.6 to 26.5)* | 44.2 (43.2 to 45.4) | 45.0 (44.0 to 46.0) | 43.4 (41.8 to 45.0) | -1.9 (-6.1 to 2.2) | -3.7 (-7.2 to -0.2)* |
| Bladder cancer | 2235.8 (2157.3 to 2326.8) | 2796.3 (2735.8 to 2859.6) | 3315.2 (3193.2 to 3425.5) | 48.3 (38.9 to 53.9)* | 18.6 (13.9 to 22.4)* | 63.3 (61.2 to 65.7) | 54.3 (53.1 to 55.6) | 49.5 (47.7 to 51.1) | -21.8 (-26.6 to -19.0)* | -9.0 (-12.4 to -6.1)* |
| Brain and nervous system cancer | 5620.2 (4833.4 to 5997.6) | 6729.7 (6094.9 to 7049.7) | 7660.0 (6922.8 to 8280.4) | 36.3 (25.9 to 54.7)* | 13.8 (9.4 to 20.8)* | 116.8 (101.5 to 123.9) | 109.1 (98.5 to 114.1) | 105.0 (94.9 to 113.3) | -10.0 (-16.4 to 2.6) | -3.7 (-7.3 to 2.3) |
| Thyroid cancer | 768.4 (713.7 to 817.8) | 960.7 (918.0 to 1004.4) | 1122.8 (1071.3 to 1183.5) | 46.1 (35.1 to 60.9)* | 16.9 (11.0 to 23.8)* | 19.9 (17.9 to 20.3) | 16.9 (16.2 to 17.6) | 15.8 (15.0 to 16.6) | -17.6 (-23.5 to -9.7)* | -6.8 (-11.4 to -1.3)* |
| Mesothelioma | 398.9 (361.5 to 440.9) | 535.4 (509.7 to 569.4) | 660.9 (619.4 to 700.6) | 65.7 (43.6 to 81.4)* | 23.4 (17.8 to 28.2)* | 10.5 (9.5 to 11.7) | 9.8 (9.3 to 10.3) | 9.5 (8.9 to 10.1) | -9.3 (-21.7 to -0.8)* | -2.8 (-7.1 to 0.9) |
| Hodgkin's lymphoma | 1589.4 (1176.0 to 1775.3) | 1269.1 (1048.1 to 1466.8) | 1126.8 (943.7 to 1337.1) | -29.1 (-35.9 to -15.9)* | -11.2 (-14.5 to -8.0)* | 31.8 (23.8 to 35.5) | 19.7 (16.4 to 22.9) | 15.2 (12.8 to 18.1) | -52.1 (-56.7 to -43.2)* | -22.7 (-25.4 to -20.1)* |
| Non-Hodgkin lymphoma | 4095.3 (3844.4 to 4377.0) | 5528.6 (5174.2 to 5736.3) | 6783.0 (6172.5 to 7083.3) | 65.6 (49.2 to 74.7)* | 22.7 (16.0 to 27.2)* | 92.8 (88.3 to 98.0) | 93.6 (88.1 to 96.9) | 94.9 (86.6 to 99.1) | 2.3 (-7.4 to 7.3) | 1.4 (-4.1 to 5.0) |
| Multiple myeloma | 1141.8 (1032.1 to 1308.8) | 1664.3 (1489.5 to 1831.8) | 2114.0 (1901.5 to 2339.1) | 85.2 (72.6 to 98.4)* | 27.0 (22.6 to 32.6)* | 30.7 (27.9 to 35.5) | 31.0 (27.6 to 34.0) | 30.5 (27.4 to 33.9) | -0.7 (-8.6 to 5.8) | -1.5 (-4.9 to 2.7) |
| Leukaemia | 10 455.3 (9415.6 to 12 242.4) | 10 401.6 (9581.8 to 11 171.9) | 10 204.0 (9319.4 to 10 809.1) | -2.4 (-14.2 to 7.2) | -1.9 (-6.2 to 2.3) | 204.7 (186.8 to 234.4) | 166.3 (153.6 to 178.1) | 141.6 (129.3 to 149.9) | -30.9 (-38.4 to -25.0)* | -14.9 (-18.4 to -11.3)* |
| Acute lymphoid leukaemia | 2158.2 (1900.3 to 3001.6) | 2339.9 (2178.5 to 2794.3) | 2426.6 (2216.0 to 2688.8) | 12.4 (-15.5 to 28.2) | 3.7 (-7.8 to 9.9) | 37.8 (33.5 to 51.5) | 35.0 (32.7 to 41.7) | 33.1 (30.2 to 36.6) | -12.5 (-33.3 to -1.5)* | -5.7 (-15.9 to -0.2)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|---------------------------------|---|---|---|---------------------------------------|-------------------------------------|--|--|--|--|--|
| | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 |
| (Continued from previous page) | | | | | | | | | | |
| Chronic lymphoid leukaemia | 468.8 (434.3 to 553.7) | 617.1 (580.4 to 709.0) | 703.1 (654.0 to 805.0) | 50.0 (38.9 to 61.4)* | 13.9 (9.2 to 19.1)* | 12.6 (11.7 to 14.6) | 11.6 (10.9 to 13.3) | 10.3 (9.7 to 11.8) | -17.6 (-22.8 to -12.2)* | -11.1 (-14.6 to -7.2)* |
| Acute myeloid leukaemia | 1707.6 (1577.2 to 2023.1) | 2327.1 (2152.7 to 2540.6) | 2651.6 (2444.2 to 2838.1) | 55.3 (34.6 to 68.5)* | 13.9 (8.7 to 18.2)* | 35.4 (33.1 to 41.0) | 38.0 (35.2 to 41.2) | 36.9 (34.1 to 39.4) | 4.3 (-8.5 to 12.2) | -2.9 (-7.0 to 0.6) |
| Chronic myeloid leukaemia | 706.2 (639.5 to 775.2) | 654.3 (596.6 to 723.6) | 609.6 (550.3 to 676.0) | -13.7 (-21.0 to -6.6)* | -6.8 (-11.1 to -2.2)* | 16.6 (15.2 to 18.1) | 11.0 (10.1 to 12.1) | 8.4 (7.6 to 9.3) | -49.5 (-53.5 to -45.7)* | -23.5 (-26.9 to -20.1)* |
| Other leukaemia | 5414.4 (4487.0 to 6178.9) | 4463.2 (3848.9 to 4762.2) | 3813.1 (3326.8 to 4039.8) | -29.6 (-38.9 to -18.9)* | -14.6 (-19.6 to -9.0)* | 102.3 (86.4 to 114.9) | 70.6 (60.9 to 75.1) | 52.9 (46.2 to 56.0) | -48.4 (-54.7 to -41.5)* | -25.1 (-29.4 to -20.4)* |
| Other neoplasms | 7536.4 (7307.3 to 7843.6) | 10 533.8 (9864.9 to 10 811.8) | 12 847.7 (11 679.5 to 13 268.0) | 70.5 (52.9 to 78.0)* | 22.0 (17.2 to 25.3)* | 163.6 (158.9 to 170.2) | 174.2 (163.3 to 178.5) | 178.5 (162.3 to 184.3) | 9.2 (-2.3 to 13.4) | 2.5 (-1.5 to 5.2) |
| Cardiovascular diseases | 266 709.6 (258 611.1 to 274 986.6) | 321 851.2 (312 966.3 to 330 487.6) | 353 120.9 (341 794.5 to 364 965.3) | 32.4 (28.8 to 36.2)* | 9.7 (7.4 to 12.2)* | 7267.3 (7049.6 to 7496.0) | 6046.0 (5878.1 to 6208.3) | 5178.4 (5010.6 to 5352.2) | -28.7 (-30.7 to -26.8)* | -14.3 (-16.1 to -12.4)* |
| Rheumatic heart disease | 13 277.2 (11 854.4 to 14 558.1) | 11 235.5 (10 229.5 to 12 162.0) | 9803.1 (9154.5 to 10 556.6) | -26.2 (-32.8 to -18.5)* | -12.8 (-18.3 to -6.9)* | 296.4 (263.4 to 324.9) | 185.4 (169.1 to 199.9) | 135.9 (127.1 to 145.8) | -54.2 (-58.2 to -49.1)* | -26.7 (-31.3 to -21.7)* |
| Ischaemic heart disease | 118 996.9 (115 809.8 to 122 723.6) | 153 547.6 (149 867.7 to 156 971.0) | 174 611.1 (169 736.1 to 179 827.6) | 46.7 (42.5 to 51.5)* | 13.7 (10.7 to 16.7)* | 3344.1 (3255.9 to 3448.7) | 2908.3 (2840.4 to 2971.7) | 2562.8 (2491.4 to 2640.8) | -23.4 (-25.6 to -21.0)* | -11.9 (-14.2 to -9.6)* |
| Cerebrovascular disease | 95 306.8 (91 616.6 to 100 606.2) | 111 916.0 (108 069.6 to 115 501.2) | 116 445.1 (111 385.4 to 121 406.9) | 22.2 (16.7 to 27.0)* | 4.0 (1.6 to 6.5)* | 2600.1 (2496.3 to 2744.0) | 2116.6 (2041.9 to 2186.6) | 1711.2 (1635.3 to 1784.4) | -34.2 (-37.2 to -31.5)* | -19.1 (-21.0 to -17.2)* |
| Ischaemic stroke | 37 687.2 (35 505.5 to 39 624.6) | 46 909.9 (44 168.6 to 49 455.6) | 51 897.4 (47 896.6 to 55 567.7) | 37.7 (32.2 to 43.4)* | 10.6 (7.1 to 14.0)* | 1112.1 (1049.6 to 1169.6) | 934.8 (880.6 to 985.5) | 787.5 (728.4 to 843.2) | -29.2 (-32.0 to -26.2)* | -15.8 (-18.4 to -13.1)* |
| Haemorrhagic stroke | 57 619.6 (55 206.4 to 61 574.1) | 65 006.1 (63 154.6 to 66 941.1) | 64 547.7 (62 622.3 to 66 497.6) | 12.0 (6.3 to 17.0)* | -0.7 (-2.8 to 1.5) | 1488.0 (1427.4 to 1594.8) | 1181.8 (1148.5 to 1216.2) | 923.6 (896.2 to 951.9) | -37.9 (-41.3 to -35.2)* | -21.8 (-23.5 to -20.1)* |
| Hypertensive heart disease | 12 136.6 (9824.4 to 13 763.9) | 13 571.4 (11 603.8 to 15 049.4) | 16 335.1 (13 456.6 to 17 843.7) | 34.6 (24.7 to 52.5)* | 20.4 (10.2 to 32.9)* | 338.6 (274.9 to 382.5) | 259.6 (221.8 to 287.4) | 242.5 (199.7 to 265.0) | -28.4 (-33.4 to -19.1)* | -6.6 (-14.6 to 2.8) |
| Cardiomyopathy and myocarditis | 6713.9 (5820.8 to 7248.0) | 8576.8 (7305.3 to 9168.4) | 8718.2 (7574.9 to 9628.0) | 29.9 (16.7 to 44.8)* | 1.6 (-7.1 to 12.9) | 159.0 (133.6 to 171.9) | 145.7 (123.9 to 155.8) | 122.7 (106.4 to 135.1) | -22.9 (-30.0 to -13.5)* | -15.8 (-22.8 to -6.8)* |
| Myocarditis | 1253.7 (1032.8 to 1618.8) | 1422.2 (1132.2 to 1574.2) | 1367.5 (1118.2 to 1513.1) | 9.1 (-20.2 to 28.8) | -3.9 (-12.7 to 5.6) | 24.8 (20.8 to 30.4) | 23.0 (18.1 to 25.4) | 19.1 (15.6 to 21.1) | -22.8 (-41.0 to -11.7)* | -16.8 (-24.3 to -8.0)* |
| Alcoholic cardiomyopathy | 1815.2 (1512.9 to 2177.5) | 2878.5 (2414.3 to 3221.2) | 2591.2 (2055.7 to 3240.6) | 42.8 (11.8 to 80.0)* | -10.0 (-27.6 to 14.3) | 44.5 (37.1 to 53.3) | 47.8 (40.1 to 53.2) | 35.2 (28.0 to 43.9) | -21.0 (-37.8 to -0.8)* | -26.3 (-40.4 to -7.6)* |
| Other cardiomyopathy | 3644.9 (2878.4 to 4234.4) | 4276.1 (3615.3 to 4655.1) | 4759.4 (4089.9 to 5094.4) | 30.6 (14.5 to 52.0)* | 11.3 (4.3 to 19.4)* | 89.7 (68.9 to 103.1) | 75.0 (63.1 to 81.5) | 68.3 (58.5 to 73.2) | -23.8 (-32.3 to -9.1)* | -8.8 (-14.3 to -2.7)* |
| Atrial fibrillation and flutter | 3044.8 (2345.7 to 3888.0) | 4460.7 (3484.6 to 5666.6) | 5951.3 (4649.6 to 7516.9) | 95.5 (91.1 to 100.0)* | 33.4 (31.8 to 35.1)* | 97.8 (76.0 to 123.5) | 93.7 (73.6 to 118.1) | 93.4 (73.2 to 118.1) | -4.5 (-6.5 to -2.4)* | -0.2 (-1.3 to 0.8) |
| Aortic aneurysm | 1942.5 (1855.5 to 2068.9) | 2510.2 (2437.9 to 2611.8) | 2881.8 (2800.8 to 2975.5) | 48.4 (39.9 to 58.1)* | 14.8 (10.7 to 20.1)* | 54.4 (52.1 to 57.6) | 47.8 (46.5 to 49.6) | 42.6 (41.4 to 44.0) | -21.7 (-25.8 to -17.0)* | -10.9 (-13.9 to -7.1)* |
| Peripheral artery disease | 678.4 (486.1 to 948.0) | 970.9 (719.7 to 1325.7) | 1235.1 (922.2 to 1683.9) | 82.1 (68.0 to 103.9)* | 27.2 (20.9 to 35.7)* | 21.4 (15.4 to 29.7) | 20.3 (15.1 to 27.9) | 19.3 (14.4 to 26.3) | -9.4 (-17.0 to 0.5) | -4.9 (-9.6 to 1.3) |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|---|--|--|--|---------------------------------|-------------------------------|--|--------------------------------------|--------------------------------------|------------------------------------|------------------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 |
| (Continued from previous page) | | | | | | | | | | |
| Endocarditis | 1536.3 (1320.2 to 1751.3) | 1998.3 (1748.2 to 2320.0) | 2368.7 (2104.5 to 2797.1) | 54.2 (39.0 to 69.5)* | 18.5 (12.0 to 23.8)* | 34.6 (29.7 to 39.2) | 34.0 (29.6 to 39.2) | 33.6 (29.8 to 39.7) | -3.0 (-11.4 to 5.3) | -1.3 (-6.7 to 2.9) |
| Other cardiovascular and circulatory diseases | 13 076.2 (10 976.0 to 14 815.8) | 13 063.8 (11 561.4 to 15 422.4) | 14 771.4 (13 010.9 to 17 397.0) | 13.0 (3.2 to 31.9)* | 13.1 (9.8 to 17.3)* | 320.9 (269.6 to 364.9) | 234.7 (208.1 to 276.1) | 214.5 (189.4 to 251.8) | -33.2 (-38.7 to -22.3)* | -8.6 (-11.2 to -5.3)* |
| Chronic respiratory diseases | 86 833.7 (79 815.5 to 92 631.5) | 86 665.1 (81 193.2 to 92 630.9) | 92 528.7 (86 142.3 to 99 725.6) | 6.6 (2.0 to 15.4)* | 6.8 (4.5 to 9.7)* | 2256.8 (2079.1 to 2386.5) | 1601.5 (1507.8 to 1698.0) | 1351.7 (1261.3 to 1452.3) | -40.1 (-43.1 to -34.6)* | -15.6 (-17.6 to -13.2)* |
| Chronic obstructive pulmonary disease | 59 810.7 (53 271.2 to 64 128.3) | 59 572.5 (56 816.4 to 62 571.2) | 63 434.3 (60 586.3 to 66 676.3) | 6.1 (0.6 to 18.4)* | 6.5 (3.8 to 10.1)* | 1666.8 (1486.0 to 1780.7) | 1151.0 (1098.0 to 1205.8) | 945.3 (904.1 to 994.8) | -43.3 (-46.2 to -36.8)* | -17.9 (-19.9 to -15.1)* |
| Pneumoconiosis | 568.0 (494.8 to 787.9) | 560.3 (506.1 to 635.2) | 577.1 (517.2 to 647.3) | 1.6 (-2.5 to 16.9) | 3.0 (-4.5 to 8.4) | 15.0 (13.2 to 20.5) | 10.4 (9.4 to 11.8) | 8.4 (7.6 to 9.4) | -44.0 (-58.3 to -36.2)* | -19.1 (-24.6 to -15.0)* |
| Silicosis | 303.5 (251.4 to 461.0) | 273.2 (247.2 to 310.8) | 270.6 (243.6 to 301.5) | -10.8 (-43.4 to 6.3) | -0.9 (-14.2 to 5.7) | 8.0 (6.7 to 11.9) | 5.0 (4.5 to 5.7) | 3.9 (3.5 to 4.4) | -50.9 (-68.2 to -42.0)* | -22.1 (-32.2 to -17.1)* |
| Asbestosis | 50.9 (40.8 to 70.6) | 73.0 (57.2 to 86.9) | 83.9 (67.9 to 97.5) | 64.8 (29.6 to 83.7)* | 14.8 (9.2 to 21.8)* | 1.3 (1.1 to 1.9) | 1.4 (1.1 to 1.6) | 1.2 (1.0 to 1.4) | -5.8 (-26.4 to 4.6) | -9.2 (-13.7 to -3.3)* |
| Coal workers' pneumoconiosis | 103.4 (70.5 to 127.8) | 87.5 (65.5 to 104.6) | 89.1 (70.2 to 108.9) | -13.9 (-30.6 to 15.1) | 1.8 (-6.8 to 12.7) | 2.9 (1.9 to 3.5) | 1.7 (1.2 to 2.0) | 1.3 (1.0 to 1.6) | -54.3 (-63.3 to -38.6)* | -21.6 (-28.2 to -13.5)* |
| Other pneumoconiosis | 110.2 (80.0 to 173.3) | 126.6 (104.4 to 161.8) | 133.5 (112.1 to 165.9) | 21.2 (-8.3 to 52.8) | 5.5 (-2.2 to 13.2) | 2.9 (2.1 to 4.5) | 2.3 (1.9 to 3.0) | 1.9 (1.6 to 2.4) | -32.1 (-47.9 to -15.2)* | -16.7 (-22.7 to -10.6)* |
| Asthma | 23 840.9 (19 339.2 to 28 695.4) | 22 827.4 (18 531.6 to 27 887.8) | 23 720.5 (18 911.2 to 29 790.8) | -0.5 (-11.3 to 9.8) | 3.9 (-0.8 to 8.2) | 514.7 (419.4 to 621.4) | 375.2 (305.9 to 454.5) | 329.2 (263.1 to 413.2) | -36.0 (-44.1 to -28.2)* | -12.2 (-16.9 to -7.7)* |
| Interstitial lung disease and pulmonary sarcoidosis | 1285.5 (924.8 to 1966.3) | 2052.5 (1506.0 to 2609.2) | 2721.6 (2075.3 to 3177.4) | 111.7 (55.3 to 165.5)* | 32.6 (18.7 to 43.0)* | 33.8 (24.7 to 50.2) | 38.7 (28.4 to 48.6) | 40.2 (30.5 to 46.9) | 19.0 (-11.4 to 49.3) | 3.9 (-6.4 to 11.7) |
| Other chronic respiratory diseases | 1328.6 (906.1 to 1714.7) | 1652.5 (1286.3 to 1932.2) | 2075.2 (1697.7 to 2315.1) | 56.2 (27.5 to 93.2)* | 25.6 (17.5 to 34.0)* | 26.6 (18.3 to 33.4) | 26.2 (20.3 to 30.5) | 28.5 (23.3 to 31.8) | 7.4 (-10.0 to 30.2) | 9.0 (2.1 to 16.4)* |
| Cirrhosis and other chronic liver diseases | 28 184.1 (26 769.7 to 29 706.9) | 36 122.1 (34 848.4 to 39 549.1) | 38 856.7 (36 890.5 to 42 795.8) | 37.9 (28.7 to 49.4)* | 7.6 (2.5 to 13.7)* | 659.0 (627.6 to 693.9) | 603.3 (582.6 to 657.4) | 531.1 (504.3 to 584.5) | -19.4 (-24.6 to -12.8)* | -12.0 (-16.1 to -7.2)* |
| Due to hepatitis B | 7875.2 (7193.4 to 8627.5) | 10 373.1 (9525.8 to 11 593.2) | 11 240.7 (10 162.6 to 13 138.6) | 42.7 (31.8 to 58.9)* | 8.4 (2.5 to 15.3)* | 189.9 (173.4 to 208.5) | 173.9 (160.1 to 195.5) | 153.3 (138.5 to 178.9) | -19.3 (-25.4 to -10.3)* | -11.8 (-16.5 to -6.1)* |
| Due to hepatitis C | 6333.5 (5732.9 to 7049.7) | 8709.0 (7906.3 to 9751.8) | 9769.1 (8819.4 to 10 985.0) | 54.2 (44.4 to 66.7)* | 12.2 (6.8 to 18.5)* | 155.1 (139.9 to 172.8) | 147.4 (133.6 to 164.2) | 133.5 (120.7 to 150.0) | -13.9 (-19.3 to -7.3)* | -9.4 (-13.8 to -4.4)* |
| Due to alcohol use | 6466.4 (5920.0 to 7057.2) | 8878.9 (8112.9 to 9688.5) | 9754.0 (8873.6 to 10 861.8) | 50.8 (42.0 to 61.5)* | 9.9 (4.8 to 15.9)* | 161.5 (148.2 to 175.7) | 151.3 (138.5 to 165.1) | 133.5 (121.6 to 148.2) | -17.3 (-22.0 to -11.6)* | -11.8 (-15.7 to -6.9)* |
| Due to other causes | 7509.0 (6800.3 to 8254.4) | 8161.1 (7451.7 to 9036.0) | 8092.8 (7310.3 to 9160.6) | 7.8 (-1.9 to 20.4) | -0.8 (-5.9 to 5.0) | 152.5 (138.5 to 167.9) | 130.7 (119.2 to 144.8) | 110.8 (100.2 to 125.3) | -27.4 (-32.9 to -19.2)* | -15.3 (-19.4 to -10.9)* |
| Digestive diseases | 32 345.7 (28 585.3 to 34 769.7) | 33 020.0 (30 716.0 to 35 440.1) | 34 368.9 (31 754.4 to 37 405.9) | 6.3 (-0.5 to 17.9) | 4.1 (0.4 to 7.7)* | 720.8 (640.2 to 770.7) | 561.3 (523.0 to 601.8) | 484.9 (448.4 to 526.8) | -32.7 (-36.3 to -26.0)* | -13.6 (-16.4 to -10.8)* |
| Peptic ulcer disease | 10 011.0 (8963.1 to 10 836.7) | 7708.0 (7085.6 to 8454.8) | 7106.0 (6459.4 to 8011.1) | -29.0 (-35.4 to -20.6)* | -7.8 (-11.8 to -3.8)* | 237.1 (212.3 to 257.4) | 134.0 (123.2 to 146.5) | 100.2 (91.2 to 112.5) | -57.8 (-61.6 to -52.7)* | -25.2 (-28.4 to -22.0)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|--|--|---|---|---------------------------------------|---------------------------------------|--|--|--|---------------------------------------|---------------------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 |
| (Continued from previous page) | | | | | | | | | | |
| Gastritis and duodenitis | 1952.5 (1571.5 to 2395.4) | 2235.1 (1806.0 to 2810.5) | 2689.8 (2104.0 to 3454.3) | 37.8 (24.2 to 50.6)* | 20.4 (13.5 to 26.7)* | 45.9 (37.1 to 56.0) | 38.4 (31.2 to 48.1) | 37.7 (29.6 to 48.3) | -17.8 (-25.0 to -10.7)* | -1.7 (-7.2 to 3.2) |
| Appendicitis | 2200.3 (1726.6 to 2565.2) | 2412.3 (2109.1 to 2744.1) | 2150.0 (1928.1 to 2484.8) | -2.3 (-16.9 to 19.1) | -10.9 (-18.1 to -3.0)* | 41.3 (32.8 to 47.4) | 37.2 (32.3 to 42.2) | 29.3 (26.3 to 33.9) | -29.0 (-38.5 to -15.1)* | -21.1 (-27.5 to -14.2)* |
| Paralytic ileus and intestinal obstruction | 6853.7 (5275.8 to 7778.5) | 7379.6 (6085.8 to 7981.1) | 7626.5 (6380.0 to 8328.6) | 11.3 (-1.7 to 32.7) | 3.4 (-2.8 to 10.6) | 131.2 (101.5 to 146.5) | 120.6 (99.3 to 130.0) | 108.0 (90.5 to 117.7) | -17.7 (-25.7 to -3.7)* | -10.5 (-15.4 to -5.0)* |
| Inguinal, femoral, and abdominal hernia | 2405.0 (1863.4 to 3003.9) | 2854.3 (2251.5 to 3551.6) | 3106.1 (2401.1 to 3932.1) | 29.1 (19.9 to 39.6)* | 8.8 (5.5 to 11.4)* | 54.2 (42.2 to 67.3) | 48.2 (38.2 to 59.9) | 43.5 (33.7 to 55.0) | -19.7 (-26.4 to -12.2)* | -9.7 (-12.8 to -7.6)* |
| Inflammatory bowel disease | 1217.7 (916.1 to 1735.5) | 1606.7 (1305.0 to 2000.0) | 1844.4 (1541.5 to 2195.5) | 51.5 (44.2 to 81.3)* | 14.8 (2.7 to 22.6)* | 27.4 (21.5 to 36.3) | 27.4 (22.3 to 33.5) | 25.8 (21.6 to 30.7) | -6.0 (-23.3 to 7.3) | -5.9 (-14.8 to 0.3) |
| Vascular intestinal disorders | 1012.2 (904.2 to 1107.2) | 1457.8 (1317.3 to 1656.0) | 1702.6 (1568.9 to 1943.4) | 68.2 (55.2 to 86.2)* | 16.8 (10.4 to 23.5)* | 28.1 (25.5 to 30.5) | 28.1 (25.7 to 31.8) | 25.5 (23.4 to 29.0) | -9.3 (-15.3 to 0.2) | -9.4 (-14.1 to -4.6)* |
| Gallbladder and biliary diseases | 1665.1 (1368.2 to 1821.1) | 1851.7 (1729.6 to 2093.8) | 2098.0 (1953.3 to 2401.6) | 26.0 (14.4 to 56.3)* | 13.3 (9.8 to 17.6)* | 42.8 (35.5 to 47.3) | 33.9 (31.8 to 38.2) | 30.5 (28.4 to 35.0) | -28.8 (-36.1 to -12.3)* | -10.0 (-12.8 to -6.6)* |
| Pancreatitis | 2035.5 (1723.4 to 2341.3) | 3046.0 (2707.5 to 3380.8) | 3347.0 (2910.4 to 3722.1) | 64.4 (44.7 to 82.4)* | 9.9 (2.9 to 16.8)* | 47.5 (40.4 to 54.7) | 50.5 (44.9 to 56.2) | 45.8 (39.8 to 50.9) | -3.6 (-15.1 to 6.6) | -9.4 (-15.1 to -3.9)* |
| Other digestive diseases | 2992.8 (2482.9 to 3546.7) | 2468.6 (2189.1 to 2834.9) | 2698.5 (2428.7 to 3008.7) | -9.8 (-24.4 to 4.2) | 9.3 (1.9 to 17.8)* | 65.2 (56.4 to 75.0) | 43.0 (38.7 to 48.8) | 38.6 (34.8 to 43.0) | -40.8 (-49.1 to -33.3)* | -10.2 (-15.5 to -4.2)* |
| Neurological disorders | 64 973.1 (50 343.0 to 80 732.5) | 87 251.9 (67 830.9 to 107 955.1) | 103 580.0 (81 171.2 to 128 122.4) | 59.4 (55.2 to 64.6)* | 18.7 (17.0 to 20.7)* | 1500.6 (1192.1 to 1832.8) | 1491.2 (1182.9 to 1824.0) | 1478.4 (1171.9 to 1813.0) | -1.5 (-3.3 to 0.4) | -0.9 (-2.1 to 0.3) |
| Alzheimer's disease and other dementias | 13 024.5 (11 052.7 to 15 479.9) | 20 912.2 (17 919.7 to 24 689.5) | 28 764.1 (24 510.8 to 33 952.4) | 120.8 (115.3 to 126.5)* | 37.5 (35.3 to 39.7)* | 460.9 (394.5 to 544.4) | 469.6 (403.2 to 552.4) | 470.6 (401.2 to 556.3) | 2.1 (0.1 to 3.8)* | 0.2 (-1.1 to 1.5) |
| Parkinson's disease | 1304.3 (1024.7 to 1606.9) | 2385.3 (1901.1 to 2910.3) | 3234.5 (2563.6 to 4012.8) | 148.0 (139.8 to 155.8)* | 35.6 (32.9 to 38.2)* | 42.0 (33.2 to 51.9) | 49.9 (39.7 to 61.0) | 51.3 (40.6 to 63.4) | 22.1 (18.2 to 25.9)* | 2.7 (1.0 to 4.5)* |
| Epilepsy | 12 420.8 (10 285.4 to 14 852.0) | 13 435.2 (11 142.0 to 16 031.4) | 13 492.2 (11 014.7 to 16 503.1) | 8.6 (-2.9 to 23.7) | 0.4 (-7.0 to 8.6) | 226.5 (187.9 to 269.3) | 201.6 (166.8 to 240.4) | 182.6 (148.9 to 223.5) | -19.4 (-27.6 to -9.0)* | -9.4 (-16.1 to -2.2)* |
| Multiple sclerosis | 694.0 (586.0 to 807.1) | 974.2 (827.1 to 1123.9) | 1151.5 (968.6 to 1345.8) | 65.9 (45.2 to 74.8)* | 18.2 (12.8 to 21.9)* | 16.3 (13.8 to 18.9) | 16.2 (13.8 to 18.6) | 15.6 (13.2 to 18.3) | -4.2 (-16.4 to 0.8) | -3.2 (-7.3 to -0.2)* |
| Motor neuron disease | 582.3 (527.8 to 651.4) | 774.9 (745.5 to 817.5) | 926.1 (881.6 to 961.8) | 59.0 (44.5 to 71.6)* | 19.5 (14.7 to 22.4)* | 13.4 (12.5 to 14.6) | 13.7 (13.2 to 14.3) | 13.2 (12.5 to 13.7) | -1.5 (-9.3 to 2.9) | -3.6 (-7.3 to -1.3)* |
| Migraine | 29 843.4 (19 092.9 to 41 793.9) | 39 485.3 (25 341.5 to 55 187.3) | 45 121.9 (29 045.8 to 62 826.9) | 51.2 (49.7 to 52.8)* | 14.3 (13.7 to 14.9)* | 599.9 (385.7 to 839.1) | 597.8 (384.6 to 833.2) | 598.6 (385.9 to 833.3) | -0.2 (-0.8 to 0.4) | 0.1 (-0.2 to 0.5) |
| Tension-type headache | 4700.9 (2968.1 to 6989.3) | 6236.2 (3972.6 to 9204.2) | 7195.1 (4614.6 to 10 499.9) | 53.1 (47.5 to 58.4)* | 15.4 (13.7 to 17.0)* | 96.2 (61.1 to 142.5) | 95.5 (61.2 to 139.9) | 95.9 (61.5 to 140.0) | -0.2 (-2.5 to 1.9) | 0.4 (-0.5 to 1.4) |
| Other neurological disorders | 2402.9 (2004.0 to 2829.7) | 3048.7 (2593.3 to 3553.3) | 3694.5 (3114.1 to 4353.4) | 53.8 (38.5 to 67.8)* | 21.2 (14.0 to 28.5)* | 45.5 (38.7 to 54.4) | 46.9 (40.2 to 54.4) | 50.6 (42.7 to 59.5) | 11.2 (2.1 to 20.0)* | 7.9 (1.9 to 14.2)* |
| Mental and substance use disorders | 110 918.3 (83 056.1 to 141 228.0) | 145 067.1 (108 650.3 to 183 887.6) | 162 509.3 (121 886.4 to 206 517.4) | 46.5 (44.9 to 48.7)* | 12.0 (11.2 to 12.9)* | 2240.8 (1681.3 to 2844.2) | 2226.6 (1669.0 to 2821.8) | 2172.7 (1629.4 to 2761.5) | -3.0 (-4.1 to -1.8)* | -2.4 (-3.2 to -1.7)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|---|---------------------------------|---------------------------------|---------------------------------|------------------------------|----------------------------|--|---------------------------|---------------------------|------------------------------|----------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 |
| (Continued from previous page) | | | | | | | | | | |
| Schizophrenia | 8447.6 (6181.9 to 10521.3) | 11492.6 (8420.0 to 14326.7) | 13414.3 (9858.7 to 16714.0) | 58.8 (55.8 to 62.0)* | 16.7 (15.5 to 18.0)* | 179.6 (132.2 to 222.3) | 178.9 (132.0 to 222.3) | 177.2 (130.5 to 220.3) | -1.3 (-2.2 to -0.5)* | -0.9 (-1.7 to -0.2)* |
| Alcohol use disorders | 11264.3 (9057.6 to 14013.8) | 15561.7 (12607.6 to 19100.6) | 16244.7 (13003.1 to 19955.3) | 44.2 (37.4 to 51.0)* | 4.4 (0.4 to 8.6)* | 237.3 (192.3 to 292.7) | 240.9 (196.4 to 293.9) | 214.4 (172.0 to 262.9) | -9.6 (-14.3 to -5.5)* | -11.0 (-14.6 to -7.5)* |
| Drug use disorders | 14247.5 (11250.4 to 17370.8) | 18009.0 (14444.2 to 21739.6) | 20394.2 (16204.4 to 24670.3) | 43.1 (36.5 to 58.5)* | 13.2 (9.8 to 16.9)* | 278.6 (221.2 to 337.7) | 270.5 (217.3 to 325.1) | 268.4 (213.5 to 324.0) | -3.6 (-8.1 to 6.7) | -0.8 (-3.8 to 2.4) |
| Opioid use disorders | 10261.8 (8052.7 to 12572.0) | 12817.5 (10018.1 to 15694.7) | 14788.8 (11380.5 to 18259.7) | 44.1 (37.5 to 56.0)* | 15.4 (11.2 to 19.5)* | 202.0 (159.3 to 247.4) | 193.1 (151.2 to 236.4) | 194.3 (149.8 to 239.6) | -3.8 (-8.3 to 3.8) | 0.7 (-3.0 to 4.2) |
| Cocaine use disorders | 790.1 (568.0 to 1051.1) | 1060.8 (780.4 to 1375.9) | 1154.2 (847.2 to 1512.1) | 46.1 (38.4 to 65.1)* | 8.8 (5.1 to 12.2)* | 15.7 (11.3 to 20.8) | 16.1 (11.8 to 20.8) | 15.3 (11.2 to 20.0) | -2.6 (-7.8 to 9.6) | -4.9 (-8.3 to -1.7)* |
| Amphetamine use disorders | 660.3 (430.6 to 985.6) | 834.2 (567.1 to 1190.4) | 881.7 (599.5 to 1243.1) | 33.5 (23.0 to 54.2)* | 5.7 (1.8 to 10.2)* | 11.8 (7.8 to 17.3) | 12.0 (8.2 to 17.0) | 11.5 (7.8 to 16.2) | -1.9 (-8.8 to 13.2) | -3.7 (-7.5 to 0.3) |
| Cannabis use disorders | 514.5 (321.8 to 757.2) | 623.9 (389.2 to 905.4) | 646.9 (400.9 to 945.5) | 25.7 (21.7 to 29.8)* | 3.7 (1.2 to 6.0)* | 9.1 (5.7 to 13.3) | 8.8 (5.5 to 12.9) | 8.5 (5.2 to 12.4) | -6.9 (-8.9 to -5.0)* | -4.2 (-5.9 to -2.4)* |
| Other drug use disorders | 2020.8 (1550.8 to 2453.2) | 2672.5 (2235.2 to 3177.1) | 2922.5 (2425.1 to 3504.3) | 44.6 (31.6 to 89.2)* | 9.3 (3.8 to 14.5)* | 40.0 (30.8 to 48.3) | 40.5 (34.0 to 47.8) | 38.8 (32.3 to 46.4) | -3.0 (-12.0 to 26.0) | -4.3 (-9.3 to 0.2) |
| Depressive disorders | 29503.5 (20318.2 to 40109.0) | 39052.4 (26930.5 to 52535.2) | 44208.4 (30573.2 to 59878.5) | 49.8 (46.9 to 53.1)* | 13.2 (12.2 to 14.4)* | 630.6 (438.2 to 852.5) | 620.1 (430.6 to 831.7) | 597.9 (414.5 to 806.2) | -5.2 (-6.2 to -4.1)* | -3.6 (-4.3 to -2.9)* |
| Major depressive disorder | 23423.5 (16150.4 to 31868.1) | 30670.3 (21102.1 to 41462.3) | 34104.6 (23469.5 to 46039.4) | 45.6 (42.4 to 49.2)* | 11.2 (10.1 to 12.3)* | 495.7 (341.2 to 671.8) | 484.6 (334.7 to 655.2) | 461.1 (317.9 to 622.5) | -7.0 (-8.1 to -5.8)* | -4.9 (-5.6 to -4.1)* |
| Dysthymia | 6080.0 (4134.8 to 8839.6) | 8382.0 (5676.3 to 12123.1) | 10103.8 (6860.6 to 14611.5) | 66.2 (62.6 to 69.8)* | 20.5 (18.3 to 23.1)* | 134.9 (91.8 to 195.5) | 135.5 (91.6 to 196.3) | 136.8 (93.0 to 197.6) | 1.4 (0.5 to 2.3)* | 1.0 (-0.5 to 2.7) |
| Bipolar disorder | 5873.0 (3645.7 to 8618.8) | 7795.1 (4869.3 to 11432.4) | 8954.0 (5588.3 to 13186.4) | 52.5 (50.0 to 55.0)* | 14.9 (13.8 to 16.0)* | 118.2 (73.8 to 173.0) | 118.3 (74.1 to 173.1) | 119.3 (74.7 to 175.1) | 0.9 (0.1 to 1.8)* | 0.8 (0.2 to 1.4)* |
| Anxiety disorders | 17893.2 (12472.9 to 24294.6) | 23364.5 (16285.6 to 31632.0) | 26417.4 (18440.4 to 35634.4) | 47.6 (45.4 to 49.8)* | 13.1 (11.9 to 14.3)* | 357.0 (248.8 to 482.4) | 356.6 (248.7 to 481.7) | 354.2 (247.1 to 477.8) | -0.8 (-1.8 to 0.2) | -0.7 (-1.7 to 0.2) |
| Eating disorders | 1400.4 (920.5 to 2016.6) | 1861.9 (1222.9 to 2684.6) | 2180.1 (1420.8 to 3122.9) | 55.7 (52.2 to 59.3)* | 17.1 (15.2 to 18.8)* | 24.6 (16.2 to 35.3) | 26.2 (17.2 to 37.7) | 28.6 (18.6 to 40.8) | 16.1 (14.3 to 17.7)* | 8.9 (7.6 to 10.1)* |
| Anorexia nervosa | 418.9 (263.0 to 626.4) | 523.9 (329.8 to 776.8) | 584.7 (367.2 to 859.0) | 39.6 (36.1 to 43.4)* | 11.6 (9.2 to 14.0)* | 7.1 (4.5 to 10.6) | 7.3 (4.6 to 10.8) | 7.7 (4.8 to 11.3) | 8.1 (5.6 to 10.5)* | 5.4 (3.2 to 7.4)* |
| Bulimia nervosa | 981.5 (602.8 to 1471.9) | 1338.0 (818.2 to 1997.1) | 1595.4 (974.5 to 2394.5) | 62.5 (58.5 to 66.9)* | 19.2 (16.9 to 21.1)* | 17.5 (10.6 to 26.3) | 18.9 (11.5 to 28.4) | 20.9 (12.7 to 31.2) | 19.4 (17.5 to 21.2)* | 10.2 (8.7 to 11.7)* |
| Autistic spectrum disorders | 6525.8 (4418.6 to 9180.8) | 8104.7 (5491.2 to 11393.0) | 9025.7 (6119.0 to 12681.1) | 38.3 (37.3 to 39.4)* | 11.4 (10.8 to 12.0)* | 120.6 (81.9 to 169.6) | 121.1 (82.1 to 170.2) | 121.4 (82.3 to 170.6) | 0.6 (0.0 to 1.2)* | 0.3 (-0.2 to 0.8) |
| Autism | 3371.4 (2170.7 to 4891.2) | 4178.3 (2693.8 to 6041.1) | 4649.0 (2976.6 to 6699.0) | 37.9 (36.3 to 39.6)* | 11.3 (10.3 to 12.3)* | 62.8 (40.4 to 90.9) | 62.6 (40.3 to 90.4) | 62.5 (40.0 to 90.2) | -0.4 (-1.3 to 0.5) | -0.1 (-0.9 to 0.8) |
| Asperger syndrome and other autistic spectrum disorders | 3154.3 (2078.1 to 4617.6) | 3926.4 (2574.5 to 5735.4) | 4376.7 (2864.1 to 6392.9) | 38.8 (37.5 to 40.1)* | 11.5 (10.9 to 12.1)* | 57.8 (38.0 to 84.2) | 58.5 (38.3 to 85.2) | 58.9 (38.5 to 85.9) | 1.8 (1.2 to 2.3)* | 0.6 (0.1 to 1.1)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | Age-standardised DALY rate (per 100 000) | | | | | |
|--|--|--|---|---------------------------------|--|--------------------------------------|--------------------------------------|--------------------------------------|------------------------------|---------------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990-2016 | Percentage change, 2006-16 | 1990 | 2006 | 2016 | Percentage change, 1990-2016 | Percentage change, 2006-16 |
| (Continued from previous page) | | | | | | | | | | |
| Attention-deficit hyperactivity disorder | 599.7 (359.0 to 952.7) | 711.9 (426.0 to 1134.5) | 755.2 (452.2 to 1196.6) | 25.9 (23.5 to 28.3)* | 6.1 (5.0 to 7.1)* | 10.2 (6.1 to 16.1) | 10.1 (6.0 to 16.1) | 10.1 (6.0 to 15.9) | -1.3 (-3.0 to 0.5) | -0.5 (-1.4 to 0.4) |
| Conduct disorder | 5072.5 (3154.4 to 7682.5) | 5820.5 (3615.8 to 8803.8) | 5947.3 (3701.9 to 8998.5) | 17.2 (15.6 to 18.7)* | 2.2 (0.9 to 3.4)* | 79.5 (49.4 to 120.5) | 79.4 (49.4 to 120.6) | 81.2 (50.5 to 122.9) | 2.1 (0.7 to 3.4)* | 2.3 (1.1 to 3.4)* |
| Idiopathic developmental intellectual disability | 3629.4 (1741.7 to 6242.4) | 4500.4 (2160.1 to 7650.1) | 4610.9 (2201.5 to 7932.1) | 27.1 (21.7 to 30.9) | 2.5 (-0.9 to 4.6) | 66.2 (31.8 to 113.6) | 66.1 (31.8 to 112.4) | 61.6 (29.4 to 105.9) | -7.0 (-11.0 to -4.1) | -6.9 (-10.0 to -4.9) |
| Other mental and substance use disorders | 6461.6 (4417.8 to 9255.4) | 8792.3 (5989.8 to 12 580.6) | 10 357.1 (7059.0 to 14 807.2) | 60.3 (59.0 to 61.9)* | 17.8 (17.2 to 18.5)* | 138.5 (94.5 to 197.7) | 138.4 (94.3 to 197.8) | 138.5 (94.4 to 197.7) | 0.0 (-0.5 to 0.5) | 0.1 (-0.4 to 0.5) |
| Diabetes, urogenital, blood, and endocrine diseases | 81 744.6 (71 204.2 to 94 535.9) | 112 751.7 (98 101.9 to 131 367.5) | 133 747.8 (115 976.8 to 155 676.1) | 63.6 (59.4 to 68.0)* | 18.6 (17.4 to 20.0)* | 1882.2 (1642.5 to 2163.3) | 1942.7 (1699.4 to 2247.3) | 1887.6 (1641.9 to 2192.9) | 0.3 (-1.8 to 2.3) | -2.8 (-3.9 to -1.7)* |
| Diabetes mellitus | 27 475.9 (23 282.7 to 32 608.1) | 45 989.7 (38 695.1 to 54 713.4) | 57 233.7 (47 967.9 to 68 279.3) | 108.3 (104.2 to 112.1)* | 24.4 (22.7 to 26.2)* | 708.6 (604.0 to 837.1) | 827.9 (700.5 to 977.9) | 814.2 (686.6 to 965.4) | 14.9 (12.6 to 16.9)* | -1.6 (-3.0 to -0.2)* |
| Acute glomerulonephritis | 560.5 (521.8 to 610.3) | 337.8 (325.7 to 355.6) | 325.6 (310.5 to 343.1) | -41.9 (-46.6 to -37.0)* | -3.6 (-8.4 to 1.5) | 11.6 (10.8 to 12.5) | 5.5 (5.3 to 5.8) | 4.5 (4.3 to 4.8) | -60.9 (-63.9 to -57.6)* | -17.4 (-21.5 to -13.2)* |
| Chronic kidney disease | 21 597.2 (20 094.0 to 23 354.9) | 29 187.2 (27 250.7 to 31 463.0) | 35 032.4 (32 622.1 to 37 954.3) | 62.2 (56.5 to 68.0)* | 20.0 (17.4 to 22.7)* | 521.4 (484.6 to 565.3) | 515.4 (480.9 to 555.7) | 500.1 (465.5 to 541.4) | -4.1 (-7.5 to -1.2)* | -3.0 (-5.0 to -1.0)* |
| Due to diabetes mellitus | 7860.2 (7048.8 to 8711.8) | 11 731.4 (10 615.3 to 12 891.9) | 14 660.6 (13 206.6 to 16 203.7) | 86.5 (77.5 to 92.6)* | 25.0 (21.9 to 27.6)* | 202.9 (182.1 to 225.3) | 212.1 (191.8 to 233.4) | 209.2 (188.8 to 230.8) | 3.1 (-1.4 to 6.1) | -1.4 (-3.5 to 0.5) |
| Due to hypertension | 3481.1 (3030.8 to 3987.5) | 5169.1 (4520.8 to 5845.6) | 6606.7 (5760.2 to 7493.9) | 89.8 (80.4 to 96.3)* | 27.8 (24.7 to 30.6)* | 97.3 (84.5 to 110.8) | 99.2 (87.0 to 112.3) | 98.2 (85.5 to 111.1) | 0.9 (-3.9 to 4.0) | -1.0 (-3.3 to 0.9) |
| Due to glomerulonephritis | 4609.2 (4053.1 to 5203.8) | 5467.8 (4843.2 to 6157.1) | 5932.9 (5226.4 to 6746.0) | 28.7 (22.3 to 36.4)* | 8.5 (5.5 to 11.8)* | 96.8 (85.7 to 109.7) | 89.0 (79.1 to 100.4) | 82.2 (72.8 to 93.3) | -15.1 (-18.2 to -11.3)* | -7.7 (-9.8 to -5.4)* |
| Due to other causes | 5646.7 (5024.1 to 6347.1) | 6819.0 (6061.0 to 7661.1) | 7832.1 (6915.6 to 8848.7) | 38.7 (31.1 to 47.3)* | 14.9 (11.6 to 18.4)* | 124.5 (110.6 to 141.4) | 115.1 (102.5 to 129.8) | 110.6 (97.5 to 125.5) | -11.2 (-14.7 to -7.3)* | -4.0 (-6.2 to -1.4)* |
| Urinary diseases and male infertility | 6451.7 (5595.1 to 7476.6) | 8383.8 (7248.0 to 9800.2) | 9965.5 (8532.5 to 11 725.6) | 54.5 (43.5 to 61.3)* | 18.9 (14.9 to 21.6)* | 152.8 (131.3 to 179.4) | 148.8 (128.2 to 175.0) | 143.5 (122.7 to 169.2) | -6.1 (-11.5 to -2.8)* | -3.6 (-6.5 to -1.5)* |
| Interstitial nephritis and urinary tract infections | 2443.8 (2111.8 to 2752.8) | 3457.4 (3137.6 to 3688.9) | 4269.2 (4005.5 to 4529.9) | 74.7 (54.0 to 95.1)* | 23.5 (15.5 to 30.3)* | 55.6 (48.9 to 61.3) | 60.7 (55.4 to 64.5) | 61.9 (58.2 to 65.6) | 11.3 (-0.3 to 22.0) | 1.9 (-4.2 to 7.3) |
| Urolithiasis | 496.9 (352.4 to 588.4) | 566.9 (491.0 to 658.3) | 622.5 (523.5 to 778.9) | 25.3 (7.1 to 86.6)* | 9.8 (2.6 to 25.5)* | 11.9 (8.6 to 14.1) | 9.7 (8.4 to 11.2) | 8.8 (7.4 to 11.0) | -26.5 (-36.9 to 8.6) | -9.8 (-15.6 to 3.0) |
| Benign prostatic hyperplasia | 1818.5 (1164.2 to 2622.2) | 2625.2 (1692.9 to 3763.4) | 3383.9 (2178.0 to 4835.1) | 86.1 (82.2 to 90.8)* | 28.9 (27.6 to 30.3)* | 50.4 (32.4 to 72.4) | 49.9 (32.2 to 72.1) | 49.2 (31.8 to 71.0) | -2.3 (-4.2 to -0.1)* | -1.3 (-2.2 to -0.3)* |
| Male infertility | 104.7 (42.2 to 207.7) | 138.3 (55.3 to 275.1) | 165.1 (66.1 to 327.1) | 57.7 (52.9 to 62.6)* | 19.3 (15.7 to 22.6)* | 2.0 (0.8 to 3.9) | 2.0 (0.8 to 4.0) | 2.1 (0.9 to 4.3) | 9.5 (6.8 to 12.1)* | 7.9 (4.9 to 10.6)* |
| Other urinary diseases | 1587.8 (1231.6 to 1910.8) | 1596.0 (1346.0 to 1852.1) | 1524.9 (1353.8 to 1722.1) | -4.0 (-17.5 to 18.2) | -4.5 (-12.0 to 7.6) | 32.9 (25.7 to 39.6) | 26.5 (22.4 to 30.7) | 21.4 (19.1 to 24.2) | -34.7 (-44.6 to -20.4)* | -19.0 (-25.5 to -9.1)* |
| Gynaecological diseases | 7068.6 (4854.9 to 10 047.8) | 9365.7 (6439.4 to 13 315.0) | 10 460.2 (7184.1 to 14 928.1) | 48.0 (45.8 to 50.2)* | 11.7 (10.4 to 13.1)* | 145.2 (99.4 to 206.1) | 140.4 (96.1 to 200.0) | 136.8 (94.0 to 195.1) | -5.8 (-7.2 to -4.3)* | -2.5 (-3.5 to -1.4)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|---|---|--|---|---------------------------------------|---------------------------------------|--|--|--|---------------------------------------|---------------------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 |
| (Continued from previous page) | | | | | | | | | | |
| Uterine fibroids | 882.6 (552.9 to 1378.7) | 1237.0 (765.6 to 1961.3) | 1462.5 (897.4 to 2340.2) | 65.7 (59.0 to 71.2)* | 18.2 (16.3 to 19.8)* | 20.1 (12.6 to 31.5) | 19.3 (12.0 to 30.7) | 19.2 (11.8 to 30.7) | -4.6 (-8.6 to -1.5)* | -0.6 (-2.2 to 0.7) |
| Polycystic ovarian syndrome | 98.7 (53.5 to 166.2) | 108.9 (57.8 to 188.7) | 112.4 (59.4 to 198.0) | 14.0 (-4.7 to 29.3) | 3.3 (-4.0 to 8.3) | 1.9 (1.1 to 3.3) | 1.6 (0.8 to 2.7) | 1.5 (0.8 to 2.6) | -24.7 (-39.1 to -12.8)* | -7.3 (-14.5 to -2.3)* |
| Female infertility | 103.8 (37.9 to 234.0) | 129.8 (47.2 to 288.4) | 179.8 (66.2 to 399.0) | 73.2 (58.8 to 91.0)* | 38.5 (29.4 to 49.5)* | 1.9 (0.7 to 4.3) | 1.9 (0.7 to 4.2) | 2.3 (0.9 to 5.2) | 23.2 (13.8 to 35.5)* | 25.3 (17.7 to 34.8)* |
| Endometriosis | 233.8 (156.3 to 325.3) | 310.3 (209.4 to 429.1) | 334.9 (226.3 to 461.8) | 43.2 (36.4 to 50.6)* | 7.9 (5.3 to 10.9)* | 4.4 (3.0 to 6.2) | 4.5 (3.0 to 6.2) | 4.3 (2.9 to 6.0) | -2.3 (-5.9 to 1.4) | -3.8 (-6.1 to -1.4)* |
| Genital prolapse | 586.6 (297.3 to 1051.1) | 694.7 (348.9 to 1246.0) | 792.6 (394.6 to 1424.8) | 35.1 (31.0 to 38.0)* | 14.1 (12.5 to 15.4)* | 14.9 (7.5 to 26.6) | 12.3 (6.2 to 22.0) | 11.1 (5.5 to 19.8) | -25.8 (-27.9 to -24.3)* | -10.0 (-11.2 to -9.0)* |
| Premenstrual syndrome | 2632.6 (1628.3 to 4036.0) | 3470.7 (2146.2 to 5295.9) | 3791.1 (2344.6 to 5812.9) | 44.0 (42.4 to 45.5)* | 9.2 (7.7 to 10.6)* | 50.7 (31.3 to 77.2) | 50.3 (31.1 to 76.7) | 49.3 (30.5 to 75.6) | -2.7 (-3.7 to -1.9)* | -2.0 (-3.3 to -0.8)* |
| Other gynaecological diseases | 2530.6 (1722.7 to 3505.2) | 3414.2 (2325.2 to 4737.6) | 3786.8 (2576.9 to 5232.9) | 49.6 (44.5 to 55.1)* | 10.9 (8.6 to 13.8)* | 51.2 (34.8 to 71.6) | 50.5 (34.4 to 70.1) | 49.2 (33.5 to 67.9) | -4.1 (-6.9 to -0.6)* | -2.7 (-4.7 to -0.1)* |
| Haemoglobinopathies and haemolytic anaemias | 12160.7 (9789.5 to 14856.4) | 12033.2 (9847.5 to 14691.3) | 12321.6 (9916.9 to 15372.2) | 1.3 (-8.3 to 15.8) | 2.4 (-2.6 to 8.3) | 213.4 (174.0 to 258.9) | 181.7 (148.9 to 221.9) | 169.7 (136.9 to 211.4) | -20.5 (-27.2 to -10.5)* | -6.6 (-10.9 to -1.3)* |
| Thalassaemias | 1432.0 (961.6 to 1835.3) | 771.2 (639.7 to 918.5) | 516.0 (443.5 to 633.5) | -64.0 (-70.0 to -40.5)* | -33.1 (-39.9 to -18.8)* | 21.7 (14.6 to 27.8) | 11.2 (9.3 to 13.4) | 7.2 (6.2 to 8.9) | -66.6 (-72.1 to -45.0)* | -35.5 (-42.1 to -21.6)* |
| Thalassaemias trait | 2703.8 (1770.0 to 3974.9) | 2957.1 (1959.0 to 4290.6) | 3280.4 (2156.8 to 4823.8) | 21.3 (17.3 to 25.1)* | 10.9 (8.0 to 13.9)* | 49.7 (32.6 to 73.1) | 44.9 (29.7 to 65.2) | 44.6 (29.3 to 65.5) | -10.2 (-12.1 to -8.4)* | -0.6 (-3.1 to 2.0) |
| Sickle cell disorders | 4349.9 (3163.6 to 5209.6) | 4224.4 (3414.2 to 4951.4) | 4117.9 (3587.8 to 4842.0) | -5.3 (-20.7 to 19.9) | -2.5 (-11.6 to 10.5) | 68.7 (50.4 to 81.8) | 60.8 (49.0 to 71.5) | 56.6 (49.1 to 66.5) | -17.7 (-30.6 to 3.8) | -6.9 (-15.9 to 6.1) |
| Sickle cell trait | 1119.1 (726.0 to 1651.0) | 1331.4 (871.9 to 1959.4) | 1555.0 (1014.1 to 2282.9) | 39.0 (36.2 to 41.7)* | 16.8 (14.8 to 18.8)* | 19.5 (12.8 to 28.8) | 19.7 (12.9 to 28.9) | 21.1 (13.8 to 31.0) | 8.1 (6.2 to 9.7)* | 7.2 (5.4 to 9.1)* |
| G6PD deficiency | 577.8 (488.8 to 691.1) | 702.7 (606.1 to 834.8) | 737.8 (637.7 to 875.0) | 27.7 (18.3 to 42.4)* | 5.0 (1.2 to 10.1)* | 10.9 (9.2 to 13.1) | 10.7 (9.3 to 12.7) | 9.9 (8.6 to 11.8) | -8.8 (-14.7 to 1.2) | -7.3 (-10.7 to -2.8)* |
| G6PD trait | 0.4 (0.3 to 0.6) | 0.5 (0.3 to 0.7) | 0.6 (0.4 to 0.8) | 37.0 (30.4 to 43.2)* | 17.6 (14.9 to 20.7)* | .. | .. | .. | -3.0 (-6.5 to 0.1) | 4.9 (2.6 to 7.5)* |
| Other haemoglobinopathies and haemolytic anaemias | 1977.7 (1591.8 to 2467.5) | 2046.0 (1610.7 to 2612.6) | 2113.9 (1640.0 to 2730.4) | 6.9 (0.9 to 13.2)* | 3.3 (0.6 to 5.8)* | 42.9 (35.4 to 52.5) | 34.4 (27.5 to 43.3) | 30.2 (23.6 to 38.9) | -29.4 (-34.6 to -23.9)* | -12.1 (-15.2 to -9.4)* |
| Endocrine, metabolic, blood, and immune disorders | 6430.0 (5399.9 to 7454.1) | 7454.3 (6326.0 to 8750.0) | 8408.8 (7025.4 to 9979.2) | 30.8 (20.9 to 40.9)* | 12.8 (9.0 to 16.5)* | 129.3 (108.6 to 151.1) | 123.0 (104.1 to 144.5) | 118.7 (99.1 to 140.9) | -8.2 (-13.4 to -3.2)* | -3.5 (-6.4 to -0.8)* |
| Musculoskeletal disorders | 86 655.4 (63 137.9 to 112 703.8) | 117 031.2 (85 442.0 to 152 359.0) | 140 030.6 (102 331.5 to 181 687.7) | 61.6 (59.1 to 63.9)* | 19.6 (18.5 to 20.8)* | 2014.2 (1474.2 to 2609.4) | 1944.7 (1422.3 to 2518.0) | 1918.7 (1404.4 to 2493.3) | -4.7 (-5.7 to -3.9)* | -1.3 (-2.0 to -0.6)* |
| Rheumatoid arthritis | 3329.7 (2458.2 to 4257.7) | 4443.0 (3238.4 to 5693.1) | 5563.4 (3985.9 to 7171.2) | 67.1 (60.8 to 72.7)* | 25.2 (22.4 to 27.6)* | 82.8 (61.6 to 104.8) | 77.7 (56.9 to 99.3) | 78.0 (55.8 to 100.1) | -5.7 (-9.4 to -2.9)* | 0.4 (-1.8 to 2.2) |
| Osteoarthritis | 7947.9 (5593.8 to 10 856.2) | 12 385.2 (8735.1 to 16 806.2) | 16 282.9 (11 486.0 to 22 047.2) | 104.9 (102.9 to 107.2)* | 31.5 (30.8 to 32.2)* | 213.4 (150.2 to 291.8) | 226.6 (159.7 to 308.0) | 232.1 (163.7 to 313.9) | 8.8 (7.6 to 10.2)* | 2.4 (1.9 to 3.0)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|---|---|---|---|--------------------------------|-------------------------------|--|--------------------------------------|--------------------------------------|-----------------------------------|---------------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 |
| (Continued from previous page) | | | | | | | | | | |
| Low back and neck pain | 55 941.4 (39 562.5 to 73 187.3) | 72 599.3 (51 532.0 to 94 807.4) | 86 584.5 (61 335.4 to 113 628.5) | 54.8 (51.8 to 57.4)* | 19.3 (17.7 to 20.7)* | 1294.3 (914.6 to 1702.0) | 1198.9 (855.0 to 1573.7) | 1182.7 (837.4 to 1550.6) | -8.6 (-9.6 to -7.7)* | -1.4 (-2.4 to -0.5)* |
| Low back pain | 39 129.9 (27 512.7 to 51 797.6) | 48 853.4 (34 564.0 to 64 073.6) | 57 648.2 (40 820.5 to 75 877.0) | 47.3 (44.4 to 50.1)* | 18.0 (16.0 to 20.1)* | 897.3 (637.6 to 1178.5) | 805.3 (574.2 to 1053.7) | 788.9 (558.7 to 1034.6) | -12.1 (-12.9 to -11.3)* | -2.0 (-3.6 to -0.9)* |
| Neck pain | 16 811.5 (11 404.8 to 23 792.2) | 23 745.9 (16 214.4 to 33 438.2) | 28 936.3 (19 578.5 to 40 543.1) | 72.1 (68.5 to 76.2)* | 21.9 (19.9 to 24.0)* | 397.0 (268.6 to 556.8) | 393.6 (269.1 to 549.8) | 393.8 (267.8 to 550.2) | -0.8 (-2.2 to 0.7) | 0.1 (-1.3 to 1.4) |
| Gout | 596.8 (413.3 to 810.6) | 848.9 (589.9 to 1155.4) | 1071.2 (742.4 to 1455.0) | 79.5 (76.2 to 82.9)* | 26.2 (24.3 to 27.9)* | 15.4 (10.6 to 21.1) | 15.2 (10.4 to 20.6) | 15.2 (10.4 to 20.6) | -1.8 (-3.3 to -0.2)* | -0.1 (-1.3 to 1.1) |
| Other musculoskeletal disorders | 18 839.6 (13 162.2 to 26 102.7) | 26 754.8 (18 664.4 to 36 839.1) | 30 528.6 (21 196.7 to 42 458.6) | 62.0 (57.4 to 67.0)* | 14.1 (11.4 to 16.8)* | 408.3 (285.4 to 564.2) | 426.3 (298.3 to 586.2) | 410.6 (285.6 to 568.6) | 0.6 (-1.3 to 2.1) | -3.7 (-5.7 to -1.8)* |
| Other non-communicable diseases | 164 624.2 (127 834.6 to 209 136.6) | 183 247.3 (141 607.2 to 236 700.6) | 196 036.1 (148 295.9 to 258 150.6) | 19.1 (6.3 to 31.0)* | 7.0 (2.4 to 10.8)* | 3107.0 (2367.7 to 4005.6) | 2912.8 (2241.2 to 3781.4) | 2758.6 (2096.2 to 3616.8) | -11.2 (-17.7 to -5.3)* | -5.3 (-8.4 to -2.6)* |
| Congenital birth defects | 68 393.0 (53 592.7 to 84 283.8) | 58 286.5 (48 759.3 to 66 905.7) | 50 429.8 (44 114.0 to 56 200.2) | -26.3 (-38.1 to -8.0)* | -13.5 (-20.8 to -5.2)* | 1042.0 (821.5 to 1280.7) | 853.9 (714.0 to 980.0) | 716.0 (626.4 to 798.7) | -31.3 (-42.1 to -15.1)* | -16.1 (-23.1 to -8.3)* |
| Neural tube defects | 10 315.0 (7239.9 to 14 462.5) | 6150.2 (4476.4 to 8763.9) | 5130.9 (3867.1 to 7016.1) | -50.3 (-58.8 to -39.9)* | -16.6 (-27.1 to -5.4)* | 156.9 (111.4 to 218.6) | 90.3 (65.8 to 128.5) | 72.6 (54.5 to 99.5) | -53.8 (-61.3 to -44.5)* | -19.6 (-29.7 to -9.2)* |
| Congenital heart anomalies | 27 581.3 (21 339.9 to 34 673.0) | 22 936.6 (19 833.8 to 26 150.9) | 18 563.8 (16 539.0 to 21 464.8) | -32.7 (-44.5 to -7.0)* | -19.1 (-26.5 to -6.9)* | 413.5 (320.8 to 519.5) | 334.6 (289.1 to 381.8) | 264.7 (235.7 to 305.9) | -36.0 (-47.0 to -11.7)* | -20.9 (-28.1 to -9.0)* |
| Orofacial clefts | 480.8 (242.6 to 688.3) | 320.2 (200.3 to 468.4) | 238.7 (148.0 to 369.6) | -50.4 (-67.5 to -18.4)* | -25.5 (-41.3 to -8.5)* | 7.2 (3.7 to 10.2) | 4.7 (2.9 to 6.8) | 3.4 (2.1 to 5.3) | -52.4 (-68.8 to -22.9)* | -27.0 (-42.6 to -10.0)* |
| Down's syndrome | 1325.4 (767.0 to 3069.7) | 1240.1 (945.4 to 2039.8) | 1166.0 (1005.7 to 1482.8) | -12.0 (-54.8 to 41.0) | -6.0 (-30.4 to 12.8) | 20.7 (12.4 to 46.5) | 18.3 (14.0 to 30.1) | 16.3 (14.0 to 20.7) | -21.4 (-58.2 to 22.5) | -11.2 (-33.7 to 6.2) |
| Turner syndrome | 37.8 (18.1 to 60.9) | 44.2 (21.1 to 71.5) | 47.3 (22.1 to 76.5) | 25.2 (20.6 to 30.2)* | 6.9 (3.2 to 10.4)* | 0.7 (0.3 to 1.1) | 0.6 (0.3 to 1.0) | 0.6 (0.3 to 1.0) | -3.4 (-6.7 to -0.1)* | -1.0 (-4.3 to 2.1) |
| Klinefelter syndrome | 13.0 (6.3 to 24.4) | 15.7 (7.6 to 29.6) | 17.1 (8.2 to 32.3) | 32.1 (28.5 to 35.8)* | 9.0 (6.2 to 11.7)* | 0.2 (0.1 to 0.4) | 0.2 (0.1 to 0.4) | 0.2 (0.1 to 0.4) | -0.3 (-2.6 to 2.3) | -0.1 (-2.5 to 2.5) |
| Other chromosomal abnormalities | 1668.6 (1053.4 to 3061.2) | 1851.6 (1371.3 to 2805.1) | 1952.2 (1542.5 to 2587.9) | 17.0 (-18.1 to 54.2) | 5.4 (-10.1 to 18.4) | 24.9 (15.8 to 45.4) | 26.8 (19.9 to 40.8) | 27.9 (22.0 to 37.1) | 12.2 (-20.7 to 47.3) | 4.0 (-11.4 to 16.7) |
| Congenital musculoskeletal and limb anomalies | 2120.7 (1465.8 to 3497.7) | 2234.9 (1621.0 to 3167.4) | 2258.4 (1661.2 to 2988.8) | 6.5 (-21.7 to 26.8) | 1.1 (-12.2 to 9.4) | 36.6 (25.7 to 57.3) | 33.6 (24.4 to 47.1) | 31.2 (23.0 to 41.5) | -14.6 (-32.7 to -2.5)* | -7.0 (-17.7 to -0.0)* |
| Urogenital congenital anomalies | 1277.1 (845.2 to 1639.7) | 1202.0 (877.2 to 1462.4) | 1085.9 (858.1 to 1305.3) | -15.0 (-35.6 to 10.8) | -9.7 (-19.8 to 3.1) | 19.4 (12.9 to 24.9) | 17.6 (12.8 to 21.4) | 15.4 (12.2 to 18.6) | -20.4 (-39.5 to 2.8) | -12.1 (-21.9 to 0.3) |
| Digestive congenital anomalies | 4666.6 (3171.1 to 8658.0) | 3973.4 (3024.5 to 6520.3) | 3343.0 (2674.6 to 4964.7) | -28.4 (-49.5 to -3.2)* | -15.9 (-27.9 to -1.6)* | 69.9 (47.8 to 128.5) | 58.2 (44.3 to 95.3) | 48.0 (38.3 to 71.5) | -31.3 (-51.3 to -8.1)* | -17.5 (-29.3 to -3.8)* |
| Other congenital birth defects | 18 906.7 (10 942.1 to 30 995.8) | 18 317.5 (12 204.8 to 26 209.6) | 16 626.6 (12 154.3 to 21 604.2) | -12.1 (-33.0 to 18.4) | -9.2 (-20.5 to 3.7) | 291.9 (171.4 to 473.1) | 268.9 (179.1 to 384.4) | 235.6 (172.4 to 307.3) | -19.3 (-37.5 to 5.5) | -12.4 (-22.8 to -0.2)* |
| Skin and subcutaneous diseases | 41 366.2 (28 152.7 to 59 346.6) | 51 550.5 (35 306.4 to 73 530.7) | 57 394.0 (39 334.2 to 81 653.4) | 38.8 (37.0 to 41.0)* | 11.3 (10.6 to 12.4)* | 756.7 (518.9 to 1081.9) | 770.3 (529.3 to 1097.1) | 781.3 (535.8 to 1110.4) | 3.2 (2.5 to 4.4)* | 1.4 (0.8 to 2.2)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|--|------------------------------------|------------------------------------|------------------------------------|------------------------------|----------------------------|--|----------------------------|----------------------------|------------------------------|----------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990-2016 | Percentage change, 2006-16 | 1990 | 2006 | 2016 | Percentage change, 1990-2016 | Percentage change, 2006-16 |
| (Continued from previous page) | | | | | | | | | | |
| Dermatitis | 8427.2 (5021.7 to 13 797.8) | 10 043.3 (6010.8 to 16 327.0) | 11 210.2 (6714.5 to 18 218.1) | 33.0 (31.2 to 35.0)* | 11.6 (10.8 to 12.4)* | 151.8 (91.0 to 246.4) | 151.3 (90.6 to 245.3) | 153.0 (91.6 to 248.4) | 0.7 (-0.3 to 1.8) | 1.1 (0.2 to 1.8)* |
| Psoriasis | 3321.0 (2384.1 to 4347.4) | 4638.9 (3323.7 to 6084.4) | 5643.4 (4039.7 to 7377.2) | 69.9 (68.4 to 71.5)* | 21.7 (20.8 to 22.6)* | 69.7 (50.0 to 91.2) | 73.5 (52.6 to 96.1) | 76.6 (54.9 to 100.0) | 9.8 (9.1 to 10.6)* | 4.2 (3.5 to 5.0)* |
| Cellulitis | 254.1 (179.0 to 320.6) | 451.3 (288.9 to 558.2) | 607.6 (398.4 to 739.6) | 139.1 (103.8 to 168.2)* | 34.6 (25.1 to 48.9)* | 5.5 (3.9 to 6.9) | 7.6 (4.8 to 9.4) | 8.6 (5.6 to 10.4) | 56.3 (32.3 to 74.2)* | 13.1 (5.5 to 24.8)* |
| Pyoderma | 1094.9 (615.2 to 1411.5) | 1577.2 (930.0 to 1948.5) | 1944.8 (1249.8 to 2603.1) | 77.6 (42.6 to 123.0)* | 23.3 (7.9 to 40.5)* | 22.1 (12.9 to 27.9) | 26.0 (15.4 to 32.2) | 27.7 (17.8 to 37.0) | 25.1 (3.6 to 51.2)* | 6.5 (-6.4 to 20.8) |
| Scabies | 3332.5 (1844.4 to 5364.2) | 3670.6 (2034.7 to 5849.4) | 3787.8 (2103.6 to 6029.0) | 13.7 (10.8 to 16.7)* | 3.2 (1.8 to 4.6)* | 58.6 (32.4 to 93.0) | 53.9 (29.8 to 85.6) | 51.0 (28.3 to 81.0) | -13.0 (-13.9 to -12.2)* | -5.4 (-6.2 to -4.8)* |
| Fungal skin diseases | 2267.4 (900.2 to 4720.9) | 2973.4 (1184.9 to 6172.3) | 3508.8 (1403.0 to 7271.0) | 54.8 (51.8 to 57.7)* | 18.0 (17.0 to 19.1)* | 46.1 (18.3 to 95.7) | 47.7 (18.9 to 98.8) | 48.9 (19.5 to 101.4) | 6.0 (5.3 to 6.8)* | 2.5 (2.2 to 2.9)* |
| Viral skin diseases | 4543.4 (2823.8 to 6780.1) | 5425.8 (3369.5 to 8082.0) | 5915.2 (3674.3 to 8828.1) | 30.2 (29.2 to 31.2)* | 9.0 (8.6 to 9.5)* | 80.2 (49.9 to 119.5) | 80.0 (49.8 to 119.1) | 79.9 (49.6 to 119.3) | -0.4 (-0.8 to 0.0) | -0.1 (-0.5 to 0.2) |
| Acne vulgaris | 12 086.8 (8150.7 to 17 552.7) | 15 067.6 (10 169.9 to 21 752.0) | 15 836.0 (10 643.5 to 22 842.6) | 31.0 (29.7 to 32.4)* | 5.1 (4.3 to 5.8)* | 202.3 (136.5 to 292.9) | 207.8 (140.1 to 300.1) | 212.1 (143.0 to 306.4) | 4.9 (4.2 to 5.6)* | 2.1 (1.5 to 2.6)* |
| Alopecia areata | 350.9 (224.8 to 530.5) | 447.0 (286.2 to 675.8) | 504.2 (322.7 to 760.0) | 43.7 (42.1 to 45.3)* | 12.8 (11.8 to 13.9)* | 6.9 (4.4 to 10.4) | 6.8 (4.3 to 10.2) | 6.7 (4.3 to 10.1) | -3.5 (-4.4 to -2.6)* | -1.3 (-2.2 to -0.5)* |
| Pruritus | 452.1 (211.1 to 825.5) | 599.8 (278.6 to 1087.3) | 709.1 (329.7 to 1298.7) | 56.8 (52.5 to 61.0)* | 18.2 (16.7 to 19.8)* | 9.4 (4.4 to 17.2) | 9.6 (4.5 to 17.5) | 9.7 (4.5 to 17.8) | 2.9 (2.4 to 3.5)* | 1.3 (0.8 to 1.8)* |
| Urticaria | 3155.2 (2020.3 to 4556.7) | 3684.4 (2348.7 to 5265.8) | 4029.9 (2575.9 to 5745.3) | 27.7 (24.9 to 31.0)* | 9.4 (8.3 to 10.5)* | 55.1 (35.3 to 78.7) | 55.0 (35.1 to 78.7) | 54.9 (35.1 to 78.6) | -0.5 (-1.1 to 0.1) | -0.3 (-0.8 to 0.3) |
| Decubitus ulcer | 377.1 (291.5 to 475.3) | 553.3 (411.4 to 680.3) | 670.4 (513.0 to 836.1) | 77.8 (65.5 to 86.8)* | 21.2 (17.1 to 27.4)* | 10.7 (8.2 to 13.6) | 10.8 (8.0 to 13.3) | 10.2 (7.8 to 12.6) | -5.1 (-11.6 to 0.4) | -5.7 (-9.0 to -0.4)* |
| Other skin and subcutaneous diseases | 1703.5 (851.6 to 3091.8) | 2417.9 (1203.9 to 4377.1) | 3026.6 (1513.0 to 5474.6) | 77.7 (76.2 to 79.3)* | 25.2 (24.5 to 25.9)* | 38.1 (19.1 to 69.0) | 40.4 (20.2 to 72.9) | 42.2 (21.1 to 76.3) | 10.8 (10.1 to 11.7)* | 4.5 (4.1 to 5.1)* |
| Sense organ diseases | 39 443.2 (27 342.1 to 54 552.7) | 54 826.6 (38 158.5 to 76 040.8) | 66 701.9 (46 534.4 to 92 391.9) | 69.1 (67.3 to 71.0)* | 21.7 (20.7 to 22.5)* | 977.4 (686.1 to 1356.7) | 977.9 (685.8 to 1355.9) | 959.3 (670.2 to 1331.0) | -1.9 (-2.5 to -1.2)* | -1.9 (-2.6 to -1.3)* |
| Glaucoma | 211.5 (142.7 to 292.9) | 339.7 (230.3 to 471.3) | 461.1 (311.4 to 642.1) | 118.0 (114.5 to 121.8)* | 35.7 (33.7 to 38.0)* | 6.3 (4.3 to 8.7) | 6.9 (4.7 to 9.5) | 7.0 (4.8 to 9.8) | 12.1 (10.4 to 14.0)* | 2.1 (0.8 to 3.6)* |
| Cataract | 2698.3 (1924.0 to 3682.2) | 4418.1 (3157.8 to 5992.5) | 5789.0 (4134.6 to 7915.3) | 114.5 (111.4 to 117.9)* | 31.0 (29.4 to 32.6)* | 80.1 (57.2 to 108.9) | 88.3 (63.4 to 119.6) | 88.3 (63.3 to 120.5) | 10.3 (9.0 to 11.7)* | 0.0 (-1.1 to 1.1) |
| Macular degeneration | 174.5 (118.9 to 238.7) | 297.1 (202.1 to 406.9) | 408.6 (277.9 to 555.6) | 134.1 (129.3 to 139.0)* | 37.5 (34.7 to 40.4)* | 5.3 (3.6 to 7.3) | 6.1 (4.2 to 8.4) | 6.3 (4.3 to 8.6) | 17.8 (15.3 to 20.4)* | 2.7 (0.7 to 4.7)* |
| Refraction and accommodation disorders | 10 172.5 (6357.4 to 15 852.4) | 13 035.5 (8163.5 to 20 302.0) | 14 972.5 (9340.6 to 23 361.5) | 47.2 (45.2 to 49.0)* | 14.9 (13.7 to 15.9)* | 230.6 (144.4 to 359.2) | 220.0 (137.9 to 342.1) | 209.1 (130.4 to 326.0) | -9.3 (-10.0 to -8.7)* | -5.0 (-5.8 to -4.3)* |
| Age-related and other hearing loss | 21 193.7 (14 943.8 to 29 581.7) | 29 673.0 (20 837.2 to 41 494.7) | 36 287.5 (25 341.8 to 50 893.6) | 71.2 (67.0 to 75.3)* | 22.3 (20.4 to 24.0)* | 534.7 (379.0 to 745.0) | 533.4 (375.3 to 743.4) | 524.3 (368.0 to 734.0) | -1.9 (-3.5 to -0.7)* | -1.7 (-2.9 to -0.6)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|----------------------------------|--|--|--|---|---------------------------------------|--|--|--|--|--|
| | 1990 | 2006 | 2016 | Percentage change, 1990-2016 | Percentage change, 2006-16 | 1990 | 2006 | 2016 | Percentage change, 1990-2016 | Percentage change, 2006-16 |
| (Continued from previous page) | | | | | | | | | | |
| Other vision loss | 1205.2 (853.2 to 1628.8) | 1776.9 (1256.9 to 2396.0) | 2241.2 (1578.3 to 3016.4) | 86.0 (80.7 to 91.0)* | 26.1 (23.9 to 28.1)* | 30.1 (21.3 to 40.6) | 31.8 (22.4 to 42.9) | 32.0 (22.5 to 43.1) | 6.3 (4.6 to 8.0)* | 0.7 (-0.8 to 2.0) |
| Other sense organ diseases | 3787.5 (2395.1 to 5704.0) | 5286.3 (3336.8 to 7935.4) | 6542.0 (4132.0 to 9828.4) | 72.7 (71.6 to 73.9)* | 23.8 (23.1 to 24.4)* | 90.3 (57.2 to 136.3) | 91.4 (57.9 to 137.9) | 92.3 (58.4 to 139.0) | 2.1 (1.7 to 2.5)* | 0.9 (0.6 to 1.3)* |
| Oral disorders | 11294.7 (6848.2 to 17556.0) | 15552.8 (9435.9 to 24158.8) | 19006.2 (11586.8 to 29629.5) | 68.3 (65.8 to 70.6)* | 22.2 (21.2 to 23.3)* | 270.1 (165.4 to 420.1) | 266.3 (162.9 to 413.0) | 265.7 (162.8 to 413.4) | -1.6 (-2.3 to -0.9)* | -0.2 (-0.7 to 0.2) |
| Caries of deciduous teeth | 120.6 (53.2 to 233.0) | 119.1 (52.5 to 231.1) | 127.2 (55.9 to 249.1) | 5.5 (2.8 to 7.3)* | 6.8 (4.5 to 8.5)* | 1.8 (0.8 to 3.5) | 1.8 (0.8 to 3.4) | 1.8 (0.8 to 3.4) | -3.7 (-5.9 to -2.1)* | 0.2 (-1.9 to 1.8) |
| Caries of permanent teeth | 1295.7 (581.5 to 2521.8) | 1571.3 (699.0 to 3047.3) | 1707.6 (760.1 to 3324.9) | 31.8 (29.7 to 33.9)* | 8.7 (8.0 to 9.4)* | 25.7 (11.4 to 50.0) | 24.0 (10.7 to 46.5) | 23.0 (10.2 to 44.6) | -10.6 (-11.9 to -9.3)* | -4.2 (-4.8 to -3.6)* |
| Periodontal diseases | 2676.0 (1065.8 to 5539.9) | 3893.0 (1550.5 to 8078.4) | 4898.0 (1946.8 to 10208.7) | 83.0 (80.9 to 84.8)* | 25.8 (24.7 to 26.8)* | 64.2 (25.6 to 134.6) | 65.1 (25.9 to 136.6) | 66.6 (26.6 to 139.4) | 3.8 (3.0 to 4.4)* | 2.3 (1.8 to 2.9)* |
| Edentulism and severe tooth loss | 4603.5 (3028.6 to 6528.4) | 6553.4 (4335.5 to 9234.5) | 8338.4 (5467.4 to 11760.4) | 81.1 (79.7 to 82.5)* | 27.2 (26.0 to 28.4)* | 125.7 (82.1 to 177.4) | 122.8 (80.3 to 172.5) | 121.7 (79.6 to 171.1) | -3.2 (-3.6 to -2.8)* | -0.9 (-1.6 to -0.3)* |
| Other oral disorders | 2598.9 (1602.8 to 3894.8) | 3416.0 (2109.2 to 5128.9) | 3935.0 (2427.2 to 5907.9) | 51.4 (50.0 to 52.8) | 15.2 (14.6 to 15.8) | 52.7 (32.6 to 79.0) | 52.7 (32.5 to 79.2) | 52.7 (32.5 to 79.2) | -0.0 (-0.4 to 0.3) | 0.0 (-0.3 to 0.3) |
| Sudden infant death syndrome | 4127.1 (3107.6 to 6179.4) | 3031.0 (2487.9 to 3993.4) | 2504.2 (2015.2 to 3003.1) | -39.3 (-57.8 to -20.1)* | -17.4 (-33.2 to -1.1)* | 60.8 (45.8 to 91.0) | 44.4 (36.4 to 58.4) | 36.4 (29.3 to 43.6) | -40.1 (-58.4 to -21.2)* | -18.0 (-33.7 to -1.9)* |
| Injuries | 259714.5 (242647.1 to 276817.9) | 259792.1 (242234.6 to 280976.8) | 255434.3 (236089.5 to 280689.1) | -1.6 (-6.2 to 3.8) | -1.7 (-4.7 to 1.4) | 4875.4 (4544.8 to 5239.0) | 3976.8 (3692.0 to 4319.2) | 3457.3 (3192.1 to 3803.2) | -29.1 (-32.0 to -25.9)* | -13.1 (-15.6 to -10.6)* |
| Transport injuries | 70430.3 (67012.4 to 74545.4) | 80802.1 (76835.3 to 85315.6) | 78051.8 (73391.8 to 83700.8) | 10.8 (6.4 to 15.9)* | -3.4 (-5.8 to -0.9)* | 1320.1 (1253.5 to 1399.1) | 1218.5 (1155.7 to 1292.0) | 1044.0 (980.2 to 1120.6) | -20.9 (-23.9 to -17.8)* | -14.3 (-16.4 to -12.3)* |
| Road injuries | 64788.4 (61744.3 to 68397.9) | 74299.1 (71021.9 to 78088.0) | 71395.0 (67520.8 to 76128.0) | 10.2 (5.5 to 15.2)* | -3.9 (-6.3 to -1.6)* | 1210.6 (1151.7 to 1279.9) | 1119.1 (1066.5 to 1179.6) | 954.5 (901.8 to 1019.0) | -21.1 (-24.1 to -18.0)* | -14.7 (-16.8 to -12.8)* |
| Pedestrian road injuries | 24717.6 (22795.5 to 28090.8) | 26131.7 (24588.0 to 28158.5) | 24001.5 (22518.2 to 25750.8) | -2.9 (-15.1 to 4.8) | -8.2 (-13.1 to -5.0)* | 459.3 (424.8 to 520.6) | 398.7 (375.0 to 429.0) | 323.3 (303.6 to 346.5) | -29.6 (-38.3 to -24.6)* | -18.9 (-23.2 to -16.2)* |
| Cyclist road injuries | 3390.5 (2919.1 to 3889.2) | 4747.7 (4149.8 to 5488.4) | 4968.2 (4229.6 to 5946.9) | 46.5 (30.4 to 67.8)* | 4.6 (-0.2 to 10.0) | 68.7 (58.8 to 79.8) | 74.3 (64.8 to 86.2) | 67.2 (57.1 to 80.7) | -2.2 (-12.4 to 11.4) | -9.5 (-13.4 to -4.7)* |
| Motorcyclist road injuries | 11467.4 (10510.3 to 12944.1) | 15105.0 (13758.6 to 16396.3) | 14937.5 (13499.8 to 16289.9) | 30.3 (18.2 to 40.1)* | -1.1 (-5.4 to 2.6) | 211.8 (194.4 to 239.6) | 222.6 (202.8 to 242.2) | 197.3 (178.6 to 215.4) | -6.8 (-15.5 to 0.2) | -11.3 (-15.1 to -8.2)* |
| Motor vehicle road injuries | 24319.9 (20644.9 to 27442.2) | 27217.7 (24983.6 to 30653.8) | 26211.8 (24325.0 to 29404.9) | 7.8 (0.7 to 25.4)* | -3.7 (-7.2 to 2.4) | 452.8 (386.1 to 509.4) | 406.3 (372.9 to 455.6) | 349.4 (324.1 to 391.5) | -22.9 (-27.6 to -11.0)* | -14.0 (-17.0 to -8.9)* |
| Other road injuries | 893.0 (714.6 to 1103.6) | 1097.0 (919.3 to 1343.6) | 1275.9 (1026.6 to 1626.3) | 42.9 (23.2 to 73.5)* | 16.3 (9.2 to 23.7)* | 17.9 (14.3 to 22.2) | 17.2 (14.3 to 21.1) | 17.3 (13.9 to 22.1) | -3.3 (-15.0 to 15.3) | 0.8 (-4.9 to 6.7) |
| Other transport injuries | 5641.9 (4761.1 to 6609.6) | 6503.0 (5751.7 to 7477.8) | 6656.8 (5797.7 to 7707.3) | 18.0 (5.2 to 38.4)* | 2.4 (-3.0 to 8.9) | 109.5 (92.6 to 128.2) | 99.4 (87.3 to 114.9) | 89.5 (77.8 to 103.9) | -18.3 (-26.4 to -5.1)* | -10.0 (-14.5 to -4.4)* |
| Unintentional injuries | 121695.5 (109006.9 to 134119.5) | 111226.3 (98814.1 to 125226.5) | 107423.5 (94550.6 to 124015.0) | -11.7 (-18.4 to -4.3)* | -3.4 (-7.1 to 0.1) | 2282.2 (2056.3 to 2530.5) | 1750.0 (1549.2 to 1975.9) | 1486.2 (1306.5 to 1716.5) | -34.9 (-39.0 to -30.6)* | -15.1 (-18.0 to -12.2)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|---|---------------------------------------|---------------------------------------|---------------------------------------|-------------------------------|-------------------------------|--|------------------------------|------------------------------|-------------------------------|-------------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990-2016 | Percentage change, 2006-16 | 1990 | 2006 | 2016 | Percentage change, 1990-2016 | Percentage change, 2006-16 |
| (Continued from previous page) | | | | | | | | | | |
| Falls | 26 607.6 (22 592.9 to 31 276.8) | 31 258.7 (26 220.2 to 37 276.0) | 35 773.8 (29 777.6 to 43 175.7) | 34.5 (22.3 to 44.8)* | 14.4 (9.0 to 18.1)* | 589.1 (498.2 to 700.1) | 530.8 (444.1 to 630.7) | 506.5 (422.1 to 610.0) | -14.0 (-20.6 to -9.2)* | -4.6 (-8.8 to -1.8)* |
| Drowning | 34 779.0 (29 934.2 to 37 903.4) | 22 886.7 (20 014.2 to 24 143.1) | 16 809.0 (15 206.1 to 18 047.4) | -51.7 (-55.6 to -43.2)* | -26.6 (-30.1 to -20.1)* | 567.0 (491.4 to 614.9) | 337.6 (295.4 to 356.2) | 229.2 (207.1 to 246.3) | -59.6 (-62.8 to -53.0)* | -32.1 (-35.4 to -26.1)* |
| Fire, heat, and hot substances | 10 059.5 (7 875.0 to 11 492.7) | 8 822.8 (7 294.7 to 10 004.2) | 8 150.8 (6 763.6 to 9 476.2) | -19.0 (-28.2 to -5.5)* | -7.6 (-13.1 to -0.1)* | 186.1 (150.6 to 212.3) | 135.9 (113.3 to 154.4) | 111.4 (92.4 to 129.4) | -40.2 (-46.0 to -31.7)* | -18.1 (-22.7 to -11.8)* |
| Poisonings | 5 471.1 (3 951.1 to 6 579.7) | 3 747.9 (2 956.8 to 4 213.0) | 3 149.4 (2 379.7 to 3 574.1) | -42.4 (-53.8 to -21.9)* | -16.0 (-25.9 to -2.9)* | 95.1 (69.4 to 112.2) | 56.3 (44.5 to 63.3) | 43.0 (32.5 to 48.9) | -54.8 (-63.8 to -39.4)* | -23.6 (-32.8 to -11.8)* |
| Exposure to mechanical forces | 12 931.3 (11 440.5 to 14 960.9) | 12 360.8 (10 580.5 to 14 181.8) | 11 921.2 (9 836.7 to 14 174.8) | -7.8 (-25.2 to 2.4) | -3.6 (-8.0 to 1.0) | 239.6 (211.3 to 277.2) | 189.7 (162.2 to 219.0) | 162.2 (133.8 to 193.1) | -32.3 (-44.1 to -26.6)* | -14.4 (-18.0 to -11.0)* |
| Unintentional firearm injuries | 1 681.2 (1 345.7 to 1 839.8) | 1 415.5 (1 182.3 to 1 575.2) | 1 351.5 (1 104.3 to 1 513.3) | -19.6 (-26.2 to -12.4)* | -4.5 (-10.9 to 0.7) | 30.9 (24.8 to 34.1) | 21.1 (17.6 to 23.5) | 18.1 (14.8 to 20.3) | -41.5 (-46.1 to -36.4)* | -14.3 (-19.9 to -9.9)* |
| Unintentional suffocation | 3 239.4 (2 677.8 to 4 135.3) | 2 101.3 (1 713.6 to 2 425.9) | 1 757.1 (1 410.8 to 2 041.2) | -45.8 (-59.1 to -34.4)* | -16.4 (-23.6 to -8.8)* | 50.6 (42.1 to 63.9) | 31.2 (25.4 to 36.0) | 24.7 (19.8 to 28.7) | -51.2 (-63.0 to -41.5)* | -20.8 (-27.6 to -13.7)* |
| Other exposure to mechanical forces | 8 010.6 (6 937.0 to 9 470.1) | 8 844.1 (7 355.1 to 10 402.4) | 8 812.6 (7 062.9 to 10 749.0) | 10.0 (-10.2 to 20.3) | -0.4 (-5.0 to 4.2) | 158.1 (135.5 to 187.8) | 137.4 (113.9 to 162.8) | 119.5 (96.0 to 145.9) | -24.4 (-37.5 to -19.1)* | -13.1 (-16.6 to -9.6)* |
| Adverse effects of medical treatment | 5 352.0 (3 741.7 to 6 317.4) | 5 077.1 (4 014.2 to 5 746.0) | 4 991.7 (4 229.9 to 5 575.1) | -6.7 (-17.7 to 15.4) | -1.7 (-8.0 to 7.5) | 99.5 (73.7 to 114.5) | 80.2 (64.1 to 90.4) | 69.7 (59.2 to 77.9) | -30.0 (-36.6 to -18.4)* | -13.2 (-18.2 to -6.2)* |
| Animal contact | 6 346.7 (4 715.4 to 7 403.2) | 6 241.9 (4 797.7 to 7 155.5) | 5 936.0 (4 650.5 to 6 820.3) | -6.5 (-17.4 to 13.4) | -4.9 (-10.6 to 3.5) | 118.2 (89.0 to 138.0) | 95.1 (73.2 to 109.3) | 80.7 (63.2 to 92.7) | -31.7 (-38.8 to -18.1)* | -15.1 (-20.1 to -7.7)* |
| Venomous animal contact | 5 129.9 (3 648.8 to 6 063.6) | 5 170.5 (3 898.3 to 5 979.9) | 4 865.9 (3 689.6 to 5 640.6) | -5.2 (-16.8 to 17.5) | -5.9 (-12.1 to 3.0) | 94.9 (68.3 to 111.9) | 78.4 (59.1 to 90.7) | 66.0 (50.0 to 76.6) | -30.4 (-37.9 to -14.7)* | -15.7 (-21.1 to -7.9)* |
| Non-venomous animal contact | 1 216.9 (949.1 to 1 693.5) | 1 071.3 (835.4 to 1 440.4) | 1 070.1 (837.8 to 1 396.1) | -12.1 (-25.2 to -0.4)* | -0.1 (-7.7 to 7.1) | 23.3 (18.2 to 31.6) | 16.7 (13.0 to 22.4) | 14.7 (11.5 to 19.1) | -37.1 (-44.5 to -30.2)* | -12.3 (-18.3 to -6.4)* |
| Foreign body | 7 078.1 (5 242.3 to 9 108.8) | 6 599.7 (5 467.6 to 7 941.9) | 6 742.8 (5 709.7 to 7 921.3) | -4.7 (-20.0 to 18.0) | 2.2 (-6.6 to 11.3) | 127.2 (98.0 to 159.1) | 102.2 (84.9 to 122.5) | 94.0 (79.8 to 110.3) | -26.1 (-35.1 to -12.1)* | -8.0 (-15.0 to -0.3)* |
| Pulmonary aspiration and foreign body in airway | 5 386.2 (4 058.7 to 7 061.1) | 5 309.7 (4 303.9 to 6 377.5) | 5 327.3 (4 587.3 to 6 136.5) | -1.1 (-17.2 to 20.3) | 0.3 (-8.8 to 11.6) | 96.0 (74.8 to 121.0) | 82.1 (66.9 to 98.0) | 74.6 (64.3 to 86.0) | -22.3 (-31.9 to -8.3)* | -9.1 (-16.7 to 0.6) |
| Foreign body in eyes | 124.2 (63.7 to 213.3) | 146.6 (78.2 to 243.6) | 185.8 (100.2 to 310.1) | 49.6 (41.7 to 57.3)* | 26.8 (24.1 to 30.1)* | 2.6 (1.4 to 4.3) | 2.4 (1.3 to 3.9) | 2.5 (1.4 to 4.2) | -2.5 (-4.3 to -0.8)* | 8.2 (6.3 to 10.3)* |
| Foreign body in other body part | 1 567.7 (948.1 to 2 028.4) | 1 143.5 (869.3 to 1 474.9) | 1 229.7 (929.1 to 1 602.4) | -21.6 (-36.7 to 11.6) | 7.5 (0.6 to 13.4)* | 28.6 (18.2 to 36.8) | 17.7 (13.4 to 22.9) | 16.8 (12.7 to 21.9) | -41.3 (-50.8 to -21.5)* | -5.2 (-10.6 to -0.8)* |
| Environmental heat and cold exposure | 4 613.3 (3 492.7 to 5 757.7) | 4 712.3 (3 611.7 to 5 881.4) | 4 676.7 (3 520.0 to 5 993.7) | 1.4 (-9.2 to 11.5) | -0.8 (-7.9 to 6.3) | 96.6 (74.1 to 121.0) | 75.7 (58.1 to 94.5) | 64.1 (48.3 to 82.0) | -33.6 (-40.8 to -27.6)* | -15.3 (-21.8 to -9.7)* |
| Other unintentional injuries | 8 456.9 (7 319.9 to 10 038.8) | 9 518.6 (8 199.6 to 11 247.0) | 9 272.1 (7 666.7 to 11 405.2) | 9.6 (-6.2 to 20.4) | -2.6 (-7.2 to 1.8) | 163.8 (140.6 to 195.7) | 146.4 (124.8 to 174.5) | 125.5 (103.7 to 154.8) | -23.4 (-33.6 to -17.0)* | -14.3 (-18.0 to -10.8)* |
| Self-harm and interpersonal violence | 5 701.9 (5 222.2 to 6 035.1) | 6 163.0 (5 673.6 to 6 450.9) | 5 871.9 (5 386.4 to 6 280.9) | 3.0 (-3.5 to 11.0) | -4.7 (-8.4 to -0.1)* | 1 091.2 (1 007.5 to 1 153.3) | 919.1 (845.6 to 962.8) | 776.8 (712.6 to 830.8) | -28.8 (-33.2 to -23.4)* | -15.5 (-18.7 to -11.4)* |

(Table 1 continues on next page)

| | All-age DALYs (thousands) | | | | | Age-standardised DALY rate (per 100 000) | | | | |
|---|--|--------------------------------------|--|--------------------------------|----------------------------------|--|---------------------------------|-----------------------------------|----------------------------------|----------------------------------|
| | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 | 1990 | 2006 | 2016 | Percentage change, 1990–2016 | Percentage change, 2006–16 |
| (Continued from previous page) | | | | | | | | | | |
| Self-harm | 36 069.6 (33 180.8 to 38 321.5) | 37 517.7 (34 901.0 to 39 462.7) | 35 149.6 (32 845.1 to 37 938.3) | -2.5 (-8.5 to 6.3) | -6.3 (-10.4 to -0.9)* | 705.3 (650.2 to 753.4) | 565.6 (526.2 to 594.0) | 465.7 (434.7 to 502.5) | -34.0 (-37.9 to -28.2)* | -17.7 (-21.2 to -12.9)* |
| Self-harm by firearm | 2951.5 (2377.2 to 3829.1) | 2875.2 (2348.6 to 3710.6) | 2853.2 (2386.0 to 3589.2) | -3.3 (-13.1 to 13.8) | -0.8 (-7.5 to 9.7) | 57.9 (47.4 to 75.0) | 43.2 (35.6 to 55.7) | 37.8 (31.6 to 47.6) | -34.8 (-41.0 to -24.0)* | -12.6 (-18.4 to -3.9)* |
| Self-harm by other specified means | 33 118.1 (30 211.8 to 35 109.1) | 34 642.5 (32 074.6 to 36 526.4) | 32 296.4 (30 216.7 to 34 924.8) | -2.5 (-8.5 to 6.3) | -6.8 (-11.0 to -1.3)* | 647.4 (590.9 to 686.4) | 522.4 (481.9 to 549.9) | 427.9 (399.9 to 462.6) | -33.9 (-37.9 to -28.1)* | -18.1 (-21.8 to -13.3)* |
| Interpersonal violence | 20 942.3 (17 360.5 to 23 608.2) | 24 113.2 (19 792.0 to 26 579.9) | 23 568.3 (19 646.6 to 26 479.2) | 12.5 (3.6 to 22.4)* | -2.3 (-6.9 to 3.5) | 385.9 (318.6 to 436.9) | 353.5 (290.4 to 391.3) | 311.1 (259.7 to 349.8) | -19.4 (-25.8 to -12.5)* | -12.0 (-16.1 to -7.0)* |
| Physical violence by firearm | 6529.3 (4689.4 to 7918.7) | 8381.5 (5616.3 to 9307.9) | 8720.1 (5844.7 to 9853.4) | 33.5 (15.0 to 48.5)* | 4.0 (-0.7 to 9.0) | 119.1 (84.7 to 144.7) | 120.5 (80.8 to 133.8) | 114.3 (76.6 to 129.1) | -4.1 (-17.5 to 6.7) | -5.2 (-9.6 to -0.6)* |
| Physical violence by sharp object | 5070.8 (3833.9 to 6262.5) | 5886.3 (4626.0 to 7279.7) | 5288.2 (4290.5 to 6922.3) | 4.3 (-8.9 to 25.1) | -10.2 (-16.5 to -0.5)* | 94.5 (71.8 to 117.5) | 86.4 (67.6 to 106.7) | 69.5 (56.4 to 90.9) | -26.5 (-36.0 to -12.1)* | -19.6 (-25.1 to -10.7)* |
| Sexual violence | 1162.6 (774.9 to 1647.4) | 1319.4 (884.1 to 1879.0) | 1365.8 (917.1 to 1946.0) | 17.5 (14.4 to 20.6)* | 3.5 (2.2 to 4.8)* | 21.2 (14.2 to 30.0) | 19.1 (12.9 to 27.3) | 18.1 (12.2 to 25.8) | -14.5 (-15.6 to -13.3)* | -5.5 (-6.2 to -4.9)* |
| Physical violence by other means | 8179.6 (6388.8 to 9519.2) | 8526.0 (7045.4 to 10 132.6) | 8194.1 (7011.8 to 10 058.4) | 0.2 (-15.5 to 20.3) | -3.9 (-12.3 to 4.9) | 151.1 (120.0 to 176.2) | 127.5 (105.6 to 151.6) | 109.3 (93.5 to 133.9) | -27.7 (-38.2 to -14.0)* | -14.3 (-21.6 to -6.5)* |
| Forces of nature, conflict and terrorism, and executions and police conflict | 10 576.7 (8362.5 to 12 866.6) | 6132.8 (4005.8 to 8352.3) | 11 241.1 (8003.1 to 14 753.8) | 6.3 (-25.0 to 46.5) | 83.3 (24.2 to 183.3)* | 181.9 (143.2 to 221.0) | 89.3 (58.2 to 121.9) | 150.3 (107.0 to 197.3) | -17.4 (-41.4 to 14.1) | 68.3 (14.3 to 159.6)* |
| Exposure to forces of nature | 3090.2 (1185.9 to 4943.3) | 946.7 (636.1 to 1276.6) | 617.1 (369.5 to 992.6) | -80.0 (-90.2 to -49.1)* | -34.8 (-50.5 to -15.8)* | 53.5 (20.8 to 85.5) | 14.0 (9.4 to 19.0) | 8.3 (5.0 to 13.4) | -84.5 (-92.3 to -60.9)* | -40.9 (-54.9 to -24.4)* |
| Conflict and terrorism | 6889.0 (5852.1 to 7909.2) | 4824.1 (2855.5 to 6799.0) | 10 326.0 (7176.9 to 13 627.5) | 49.9 (5.2 to 100.3)* | 114.0 (36.2 to 271.3)* | 117.4 (99.2 to 135.3) | 69.9 (41.6 to 98.8) | 138.1 (95.9 to 182.2) | 17.6 (-17.4 to 57.1) | 97.4 (25.7 to 240.9)* |
| Executions and police conflict | 733.5 (292.1 to 958.6) | 418.5 (263.0 to 654.0) | 355.7 (221.1 to 576.5) | -51.5 (-65.3 to -12.8)* | -15.0 (-22.4 to -7.5)* | 13.4 (5.4 to 17.6) | 6.2 (3.9 to 9.9) | 4.7 (2.9 to 7.7) | -64.8 (-74.6 to -37.4)* | -24.2 (-30.2 to -17.8)* |

Data in parentheses are 95% uncertainty intervals. To download the data in this table, please visit the Global Health Data Exchange. --not defined (used for asymptomatic causes, epidemics, and outbreaks). DALYs=disability-adjusted life-years. G6PD=glucose-6-phosphate dehydrogenase. *Percentage changes that are statistically significant.

Table 1: Global all-age DALYs and age-standardised DALY rates in 1990, 2006, and 2016 with mean percentage changes between 1990 and 2016, 2006 and 2016, and 1990 and 2016 for all causes

[26.1–35.4]) from 2006 to 2016. Age-standardised DALY rates from self-harm (decreased by 17.7% [12.9–21.2]) and interpersonal violence (12.0% [7.0–16.1]) have both decreased by more than 10% since 2006. Age-standardised DALY rates resulting from conflict and terrorism increased by 97.4% (25.7–240.9) from 2006 to 2016; this rise was primarily driven by ongoing conflicts in north Africa, the Middle East, and sub-Saharan Africa. This result represents an increase in all-age DALYs of 114.0% (36.2–271.3) from 2006 to 2016.

Global DALY counts in 2016 for causes that were estimated separately for the first time are as follows: alcoholic cardiomyopathy 2.59 million (95% UI

2.06 million to 3.24 million), urogenital congenital anomalies 1.09 million (0.86 million to 1.31 million), congenital musculoskeletal and limb anomalies 2.26 million (1.66 million to 2.99 million), digestive congenital anomalies 3.34 million (2.67 million to 4.96 million), Zika virus disease 5100 (3400–8000), Guinea worm disease 0.87 (0.5–1.35), self-harm by firearm 2.85 million (2.39 million to 3.59 million), sexual violence 1.37 million (0.92 million to 1.95 million), myocarditis 1.37 million (1.12 million to 1.51 million), drug-susceptible tuberculosis 39.9 million (38.1 million to 41.9 million), multidrug-resistant tuberculosis without extensive drug resistance 3.32 million (2.79 million to

For the Global Health Data Exchange see <http://ghdx.healthdata.org/node/311214>

| | 1990, at birth | | | | 2006, at birth | | | | 2016, at birth | | | |
|---------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| Global | 67.57 (67.33-67.77) | 58.42 (55.80-60.77) | 62.70 (62.42-62.99) | 55.38 (53.27-57.31) | 71.44 (71.25-71.65) | 61.73 (58.91-64.21) | 66.35 (66.05-66.63) | 58.51 (56.27-60.55) | 75.33 (74.95-75.64) | 64.91 (61.88-67.54) | 69.79 (69.29-70.22) | 61.42 (59.01-63.58) |
| High SDI | 79.15 (79.05-79.24) | 68.23 (65.08-71.02) | 72.32 (72.21-72.44) | 63.82 (61.37-66.00) | 82.13 (82.06-82.21) | 70.61 (67.28-73.57) | 76.32 (76.23-76.41) | 66.98 (64.32-69.32) | 83.37 (83.18-83.57) | 71.49 (68.03-74.53) | 78.06 (77.81-78.29) | 68.33 (65.60-70.82) |
| High-middle SDI | 73.54 (73.16-73.92) | 63.64 (60.78-66.23) | 66.44 (65.99-66.90) | 58.73 (56.43-60.76) | 76.76 (76.39-77.11) | 66.53 (63.49-69.12) | 69.36 (68.83-69.82) | 61.45 (59.13-63.51) | 79.86 (78.65-80.80) | 68.96 (65.69-71.84) | 73.12 (72.09-74.05) | 64.57 (62.03-67.04) |
| Middle SDI | 68.75 (68.41-69.09) | 60.16 (57.68-62.37) | 64.15 (63.79-64.50) | 57.36 (55.31-59.11) | 73.67 (73.45-73.88) | 64.32 (61.62-66.73) | 68.32 (68.06-68.58) | 60.95 (58.80-62.90) | 77.28 (76.99-77.57) | 67.19 (64.26-69.70) | 71.06 (70.68-71.40) | 63.21 (60.91-65.16) |
| Low-middle SDI | 60.39 (59.94-60.81) | 51.69 (49.11-53.94) | 58.41 (57.92-58.90) | 51.03 (48.92-52.98) | 65.55 (65.17-65.94) | 56.12 (53.36-58.47) | 62.39 (61.93-62.85) | 54.51 (52.29-56.57) | 70.26 (69.74-70.74) | 60.11 (57.16-62.70) | 66.25 (65.64-66.79) | 57.90 (55.43-60.03) |
| Low SDI | 53.71 (53.24-54.20) | 46.18 (43.95-48.11) | 51.14 (50.52-51.75) | 44.45 (42.48-46.30) | 57.62 (57.12-58.15) | 49.75 (47.41-51.86) | 55.98 (55.32-56.59) | 48.81 (46.70-50.76) | 64.10 (63.33-64.82) | 55.44 (52.83-57.79) | 61.63 (60.68-62.52) | 53.86 (51.50-56.11) |
| High income | 79.34 (79.24-79.43) | 68.40 (65.23-71.19) | 72.64 (72.54-72.75) | 64.15 (61.70-66.33) | 82.31 (82.23-82.38) | 70.76 (67.43-73.73) | 76.62 (76.52-76.70) | 67.29 (64.63-69.64) | 83.48 (83.28-83.67) | 71.61 (68.12-74.66) | 78.27 (78.04-78.49) | 68.58 (65.82-71.06) |
| High-income North America | 79.04 (78.94-79.13) | 67.67 (64.44-70.61) | 72.15 (72.03-72.26) | 63.25 (60.75-65.49) | 80.67 (80.58-80.76) | 68.83 (65.45-71.90) | 75.63 (75.54-75.74) | 65.87 (63.11-68.29) | 81.50 (81.28-81.72) | 69.34 (65.86-72.43) | 76.79 (76.54-77.04) | 66.71 (63.87-69.21) |
| Canada | 80.64 (80.33-80.93) | 69.83 (66.72-72.67) | 74.23 (73.91-74.55) | 65.75 (63.26-67.95) | 82.75 (82.46-83.01) | 71.51 (68.28-74.46) | 78.10 (77.84-78.42) | 68.86 (66.17-71.23) | 83.89 (83.49-84.30) | 72.30 (69.00-75.27) | 79.76 (79.30-80.22) | 70.04 (67.23-72.61) |
| Greenland | 66.09 (64.72-67.48) | 57.28 (54.54-59.83) | 57.74 (56.27-59.11) | 51.39 (49.21-53.48) | 68.78 (67.50-70.02) | 59.52 (56.78-62.08) | 65.08 (63.83-66.20) | 57.69 (55.47-59.79) | 72.82 (70.37-75.59) | 62.90 (59.41-66.12) | 67.80 (64.69-70.64) | 60.03 (56.88-63.34) |
| USA | 78.87 (78.77-78.98) | 67.45 (64.20-70.42) | 71.93 (71.81-72.05) | 62.99 (60.48-65.23) | 80.45 (80.35-80.54) | 68.54 (65.16-71.62) | 75.37 (75.26-75.48) | 65.54 (62.79-67.99) | 81.23 (80.99-81.46) | 69.01 (65.50-72.12) | 76.45 (76.19-76.73) | 66.34 (63.49-68.86) |
| Australasia | 79.85 (79.63-80.08) | 68.90 (65.74-71.70) | 73.76 (73.52-74.00) | 64.95 (62.38-67.23) | 83.32 (83.13-83.52) | 71.69 (68.27-74.72) | 78.76 (78.55-78.99) | 69.05 (66.22-71.56) | 84.39 (83.80-84.99) | 72.55 (69.07-75.61) | 80.32 (79.63-81.02) | 70.28 (67.38-72.83) |
| Australia | 80.11 (79.87-80.39) | 69.11 (65.92-71.94) | 73.97 (73.69-74.25) | 65.10 (62.51-67.40) | 83.58 (83.35-83.80) | 71.88 (68.45-74.96) | 78.93 (78.68-79.19) | 69.17 (66.31-71.70) | 84.58 (83.84-85.27) | 72.70 (69.18-75.75) | 80.48 (79.67-81.21) | 70.38 (67.51-72.95) |
| New Zealand | 78.55 (78.18-78.89) | 67.90 (64.78-70.66) | 72.73 (72.38-73.08) | 64.24 (61.75-66.50) | 82.05 (81.73-82.34) | 70.75 (67.40-73.65) | 77.93 (77.60-78.25) | 68.50 (65.75-70.98) | 83.40 (82.25-84.62) | 71.83 (68.56-74.89) | 79.55 (78.27-80.83) | 69.78 (66.84-72.44) |
| High-income Asia Pacific | 80.68 (80.38-80.95) | 70.30 (67.29-73.05) | 74.10 (73.68-74.48) | 65.82 (63.48-67.98) | 84.95 (84.70-85.18) | 73.64 (70.34-76.57) | 78.06 (77.75-78.37) | 68.88 (66.28-71.30) | 86.42 (85.62-87.09) | 74.70 (71.14-77.75) | 80.07 (79.10-80.94) | 70.47 (67.66-73.06) |
| Brunei | 75.39 (73.88-76.73) | 65.61 (62.70-68.40) | 71.68 (70.69-72.82) | 63.24 (60.71-65.65) | 78.90 (77.93-79.65) | 68.72 (65.74-71.51) | 74.64 (73.89-75.45) | 65.84 (63.47-68.18) | 79.48 (77.79-81.73) | 69.34 (66.13-72.26) | 74.56 (72.51-77.34) | 65.84 (62.82-68.69) |
| Japan | 81.81 (81.76-81.85) | 71.31 (68.26-74.00) | 75.91 (75.86-75.96) | 67.45 (65.01-69.60) | 85.57 (85.51-85.64) | 74.20 (70.86-77.10) | 78.76 (78.71-78.81) | 69.51 (66.85-71.84) | 86.94 (86.73-87.16) | 75.10 (71.59-78.08) | 80.83 (80.57-81.08) | 71.11 (68.36-73.60) |
| Singapore | 78.10 (76.47-79.73) | 68.51 (65.44-71.31) | 72.86 (71.33-74.39) | 65.11 (62.52-67.52) | 83.71 (82.39-84.98) | 73.34 (70.20-76.23) | 78.55 (77.18-79.92) | 69.99 (67.44-72.54) | 86.08 (83.92-88.42) | 75.16 (71.85-78.57) | 81.26 (78.75-83.67) | 72.01 (68.76-75.05) |
| South Korea | 76.33 (74.99-77.68) | 66.56 (63.56-69.48) | 67.74 (66.18-69.33) | 60.30 (57.86-62.70) | 82.11 (81.07-83.18) | 71.18 (67.93-74.18) | 75.48 (74.23-76.96) | 66.63 (64.01-69.40) | 84.22 (81.22-87.13) | 72.97 (68.98-76.74) | 77.67 (74.32-81.52) | 68.49 (64.84-72.03) |

(Table 2 continues on next page)

| | 1990, at birth | | | | 2006, at birth | | | | 2016, at birth | | | |
|--------------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Western Europe | 79.50 (79.34-79.65) | 68.49 (65.28-71.34) | 72.94 (72.77-73.11) | 64.52 (62.05-66.70) | 82.79 (82.66-82.92) | 71.24 (67.89-74.22) | 77.24 (77.10-77.39) | 68.07 (65.41-70.45) | 84.10 (83.84-84.38) | 72.27 (68.78-75.35) | 79.21 (78.90-79.49) | 69.69 (66.89-72.13) |
| Andorra | 82.80 (80.33-85.33) | 70.60 (66.75-74.37) | 76.19 (73.96-78.77) | 66.81 (63.64-70.01) | 85.95 (83.76-87.60) | 73.11 (69.15-76.73) | 79.16 (77.43-80.82) | 69.25 (65.99-72.17) | 85.77 (82.96-87.69) | 73.03 (69.09-76.68) | 79.33 (77.41-81.92) | 69.42 (66.07-72.49) |
| Austria | 78.86 (78.61-79.12) | 68.23 (65.16-71.00) | 72.15 (71.87-72.41) | 63.91 (61.49-66.04) | 82.53 (82.30-82.75) | 71.30 (68.06-74.20) | 77.00 (76.74-77.24) | 67.80 (65.05-70.15) | 83.85 (83.20-84.52) | 72.24 (68.88-75.30) | 79.12 (78.50-79.87) | 69.55 (66.69-72.18) |
| Belgium | 79.26 (78.82-79.66) | 68.29 (65.15-71.13) | 72.65 (72.14-73.14) | 64.27 (61.68-66.46) | 82.17 (81.80-82.56) | 70.29 (66.89-73.26) | 76.57 (76.13-77.01) | 67.17 (64.45-69.56) | 83.37 (82.33-84.52) | 71.40 (67.92-74.49) | 78.42 (77.24-79.54) | 68.80 (65.77-71.49) |
| Cyprus | 76.97 (76.52-77.41) | 66.64 (63.65-69.30) | 72.89 (72.36-73.42) | 64.67 (62.30-66.83) | 81.02 (80.57-81.48) | 69.93 (66.72-72.78) | 76.30 (75.77-76.81) | 67.41 (64.75-69.72) | 82.82 (82.27-83.36) | 71.45 (68.14-74.43) | 78.12 (77.46-78.82) | 68.97 (66.28-71.45) |
| Denmark | 77.96 (77.13-78.75) | 67.14 (64.04-70.04) | 72.44 (71.66-73.31) | 64.02 (61.53-66.33) | 80.55 (79.78-81.31) | 69.19 (65.95-72.08) | 76.12 (75.38-76.89) | 66.84 (64.10-69.41) | 82.85 (81.51-84.23) | 70.83 (67.40-74.11) | 78.79 (77.31-80.14) | 68.92 (66.08-71.68) |
| Finland | 79.00 (78.66-79.33) | 67.73 (64.43-70.69) | 70.97 (70.60-71.34) | 62.36 (59.81-64.66) | 82.59 (82.27-82.88) | 70.51 (66.95-73.66) | 75.68 (75.31-76.06) | 66.02 (63.18-68.51) | 84.58 (83.85-85.41) | 72.21 (68.52-75.49) | 78.85 (77.95-79.76) | 68.81 (65.95-71.54) |
| France | 81.22 (80.93-81.50) | 69.93 (66.71-72.84) | 73.09 (72.73-73.46) | 64.82 (62.36-66.98) | 84.09 (83.78-84.36) | 72.51 (69.07-75.52) | 77.11 (76.77-77.52) | 68.28 (65.62-70.59) | 85.39 (84.85-85.99) | 73.42 (69.88-76.59) | 79.20 (78.58-79.84) | 69.93 (67.19-72.37) |
| Germany | 78.53 (77.95-79.07) | 67.63 (64.42-70.49) | 72.04 (71.38-72.70) | 63.62 (61.19-65.80) | 82.13 (81.67-82.62) | 70.57 (67.24-73.61) | 76.67 (76.12-77.27) | 67.43 (64.78-69.89) | 83.34 (82.47-84.29) | 71.52 (68.00-74.73) | 78.47 (77.44-79.52) | 68.90 (66.03-71.51) |
| Greece | 79.54 (79.22-79.86) | 68.72 (65.64-71.45) | 74.48 (74.11-74.88) | 66.03 (63.54-68.20) | 82.64 (82.36-82.92) | 71.22 (67.96-74.12) | 77.11 (76.75-77.46) | 68.17 (65.55-70.47) | 83.50 (82.77-84.25) | 72.02 (68.77-75.06) | 78.43 (77.45-79.46) | 69.22 (66.58-71.70) |
| Iceland | 80.41 (79.80-81.00) | 69.17 (65.92-72.04) | 75.68 (75.14-76.24) | 66.75 (64.15-69.07) | 83.15 (82.53-83.81) | 71.45 (68.04-74.44) | 79.55 (79.05-80.05) | 69.93 (67.10-72.36) | 83.99 (82.92-85.02) | 72.17 (68.76-75.31) | 80.56 (79.65-81.40) | 70.79 (67.97-73.50) |
| Ireland | 77.64 (77.03-78.27) | 67.08 (63.98-69.89) | 72.18 (71.58-72.85) | 63.93 (61.51-66.08) | 81.54 (80.95-82.14) | 70.17 (66.93-73.15) | 76.97 (76.34-77.55) | 67.74 (65.08-70.24) | 83.33 (81.96-84.72) | 71.59 (68.10-74.73) | 78.97 (77.45-80.41) | 69.34 (66.29-72.10) |
| Israel | 77.55 (76.40-78.64) | 67.25 (64.05-70.06) | 74.11 (73.01-75.34) | 65.62 (63.00-68.01) | 81.96 (80.91-83.03) | 70.70 (67.35-73.75) | 77.60 (76.42-78.76) | 68.37 (65.55-71.00) | 84.14 (82.16-85.93) | 72.46 (68.90-75.73) | 80.03 (77.74-82.35) | 70.37 (67.28-73.45) |
| Italy | 80.26 (80.02-80.51) | 69.20 (66.00-72.03) | 73.69 (73.41-73.97) | 65.23 (62.77-67.52) | 83.75 (83.51-83.96) | 72.24 (68.85-75.22) | 78.24 (77.97-78.51) | 69.07 (66.42-71.48) | 84.62 (83.96-85.28) | 72.93 (69.56-76.04) | 79.92 (79.14-80.69) | 70.48 (67.66-72.99) |
| Luxembourg | 78.49 (77.99-78.99) | 67.27 (63.97-70.15) | 71.77 (71.28-72.25) | 63.07 (60.55-65.33) | 82.91 (82.40-83.38) | 70.86 (67.36-73.97) | 77.78 (77.29-78.27) | 67.96 (65.12-70.51) | 83.89 (82.76-85.25) | 71.72 (68.23-74.88) | 80.27 (79.11-81.45) | 70.13 (67.02-72.92) |
| Malta | 77.78 (76.65-78.88) | 67.36 (64.26-70.34) | 73.69 (72.64-74.75) | 65.33 (62.75-67.78) | 81.65 (80.63-82.66) | 70.51 (67.17-73.48) | 77.15 (76.16-78.14) | 68.10 (65.44-70.44) | 83.83 (82.07-85.79) | 72.16 (68.59-75.60) | 79.04 (77.13-81.02) | 69.70 (66.63-72.57) |
| Netherlands | 80.09 (79.62-80.59) | 68.69 (65.39-71.61) | 73.87 (73.37-74.35) | 65.23 (62.77-67.50) | 81.91 (81.47-82.33) | 70.14 (66.76-73.22) | 77.65 (77.23-78.12) | 68.24 (65.57-70.67) | 83.61 (82.62-84.52) | 71.53 (67.93-74.73) | 79.63 (78.57-80.58) | 69.85 (67.02-72.51) |
| Norway | 80.06 (79.58-80.53) | 69.28 (66.16-72.13) | 73.63 (73.13-74.20) | 65.00 (62.42-67.32) | 82.62 (82.16-83.03) | 71.46 (68.27-74.40) | 78.08 (77.63-78.52) | 68.58 (65.80-71.04) | 84.10 (82.90-85.35) | 72.67 (69.39-75.72) | 80.07 (78.99-81.28) | 70.29 (67.42-73.13) |
| Portugal | 77.88 (77.56-78.20) | 67.18 (64.16-70.00) | 70.74 (70.33-71.13) | 62.69 (60.23-64.73) | 82.14 (81.79-82.47) | 70.72 (67.42-73.65) | 75.35 (74.92-75.78) | 66.65 (64.07-68.98) | 84.03 (83.35-84.77) | 72.27 (68.81-75.39) | 77.76 (76.94-78.65) | 68.65 (65.82-71.07) |

(Table 2 continues on next page)

| | 1990, at birth | | | | 2006, at birth | | | | 2016, at birth | | | |
|---|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Spain | 80-63 (80-44-80-83) | 69-85 (66-61-72-64) | 73-59 (73-33-73-86) | 65-44 (63-01-67-60) | 84-03 (83-84-84-19) | 72-75 (69-43-75-70) | 77-60 (77-37-77-82) | 68-86 (66-28-71-16) | 85-59 (85-13-86-05) | 73-98 (70-65-77-05) | 80-28 (79-67-80-83) | 71-20 (68-55-73-64) |
| Sweden | 80-43 (80-06-80-79) | 68-95 (65-54-71-97) | 74-85 (74-47-75-27) | 66-02 (63-36-68-35) | 82-83 (82-53-83-14) | 70-85 (67-42-73-87) | 78-66 (78-30-78-99) | 69-03 (66-24-71-48) | 83-96 (82-63-85-19) | 71-69 (68-22-75-14) | 80-13 (78-83-81-40) | 70-06 (66-89-72-76) |
| Switzerland | 80-87 (80-18-81-52) | 69-16 (65-79-72-25) | 74-01 (73-23-74-76) | 65-09 (62-45-67-47) | 83-77 (83-16-84-33) | 71-23 (67-55-74-56) | 78-98 (78-26-79-69) | 68-95 (66-07-71-59) | 85-23 (82-61-87-65) | 72-86 (68-72-76-47) | 81-01 (78-06-83-61) | 70-96 (67-24-74-23) |
| UK | 78-47 (78-35-78-61) | 67-45 (64-27-70-31) | 72-85 (72-72-72-98) | 64-24 (61-72-66-47) | 81-48 (81-36-81-62) | 69-89 (66-55-72-82) | 77-17 (77-04-77-29) | 67-68 (64-93-70-16) | 82-86 (82-65-83-07) | 70-97 (67-58-74-07) | 78-92 (78-71-79-13) | 69-11 (66-28-71-60) |
| England | 78-69 (78-60-78-78) | 67-57 (64-37-70-43) | 73-12 (73-03-73-21) | 64-43 (61-91-66-67) | 81-72 (81-63-81-81) | 70-01 (66-66-73-02) | 77-47 (77-38-77-57) | 67-89 (65-12-70-36) | 83-11 (82-96-83-25) | 71-12 (67-68-74-16) | 79-24 (79-09-79-38) | 69-35 (66-49-71-87) |
| Northern Ireland | 77-86 (76-97-78-86) | 67-25 (64-19-70-16) | 71-89 (70-92-72-94) | 63-72 (61-27-65-96) | 81-13 (80-33-82-04) | 69-82 (66-50-72-92) | 76-24 (75-24-77-20) | 67-16 (64-47-69-69) | 82-42 (80-83-83-98) | 70-89 (67-48-74-17) | 77-85 (76-04-79-70) | 68-48 (65-52-71-38) |
| Scotland | 76-82 (75-91-77-71) | 66-36 (63-34-69-05) | 70-91 (69-99-71-81) | 62-73 (60-22-65-05) | 79-71 (78-83-80-64) | 68-79 (65-75-71-68) | 74-88 (73-94-75-77) | 65-93 (63-25-68-37) | 81-19 (79-67-82-62) | 69-85 (66-62-73-15) | 76-88 (75-38-78-37) | 67-50 (64-51-70-41) |
| Wales | 78-36 (77-53-79-18) | 67-60 (64-53-70-42) | 72-57 (71-70-73-35) | 64-21 (61-67-66-48) | 81-17 (80-50-81-98) | 70-03 (66-82-73-00) | 76-81 (75-88-77-63) | 67-61 (64-93-70-14) | 82-21 (80-81-83-50) | 70-78 (67-46-73-99) | 77-93 (76-40-79-41) | 68-50 (65-45-71-32) |
| Southern Latin America | 76-13 (75-77-76-51) | 66-19 (63-23-68-81) | 68-97 (68-53-69-40) | 61-19 (58-93-63-23) | 79-48 (79-14-79-86) | 68-97 (65-87-71-78) | 72-79 (72-37-73-22) | 64-40 (62-02-66-62) | 80-88 (79-86-81-84) | 70-09 (66-75-72-91) | 74-36 (73-26-75-41) | 65-72 (63-11-68-16) |
| Argentina | 75-75 (75-29-76-26) | 65-95 (63-01-68-56) | 68-54 (68-00-69-07) | 60-85 (58-59-62-95) | 78-76 (78-29-79-23) | 68-49 (65-47-71-24) | 71-82 (71-27-72-35) | 63-67 (61-27-65-87) | 79-99 (79-07-80-94) | 69-45 (66-22-72-26) | 73-31 (72-28-74-38) | 64-91 (62-40-67-25) |
| Chile | 76-84 (76-09-77-67) | 66-46 (63-38-69-30) | 70-19 (69-28-71-14) | 62-03 (59-52-64-35) | 81-54 (80-87-82-23) | 70-29 (67-00-73-23) | 75-82 (75-07-76-66) | 66-64 (63-96-69-06) | 83-19 (80-15-86-14) | 71-75 (67-58-75-74) | 77-28 (73-98-80-51) | 67-98 (64-16-71-77) |
| Uruguay | 76-50 (76-12-76-91) | 66-84 (63-99-69-27) | 69-10 (68-67-69-52) | 61-72 (59-57-63-65) | 79-58 (79-19-79-94) | 69-39 (66-41-71-92) | 72-02 (71-61-72-46) | 64-20 (61-87-66-23) | 81-08 (80-22-81-94) | 70-49 (67-40-73-37) | 73-45 (72-58-74-37) | 65-28 (62-78-67-50) |
| Central Europe, eastern Europe, and central Asia | 73-62 (73-15-74-08) | 63-68 (60-79-66-31) | 64-56 (63-90-65-19) | 56-76 (54-42-58-84) | 74-26 (73-67-74-80) | 64-48 (61-53-67-03) | 63-87 (63-03-64-67) | 56-45 (54-24-58-54) | 77-22 (75-05-78-94) | 66-83 (63-42-69-95) | 68-17 (66-08-70-23) | 59-99 (57-07-62-92) |
| Eastern Europe | 74-39 (73-62-75-17) | 64-24 (61-30-67-02) | 64-17 (63-08-65-26) | 56-40 (54-03-58-61) | 73-71 (72-78-74-57) | 64-01 (61-00-66-60) | 61-30 (59-98-62-59) | 54-31 (52-14-56-50) | 76-64 (73-16-79-74) | 66-36 (62-34-70-18) | 66-07 (62-56-69-75) | 58-32 (54-75-62-09) |
| Belarus | 75-42 (74-82-76-01) | 65-14 (62-14-67-87) | 65-70 (64-92-66-48) | 57-89 (55-55-60-12) | 75-38 (74-79-75-96) | 65-33 (62-38-67-97) | 63-34 (62-55-64-18) | 56-18 (53-89-58-22) | 78-76 (76-56-81-04) | 67-94 (64-17-71-31) | 68-18 (65-61-70-84) | 60-07 (56-96-63-14) |
| Estonia | 75-26 (74-76-75-78) | 64-68 (61-52-67-39) | 64-84 (64-18-65-52) | 56-91 (54-53-59-04) | 78-54 (78-03-79-08) | 67-67 (64-52-70-51) | 67-68 (66-99-68-32) | 59-59 (57-26-61-84) | 81-79 (80-70-83-50) | 70-24 (66-74-73-49) | 72-96 (71-30-74-75) | 63-79 (60-86-66-60) |
| Latvia | 74-65 (74-03-75-21) | 64-20 (61-13-66-92) | 64-27 (63-36-65-20) | 56-36 (53-96-58-53) | 75-93 (75-33-76-57) | 65-68 (62-67-68-35) | 64-98 (64-20-65-77) | 57-45 (55-16-59-55) | 79-79 (78-21-81-34) | 68-63 (65-36-71-88) | 69-96 (68-13-72-10) | 61-53 (58-62-64-29) |
| Lithuania | 76-37 (75-92-76-81) | 65-66 (62-50-68-43) | 66-40 (65-75-67-05) | 58-20 (55-89-60-35) | 77-25 (76-78-77-69) | 66-55 (63-41-69-39) | 65-08 (64-46-65-73) | 57-47 (55-20-59-56) | 80-35 (79-43-81-33) | 69-08 (65-80-72-07) | 69-74 (68-57-70-95) | 61-15 (58-32-63-59) |
| Moldova | 70-79 (69-73-71-89) | 61-25 (58-46-63-91) | 63-58 (62-30-65-03) | 56-05 (53-64-58-31) | 72-79 (71-81-73-84) | 63-39 (60-49-66-05) | 65-05 (63-70-66-39) | 57-81 (55-42-60-17) | 76-14 (74-59-77-77) | 66-07 (62-90-69-03) | 68-27 (66-56-70-28) | 60-43 (57-66-63-22) |

(Table 2 continues on next page)

| | 1990, at birth | | | | 2006, at birth | | | | 2016, at birth | | | |
|--------------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Russia | 74.30 (73.15-75.53) | 64.17 (61.24-67.09) | 63.68 (62.12-65.35) | 55.93 (53.49-58.36) | 73.46 (72.10-74.72) | 63.81 (60.78-66.50) | 60.62 (58.76-62.48) | 53.67 (51.21-56.11) | 76.24 (71.41-80.77) | 66.09 (61.24-70.37) | 65.39 (60.82-70.72) | 57.78 (53.22-62.52) |
| Ukraine | 74.59 (73.87-75.29) | 64.44 (61.53-67.12) | 65.17 (64.23-66.02) | 57.37 (54.94-59.57) | 73.73 (73.02-74.47) | 64.05 (61.13-66.66) | 62.11 (61.14-63.17) | 55.18 (53.00-57.24) | 76.98 (73.53-80.45) | 66.71 (62.72-70.66) | 67.03 (63.32-71.46) | 59.28 (55.74-63.20) |
| Central Europe | 74.78 (74.56-74.99) | 64.82 (61.91-67.34) | 67.20 (66.93-67.47) | 59.01 (56.59-61.15) | 78.06 (77.86-78.26) | 67.66 (64.65-70.33) | 70.72 (70.48-70.96) | 62.05 (59.48-64.26) | 80.26 (79.81-80.73) | 69.37 (66.15-72.15) | 73.46 (72.92-74.01) | 64.18 (61.51-66.52) |
| Albania | 76.80 (76.16-77.40) | 66.70 (63.79-69.40) | 70.99 (70.33-71.57) | 62.37 (59.85-64.70) | 77.77 (77.25-78.26) | 67.82 (64.84-70.51) | 72.54 (71.95-73.05) | 63.92 (61.30-66.25) | 80.73 (79.37-82.15) | 70.09 (67.00-73.06) | 74.70 (73.12-76.39) | 65.64 (62.73-68.36) |
| Bosnia and Herzegovina | 75.13 (73.76-76.67) | 65.36 (62.38-68.11) | 69.04 (67.54-70.60) | 60.82 (58.20-63.58) | 78.49 (77.21-79.85) | 68.19 (65.07-71.10) | 73.34 (71.84-74.76) | 64.16 (61.33-66.77) | 80.22 (78.35-82.06) | 69.40 (66.06-72.61) | 74.92 (73.02-76.90) | 65.25 (62.10-68.20) |
| Bulgaria | 74.75 (74.36-75.12) | 65.12 (62.33-67.64) | 68.21 (67.75-68.62) | 59.90 (57.43-62.08) | 76.38 (76.03-76.73) | 66.65 (63.83-69.18) | 69.21 (68.75-69.66) | 61.13 (58.74-63.22) | 78.48 (76.47-80.36) | 68.23 (64.85-71.41) | 71.66 (69.55-73.90) | 63.05 (60.16-66.08) |
| Croatia | 76.05 (75.15-76.92) | 66.16 (63.16-68.86) | 68.33 (67.25-69.54) | 60.18 (57.73-62.48) | 78.93 (78.19-79.66) | 68.66 (65.57-71.40) | 71.76 (70.89-72.70) | 63.07 (60.52-65.48) | 80.48 (79.01-82.09) | 69.84 (66.51-72.81) | 74.18 (72.61-76.00) | 64.97 (62.11-67.80) |
| Czech Republic | 75.56 (75.29-75.80) | 65.19 (62.16-67.81) | 68.09 (67.77-68.40) | 59.76 (57.30-61.86) | 79.71 (79.42-79.96) | 68.54 (65.33-71.43) | 73.32 (73.01-73.66) | 63.79 (61.01-66.25) | 81.89 (81.28-82.51) | 70.17 (66.63-73.22) | 76.25 (75.52-76.95) | 65.99 (62.86-68.76) |
| Hungary | 73.86 (73.19-74.59) | 63.70 (60.67-66.39) | 65.29 (64.49-66.12) | 57.37 (55.04-59.60) | 77.51 (76.87-78.17) | 66.99 (64.02-69.86) | 69.15 (68.34-69.98) | 60.82 (58.29-63.12) | 79.08 (77.74-80.34) | 68.25 (65.05-71.40) | 72.24 (70.72-73.89) | 63.29 (60.37-66.06) |
| Macedonia | 72.90 (72.13-73.66) | 63.75 (61.00-66.39) | 68.25 (67.47-69.07) | 60.33 (57.97-62.57) | 75.51 (74.84-76.21) | 66.07 (63.33-68.61) | 70.72 (70.05-71.46) | 62.54 (60.07-64.69) | 77.38 (76.62-78.14) | 67.46 (64.51-70.11) | 72.36 (71.51-73.31) | 63.78 (61.12-66.12) |
| Montenegro | 77.78 (76.60-78.77) | 67.53 (64.55-70.39) | 72.15 (70.99-73.33) | 63.20 (60.35-65.67) | 78.22 (77.57-78.80) | 68.03 (64.91-70.75) | 72.32 (71.56-73.15) | 63.53 (60.90-65.94) | 79.74 (78.74-81.20) | 69.19 (65.84-72.36) | 74.35 (73.14-75.51) | 65.09 (62.29-67.77) |
| Poland | 75.45 (75.00-75.89) | 65.44 (62.51-68.02) | 66.52 (65.99-67.14) | 58.50 (56.18-60.68) | 79.51 (79.12-79.95) | 68.82 (65.68-71.57) | 71.08 (70.52-71.66) | 62.28 (59.68-64.58) | 81.70 (80.62-82.78) | 70.53 (67.24-73.68) | 74.08 (72.83-75.37) | 64.61 (61.82-67.22) |
| Romania | 73.19 (72.61-73.78) | 63.48 (60.52-66.01) | 66.61 (65.86-67.30) | 58.28 (55.79-60.54) | 76.46 (75.89-77.05) | 66.33 (63.40-69.05) | 69.22 (68.56-69.89) | 60.81 (58.29-63.07) | 78.88 (77.68-80.08) | 68.33 (65.15-71.19) | 71.66 (70.14-73.15) | 62.82 (60.01-65.52) |
| Serbia | 75.57 (74.59-76.72) | 65.60 (62.61-68.47) | 69.62 (68.54-70.89) | 61.20 (58.60-63.73) | 76.21 (75.52-76.87) | 66.32 (63.47-68.92) | 70.89 (70.13-71.71) | 62.34 (59.78-64.74) | 78.78 (78.14-79.39) | 68.21 (65.12-70.93) | 73.03 (72.20-73.92) | 63.93 (61.27-66.40) |
| Slovakia | 75.40 (74.92-75.91) | 65.48 (62.55-68.09) | 66.74 (66.12-67.40) | 58.67 (56.21-60.75) | 78.32 (77.86-78.82) | 68.02 (65.01-70.67) | 70.47 (69.87-71.01) | 61.80 (59.17-64.11) | 80.39 (79.07-81.78) | 69.59 (66.48-72.65) | 73.43 (71.87-75.05) | 64.10 (61.10-66.96) |
| Slovenia | 77.54 (76.57-78.53) | 66.71 (63.55-69.70) | 69.54 (68.45-70.81) | 60.68 (57.92-63.14) | 81.53 (80.68-82.48) | 69.77 (66.41-72.88) | 74.22 (73.07-75.34) | 64.26 (61.17-67.11) | 83.82 (82.53-85.29) | 71.49 (67.82-74.79) | 77.82 (76.19-79.51) | 67.16 (63.87-70.44) |
| Central Asia | 70.96 (70.47-71.43) | 61.67 (58.93-64.15) | 63.41 (62.78-64.01) | 56.13 (53.81-58.15) | 71.21 (70.56-71.85) | 62.21 (59.51-64.62) | 63.14 (62.28-63.98) | 56.21 (54.16-58.20) | 75.11 (74.09-76.00) | 65.35 (62.41-68.00) | 67.50 (66.36-68.69) | 59.80 (57.45-62.23) |
| Armenia | 73.65 (72.86-74.55) | 64.21 (61.39-66.67) | 66.84 (65.87-67.83) | 58.93 (56.53-61.07) | 75.86 (75.06-76.62) | 66.28 (63.53-68.81) | 69.04 (68.00-70.04) | 61.17 (58.69-63.40) | 79.32 (77.99-80.71) | 69.06 (65.86-71.81) | 71.97 (70.46-73.52) | 63.59 (60.90-66.11) |
| Azerbaijan | 70.14 (68.89-71.42) | 61.29 (58.55-63.91) | 62.31 (60.84-63.68) | 55.45 (53.04-57.80) | 71.21 (69.75-72.64) | 62.51 (59.64-65.19) | 64.79 (63.25-66.29) | 57.83 (55.48-60.13) | 75.80 (73.51-77.94) | 66.17 (62.90-69.11) | 68.41 (65.71-71.21) | 60.78 (57.79-63.89) |

(Table 2 continues on next page)

| | 1990, at birth | | | | 2006, at birth | | | | 2016, at birth | | | |
|------------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Georgia | 73.69 (72.60-74.77) | 64.60 (61.82-67.19) | 65.44 (64.13-66.83) | 58.19 (55.77-60.33) | 77.15 (76.33-77.91) | 67.50 (64.60-70.01) | 67.75 (66.57-68.93) | 60.39 (57.98-62.58) | 78.83 (76.59-81.37) | 68.78 (65.53-71.92) | 69.07 (66.20-72.20) | 61.32 (58.06-64.46) |
| Kazakhstan | 72.57 (71.28-73.83) | 62.91 (59.93-65.60) | 63.02 (61.42-64.78) | 55.66 (52.95-58.03) | 70.91 (69.54-72.39) | 61.82 (58.83-64.44) | 59.58 (57.80-61.41) | 52.99 (50.58-55.46) | 76.18 (73.87-78.50) | 66.04 (62.70-69.28) | 67.14 (64.13-69.98) | 59.19 (56.00-62.47) |
| Kyrgyzstan | 70.12 (69.47-70.74) | 60.78 (58.04-63.25) | 61.84 (61.08-62.69) | 54.55 (52.38-56.53) | 70.78 (70.19-71.40) | 61.83 (59.15-64.16) | 62.51 (61.72-63.27) | 55.63 (53.46-57.56) | 74.95 (73.75-76.12) | 65.15 (62.22-67.95) | 67.51 (66.07-68.99) | 59.76 (57.08-62.16) |
| Mongolia | 65.20 (63.76-66.84) | 57.09 (54.37-59.60) | 59.84 (58.18-61.22) | 53.18 (50.89-55.48) | 68.78 (67.56-69.88) | 60.40 (57.70-62.78) | 60.23 (58.86-61.76) | 53.67 (51.52-55.83) | 73.04 (71.09-75.10) | 63.80 (60.59-66.78) | 63.65 (61.39-65.84) | 56.38 (53.52-59.08) |
| Tajikistan | 67.42 (65.76-69.13) | 58.79 (56.10-61.55) | 62.44 (60.73-64.21) | 55.20 (52.77-57.69) | 70.35 (68.77-71.80) | 61.69 (58.89-64.26) | 65.68 (64.11-67.36) | 58.31 (55.63-60.71) | 74.34 (72.42-76.08) | 64.95 (61.89-67.79) | 69.41 (67.10-71.52) | 61.41 (58.46-64.17) |
| Turkmenistan | 67.10 (66.05-68.06) | 58.73 (55.99-61.13) | 60.29 (59.01-61.58) | 53.72 (51.34-55.90) | 69.90 (68.65-71.06) | 61.43 (58.65-63.85) | 61.78 (60.32-63.14) | 55.19 (52.77-57.37) | 74.00 (72.71-75.15) | 64.86 (62.01-67.41) | 66.48 (65.09-67.80) | 59.26 (56.77-61.54) |
| Uzbekistan | 71.30 (70.34-72.21) | 61.69 (58.72-64.34) | 65.26 (64.22-66.31) | 57.73 (55.40-60.00) | 70.16 (68.96-71.38) | 61.23 (58.40-63.90) | 64.02 (62.34-65.56) | 57.06 (54.76-59.46) | 73.50 (71.85-75.16) | 63.99 (60.85-66.87) | 67.08 (65.28-68.84) | 59.63 (56.99-62.15) |
| Latin America and Caribbean | 72.68 (72.47-72.88) | 63.19 (60.37-65.62) | 66.44 (66.15-66.70) | 58.85 (56.56-60.86) | 77.02 (76.79-77.24) | 66.87 (63.91-69.49) | 70.87 (70.59-71.12) | 62.65 (60.25-64.80) | 78.89 (78.44-79.29) | 68.50 (65.36-71.17) | 72.75 (72.23-73.22) | 64.24 (61.68-66.39) |
| Central Latin America | 74.27 (74.01-74.52) | 64.94 (62.15-67.38) | 68.10 (67.78-68.43) | 60.43 (58.05-62.41) | 77.98 (77.72-78.24) | 68.23 (65.34-70.75) | 72.07 (71.74-72.39) | 63.97 (61.57-66.13) | 79.44 (78.96-79.94) | 69.42 (66.40-71.99) | 73.58 (72.90-74.21) | 65.21 (62.67-67.40) |
| Colombia | 75.14 (74.65-75.61) | 65.94 (63.10-68.39) | 67.62 (67.02-68.16) | 60.35 (58.03-62.34) | 78.34 (77.93-78.75) | 68.76 (65.91-71.28) | 72.04 (71.54-72.55) | 64.29 (61.90-66.41) | 81.05 (79.95-82.12) | 71.08 (68.06-73.83) | 75.42 (74.17-76.72) | 67.08 (64.50-69.48) |
| Costa Rica | 78.90 (78.33-79.44) | 69.10 (66.16-71.74) | 74.24 (73.67-74.83) | 66.15 (63.73-68.37) | 82.16 (81.63-82.63) | 71.73 (68.60-74.44) | 77.09 (76.50-77.68) | 68.42 (65.76-70.84) | 83.57 (82.65-84.51) | 72.85 (69.62-75.81) | 78.47 (77.41-79.58) | 69.52 (66.66-72.08) |
| El Salvador | 72.47 (71.78-73.20) | 63.57 (60.78-66.00) | 63.87 (62.97-64.87) | 56.75 (54.59-58.80) | 77.36 (76.76-78.00) | 67.83 (64.98-70.32) | 69.32 (68.50-70.20) | 61.47 (59.00-63.74) | 79.07 (77.61-80.53) | 69.30 (66.19-72.15) | 71.46 (69.33-73.45) | 63.36 (60.64-66.09) |
| Guatemala | 66.49 (64.79-68.38) | 58.05 (55.25-60.73) | 62.75 (60.95-65.05) | 55.37 (52.93-58.14) | 73.75 (71.64-75.73) | 64.16 (60.86-67.31) | 66.57 (64.06-69.21) | 58.78 (55.71-61.97) | 76.03 (72.36-79.77) | 66.34 (62.24-70.02) | 69.40 (65.37-73.96) | 61.41 (57.28-65.80) |
| Honduras | 67.23 (65.64-68.91) | 59.20 (56.47-61.74) | 67.35 (63.79-70.22) | 59.91 (56.85-63.19) | 71.72 (68.57-75.52) | 63.09 (59.35-66.67) | 69.97 (65.91-74.24) | 62.44 (58.43-66.60) | 73.71 (70.41-77.70) | 64.78 (60.90-68.63) | 71.64 (67.32-75.82) | 63.86 (59.59-67.72) |
| Mexico | 74.77 (74.46-75.10) | 65.19 (62.44-67.68) | 68.33 (67.95-68.72) | 60.44 (58.05-62.49) | 78.14 (77.79-78.46) | 68.29 (65.38-70.81) | 72.79 (72.42-73.13) | 64.44 (61.96-66.62) | 79.11 (78.65-79.56) | 69.00 (65.92-71.58) | 73.71 (73.20-74.19) | 65.17 (62.59-67.43) |
| Nicaragua | 76.43 (75.87-77.02) | 66.58 (63.66-69.19) | 70.74 (70.04-71.41) | 62.56 (59.98-64.83) | 79.65 (79.10-80.27) | 69.67 (66.69-72.42) | 74.08 (73.42-74.74) | 65.79 (63.30-68.12) | 81.17 (79.21-83.22) | 70.94 (67.63-74.13) | 75.25 (72.85-77.72) | 66.84 (63.52-69.99) |
| Panama | 77.54 (76.65-78.49) | 67.96 (65.04-70.56) | 72.50 (71.40-73.60) | 64.56 (62.18-66.83) | 80.33 (79.47-81.20) | 70.30 (67.20-73.04) | 74.78 (73.66-75.91) | 66.44 (63.89-68.76) | 81.98 (80.38-83.60) | 71.57 (68.22-74.62) | 76.03 (73.97-78.07) | 67.33 (64.33-70.19) |
| Venezuela | 75.41 (74.60-76.29) | 66.21 (63.41-68.75) | 69.98 (68.88-70.90) | 62.40 (59.97-64.62) | 79.05 (78.13-79.89) | 69.26 (66.19-71.84) | 70.98 (69.93-72.11) | 63.25 (60.76-65.44) | 79.79 (77.49-81.84) | 69.87 (66.40-73.26) | 71.30 (68.22-73.83) | 63.46 (60.14-66.42) |
| Andean Latin America | 70.08 (69.45-70.78) | 61.05 (58.27-63.48) | 66.45 (65.71-67.13) | 58.65 (56.38-60.66) | 77.68 (77.01-78.35) | 67.63 (64.50-70.22) | 73.50 (72.83-74.19) | 64.81 (62.21-67.09) | 79.76 (78.33-81.16) | 69.43 (66.22-72.41) | 76.02 (74.60-77.53) | 66.99 (64.01-69.65) |

(Table 2 continues on next page)

| | 1990, at birth | | | | 2006, at birth | | | | 2016, at birth | | | |
|----------------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Bolivia | 62.16 (60.65–63.60) | 54.02 (51.26–56.57) | 60.53 (59.19–61.86) | 53.13 (50.77–55.31) | 71.77 (70.20–73.10) | 62.29 (59.32–65.00) | 69.14 (67.72–70.73) | 60.72 (57.94–63.35) | 74.27 (71.23–76.97) | 64.59 (60.91–67.96) | 72.23 (69.63–75.00) | 63.45 (60.07–66.93) |
| Ecuador | 73.91 (73.34–74.50) | 64.46 (61.68–67.01) | 68.98 (68.32–69.61) | 60.96 (58.50–63.15) | 78.62 (78.06–79.21) | 68.58 (65.52–71.15) | 73.05 (72.29–73.76) | 64.42 (61.86–66.79) | 80.36 (79.36–81.30) | 70.02 (66.82–72.77) | 75.35 (74.11–76.49) | 66.43 (63.73–69.03) |
| Peru | 71.42 (70.42–72.50) | 62.21 (59.36–64.81) | 67.44 (66.30–68.55) | 59.59 (57.17–61.81) | 79.55 (78.55–80.63) | 69.25 (66.06–72.00) | 75.44 (74.35–76.53) | 66.62 (63.82–69.13) | 81.60 (79.22–83.98) | 71.03 (67.47–74.59) | 77.80 (75.33–80.32) | 68.64 (65.32–71.83) |
| Caribbean | 69.50 (68.81–70.13) | 60.44 (57.64–62.88) | 65.93 (65.29–66.61) | 58.28 (55.90–60.50) | 72.40 (71.53–73.27) | 62.90 (60.07–65.48) | 68.67 (67.79–69.54) | 60.57 (58.07–62.88) | 75.35 (74.02–76.61) | 65.39 (62.39–68.26) | 71.00 (69.69–72.33) | 62.58 (59.79–65.00) |
| Antigua and Barbuda | 76.68 (75.47–77.85) | 66.92 (63.82–69.77) | 70.84 (69.63–71.97) | 62.64 (60.04–65.02) | 78.76 (77.61–79.93) | 68.43 (65.23–71.40) | 73.34 (72.15–74.65) | 64.53 (61.68–67.13) | 79.92 (78.32–81.59) | 69.37 (65.99–72.65) | 74.65 (72.67–76.41) | 65.61 (62.64–68.53) |
| The Bahamas | 74.00 (73.00–74.89) | 64.52 (61.51–67.24) | 67.09 (66.00–68.09) | 59.62 (57.31–61.77) | 76.19 (75.39–77.01) | 66.38 (63.53–69.02) | 70.46 (69.68–71.19) | 62.45 (60.10–64.62) | 76.30 (74.58–77.97) | 66.49 (63.31–69.38) | 71.33 (69.70–72.93) | 63.22 (60.52–65.91) |
| Barbados | 76.16 (75.12–77.15) | 66.36 (63.33–69.04) | 70.92 (69.66–71.99) | 62.90 (60.42–65.23) | 78.49 (77.53–79.38) | 68.02 (64.90–70.80) | 73.54 (72.47–74.50) | 64.95 (62.20–67.34) | 78.68 (77.25–80.06) | 68.10 (64.80–71.19) | 74.38 (72.78–75.92) | 65.64 (62.79–68.37) |
| Belize | 73.93 (72.68–75.23) | 64.26 (61.15–67.05) | 69.38 (67.81–70.50) | 61.39 (58.70–63.71) | 73.29 (72.38–74.17) | 63.80 (60.87–66.53) | 67.40 (66.22–68.53) | 59.76 (57.23–62.05) | 74.95 (73.10–76.81) | 65.15 (61.97–68.12) | 69.11 (67.10–71.40) | 61.25 (58.23–64.02) |
| Bermuda | 72.90 (71.87–73.98) | 63.98 (61.26–66.60) | 66.78 (65.67–67.86) | 59.48 (57.02–61.60) | 78.00 (76.91–79.03) | 68.18 (65.20–70.92) | 72.81 (71.81–73.79) | 64.62 (62.21–66.92) | 82.39 (80.69–84.46) | 71.70 (68.30–74.84) | 75.74 (74.10–77.22) | 67.03 (64.10–69.63) |
| Cuba | 76.98 (76.43–77.56) | 66.87 (63.85–69.51) | 72.70 (72.06–73.27) | 64.51 (61.96–66.78) | 79.92 (79.40–80.40) | 69.58 (66.44–72.32) | 75.70 (75.10–76.28) | 67.09 (64.47–69.42) | 81.30 (80.31–82.36) | 70.66 (67.44–73.66) | 76.66 (75.59–77.74) | 67.80 (65.03–70.35) |
| Dominica | 73.68 (72.69–74.70) | 64.65 (61.88–67.09) | 70.44 (69.45–71.54) | 62.42 (59.96–64.72) | 75.98 (74.80–77.05) | 66.47 (63.56–69.05) | 70.83 (69.73–71.92) | 62.68 (60.23–64.98) | 75.84 (73.91–77.37) | 66.21 (63.22–69.11) | 70.19 (67.89–72.38) | 62.08 (59.01–65.02) |
| Dominican Republic | 72.72 (72.03–73.64) | 63.61 (60.92–66.27) | 68.92 (67.91–69.97) | 61.20 (58.76–63.51) | 76.56 (75.18–78.17) | 66.72 (63.43–69.68) | 70.72 (69.57–72.27) | 62.60 (60.06–65.04) | 78.57 (76.90–80.63) | 68.47 (65.35–71.80) | 72.88 (71.17–74.87) | 64.51 (61.65–67.18) |
| Grenada | 73.07 (71.50–74.35) | 63.29 (60.23–66.00) | 68.65 (66.90–70.37) | 60.45 (57.69–63.13) | 73.99 (72.55–75.31) | 64.33 (61.30–67.01) | 68.33 (67.02–69.71) | 60.44 (57.82–62.97) | 74.43 (72.29–76.68) | 64.67 (61.62–67.69) | 68.80 (66.68–71.09) | 60.80 (57.83–63.61) |
| Guyana | 67.79 (66.80–68.87) | 58.86 (56.01–61.35) | 60.94 (59.78–62.10) | 53.86 (51.57–56.07) | 68.30 (67.14–69.35) | 59.16 (56.33–61.63) | 61.77 (60.64–62.99) | 54.48 (52.07–56.63) | 71.17 (69.59–72.89) | 61.60 (58.62–64.52) | 64.67 (62.99–66.41) | 57.01 (54.30–59.41) |
| Haiti | 54.27 (52.55–56.07) | 47.10 (44.54–49.51) | 53.86 (52.03–55.82) | 47.16 (44.50–49.59) | 58.20 (55.79–60.80) | 50.59 (47.70–53.48) | 58.51 (56.00–61.09) | 51.37 (48.35–54.31) | 64.42 (61.40–67.53) | 55.98 (52.71–59.53) | 63.58 (60.17–66.93) | 55.92 (52.30–59.54) |
| Jamaica | 75.36 (74.03–76.62) | 65.33 (62.13–68.20) | 73.44 (72.09–74.70) | 64.70 (61.97–67.25) | 76.36 (75.09–77.66) | 66.06 (62.79–69.00) | 73.28 (71.90–74.66) | 64.54 (61.62–67.13) | 76.79 (74.50–79.15) | 66.43 (63.09–69.99) | 73.06 (70.43–75.53) | 64.33 (61.11–67.43) |
| Puerto Rico | 78.45 (77.86–79.07) | 68.36 (65.33–71.06) | 69.65 (68.82–70.46) | 61.58 (59.12–63.83) | 81.08 (80.48–81.63) | 70.21 (66.94–73.17) | 73.52 (72.69–74.23) | 64.40 (61.61–66.85) | 82.36 (81.27–83.54) | 71.33 (68.04–74.29) | 75.04 (73.74–76.39) | 65.81 (62.94–68.39) |
| Saint Lucia | 74.26 (73.22–75.19) | 64.00 (60.85–66.81) | 69.52 (68.34–70.56) | 60.90 (58.14–63.36) | 77.55 (76.60–78.37) | 67.25 (64.22–70.02) | 71.39 (70.30–72.29) | 62.91 (60.23–65.30) | 79.29 (78.14–80.36) | 68.59 (65.28–71.52) | 73.05 (71.64–74.27) | 64.24 (61.19–66.75) |
| Saint Vincent and the Grenadines | 72.72 (71.69–73.72) | 63.32 (60.36–65.87) | 68.54 (67.17–69.93) | 60.62 (58.00–62.99) | 73.96 (73.29–74.76) | 64.30 (61.47–67.01) | 68.62 (67.69–69.48) | 60.56 (58.17–62.78) | 74.80 (73.67–76.09) | 65.00 (62.00–67.90) | 68.80 (67.42–70.06) | 60.65 (58.05–63.14) |

(Table 2 continues on next page)

| | 1990, at birth | | | | 2006, at birth | | | | 2016, at birth | | | |
|---|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Suriname | 72.53 (71.89–73.22) | 62.93 (59.91–65.65) | 67.75 (66.99–68.41) | 59.79 (57.40–61.92) | 72.46 (71.60–73.26) | 62.79 (59.75–65.34) | 66.70 (65.95–67.45) | 58.89 (56.40–61.02) | 74.37 (73.12–75.41) | 64.32 (61.24–67.12) | 68.41 (67.13–69.69) | 60.24 (57.53–62.67) |
| Trinidad and Tobago | 71.87 (71.25–72.56) | 62.39 (59.64–64.91) | 66.28 (65.59–66.97) | 58.68 (56.31–60.87) | 75.17 (74.41–75.89) | 65.04 (61.98–67.79) | 68.14 (67.33–68.96) | 59.89 (57.46–62.16) | 76.98 (75.63–78.20) | 66.48 (63.13–69.33) | 69.31 (67.78–70.76) | 60.75 (57.91–63.36) |
| Virgin Islands | 76.11 (75.19–77.53) | 66.70 (63.82–69.42) | 69.19 (68.00–70.51) | 61.53 (59.07–63.76) | 77.74 (76.93–78.51) | 67.90 (64.87–70.49) | 70.08 (68.90–71.14) | 62.18 (59.66–64.42) | 78.77 (77.54–80.06) | 68.65 (65.55–71.60) | 70.63 (68.75–72.91) | 62.58 (59.71–65.48) |
| Tropical Latin America | 72.47 (72.03–72.91) | 62.56 (59.56–65.08) | 64.85 (64.34–65.35) | 57.38 (55.02–59.42) | 76.95 (76.51–77.34) | 66.18 (62.94–68.96) | 69.57 (69.10–70.00) | 61.27 (58.86–63.46) | 78.92 (78.42–79.40) | 67.98 (64.72–70.81) | 71.59 (70.97–72.12) | 62.99 (60.40–65.22) |
| Brazil | 72.40 (71.94–72.85) | 62.50 (59.50–65.07) | 64.64 (64.13–65.16) | 57.20 (54.86–59.22) | 77.01 (76.56–77.40) | 66.21 (62.96–69.01) | 69.49 (69.01–69.92) | 61.20 (58.75–63.39) | 79.00 (78.49–79.49) | 68.05 (64.76–70.88) | 71.56 (70.94–72.10) | 62.96 (60.36–65.16) |
| Paraguay | 76.30 (75.51–76.96) | 65.92 (62.83–68.70) | 72.86 (72.16–73.63) | 64.25 (61.74–66.64) | 76.20 (75.30–77.15) | 65.88 (62.68–68.75) | 72.04 (71.31–72.71) | 63.45 (60.88–65.75) | 77.10 (75.69–78.52) | 66.50 (63.13–69.55) | 72.09 (70.29–73.53) | 63.58 (60.67–66.17) |
| Southeast Asia, east Asia, and Oceania | 68.81 (68.39–69.21) | 60.57 (58.22–62.65) | 64.47 (64.04–64.88) | 57.98 (56.05–59.68) | 74.53 (74.29–74.77) | 65.49 (62.86–67.79) | 69.18 (68.89–69.46) | 62.13 (60.08–64.01) | 78.37 (78.08–78.64) | 68.55 (65.74–71.01) | 72.11 (71.76–72.42) | 64.57 (62.36–66.48) |
| East Asia | 69.41 (68.83–69.96) | 61.28 (58.95–63.37) | 65.16 (64.58–65.69) | 58.89 (57.00–60.58) | 75.81 (75.53–76.10) | 66.80 (64.19–69.11) | 70.34 (70.04–70.64) | 63.49 (61.45–65.35) | 79.81 (79.51–80.12) | 69.96 (67.21–72.47) | 73.29 (72.97–73.63) | 65.94 (63.80–67.83) |
| China | 69.20 (68.61–69.76) | 61.11 (58.80–63.20) | 64.98 (64.41–65.53) | 58.75 (56.87–60.44) | 75.81 (75.51–76.11) | 66.83 (64.23–69.14) | 70.34 (70.03–70.65) | 63.50 (61.47–65.37) | 79.93 (79.63–80.24) | 70.06 (67.29–72.55) | 73.36 (73.04–73.69) | 66.01 (63.88–67.91) |
| North Korea | 74.06 (72.75–75.74) | 65.06 (62.20–67.84) | 69.18 (67.60–71.13) | 62.46 (60.03–64.70) | 72.48 (71.28–73.81) | 63.39 (60.56–66.10) | 67.13 (66.02–68.33) | 60.62 (58.62–62.50) | 73.59 (72.47–74.73) | 64.70 (61.97–67.31) | 67.88 (66.67–69.08) | 61.17 (58.94–63.24) |
| Taiwan (Province of China) | 76.89 (76.37–77.41) | 67.05 (64.12–69.73) | 71.51 (70.87–72.12) | 63.93 (61.56–65.98) | 81.18 (80.69–81.66) | 70.66 (67.64–73.26) | 74.93 (74.33–75.59) | 66.97 (64.74–69.04) | 82.84 (81.44–84.41) | 72.06 (68.74–75.22) | 76.68 (74.97–78.54) | 68.36 (65.70–70.89) |
| Southeast Asia | 67.54 (67.07–68.01) | 59.01 (56.53–61.29) | 63.14 (62.68–63.59) | 55.93 (53.88–57.83) | 72.13 (71.76–72.56) | 63.04 (60.38–65.34) | 67.15 (66.61–67.64) | 59.50 (57.41–61.49) | 75.51 (75.03–75.96) | 65.84 (63.04–68.32) | 69.96 (69.31–70.56) | 61.94 (59.60–63.93) |
| Cambodia | 59.25 (58.26–60.38) | 51.14 (48.49–53.52) | 55.50 (54.32–56.71) | 48.59 (46.34–50.71) | 66.38 (65.50–67.29) | 57.46 (54.75–59.88) | 61.11 (59.88–62.24) | 53.61 (51.44–55.67) | 71.60 (70.52–72.72) | 62.14 (59.37–64.72) | 65.72 (64.50–67.01) | 57.97 (55.56–60.13) |
| Indonesia | 64.90 (64.34–65.40) | 56.93 (54.56–59.05) | 62.37 (61.83–62.90) | 55.36 (53.33–57.28) | 70.24 (69.77–70.68) | 61.59 (58.97–63.78) | 67.39 (66.80–68.02) | 59.74 (57.49–61.82) | 73.56 (72.96–74.12) | 64.15 (61.36–66.57) | 69.82 (68.83–70.75) | 61.78 (59.34–63.95) |
| Laos | 54.18 (52.72–55.48) | 47.28 (44.92–49.41) | 50.74 (49.16–52.24) | 44.90 (42.56–47.03) | 63.14 (61.87–64.24) | 55.13 (52.55–57.41) | 58.71 (57.49–59.91) | 51.99 (49.78–54.10) | 69.70 (68.04–71.13) | 60.82 (57.98–63.41) | 64.82 (62.91–66.40) | 57.37 (54.73–59.81) |
| Malaysia | 73.70 (73.40–74.04) | 64.48 (61.72–66.81) | 69.61 (69.27–69.95) | 61.94 (59.67–63.97) | 75.99 (75.74–76.23) | 66.58 (63.75–68.95) | 72.12 (71.80–72.45) | 63.96 (61.61–66.06) | 78.05 (77.50–78.57) | 68.14 (65.17–70.73) | 73.16 (72.41–73.94) | 64.78 (62.37–66.97) |
| Maldives | 62.00 (60.92–63.23) | 53.61 (51.08–56.05) | 62.85 (61.73–64.05) | 55.11 (52.72–57.26) | 77.42 (76.45–78.47) | 66.95 (63.89–69.74) | 75.23 (74.21–76.20) | 66.13 (63.33–68.64) | 81.33 (78.62–83.96) | 70.26 (66.32–73.72) | 77.58 (74.97–80.31) | 68.23 (64.84–71.50) |
| Mauritius | 73.42 (72.89–73.94) | 63.63 (60.79–66.22) | 65.39 (64.86–65.92) | 58.03 (55.89–59.98) | 75.89 (75.31–76.40) | 65.67 (62.62–68.28) | 69.19 (68.61–69.83) | 60.98 (58.61–63.08) | 77.81 (76.00–79.57) | 67.12 (63.74–70.36) | 71.42 (69.59–73.36) | 62.67 (59.53–65.47) |
| Myanmar | 60.93 (59.45–62.34) | 53.09 (50.61–55.57) | 56.26 (54.54–57.98) | 49.56 (47.16–51.97) | 67.28 (65.95–68.50) | 58.84 (56.20–61.27) | 61.28 (58.40–63.10) | 54.21 (51.28–56.61) | 73.35 (72.38–74.34) | 64.11 (61.36–66.65) | 66.73 (65.16–67.87) | 59.04 (56.43–61.32) |

(Table 2 continues on next page)

| | 1990, at birth | | | | 2006, at birth | | | | 2016, at birth | | | |
|--------------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Philippines | 70.92 (70.24-71.61) | 61.69 (58.96-64.14) | 63.45 (62.60-64.36) | 55.80 (53.48-57.96) | 72.07 (71.24-72.76) | 62.73 (59.95-65.22) | 64.58 (63.69-65.46) | 57.07 (54.88-59.12) | 73.87 (72.05-75.85) | 64.47 (61.52-67.57) | 66.64 (64.58-68.74) | 59.07 (56.50-61.71) |
| Sri Lanka | 75.38 (74.55-76.26) | 65.51 (62.68-68.15) | 67.77 (66.67-68.89) | 59.80 (57.30-62.16) | 78.31 (77.17-79.27) | 68.15 (65.22-70.82) | 70.46 (69.17-71.65) | 62.27 (59.85-64.54) | 81.09 (78.67-83.77) | 70.47 (66.87-74.07) | 73.72 (70.53-76.90) | 65.07 (61.57-68.32) |
| Seychelles | 74.15 (73.50-74.82) | 65.14 (62.54-67.54) | 64.48 (63.41-65.58) | 57.71 (55.61-59.61) | 76.20 (75.56-77.25) | 66.77 (63.92-69.39) | 68.64 (67.74-69.55) | 61.10 (58.91-63.25) | 77.40 (76.02-78.87) | 67.72 (64.76-70.58) | 70.25 (68.17-72.03) | 62.41 (59.83-64.93) |
| Thailand | 74.08 (73.42-74.68) | 64.41 (61.59-66.96) | 67.20 (66.41-68.01) | 59.48 (57.24-61.59) | 77.81 (77.23-78.38) | 67.73 (64.85-70.31) | 71.48 (70.70-72.29) | 63.15 (60.71-65.36) | 80.91 (79.63-82.00) | 70.24 (67.11-73.26) | 74.59 (72.93-76.22) | 65.71 (62.91-68.38) |
| Timor-Leste | 57.61 (54.99-60.60) | 50.23 (47.11-53.26) | 58.39 (56.06-60.79) | 50.62 (47.62-53.51) | 69.06 (66.95-71.25) | 60.09 (57.11-63.16) | 67.57 (65.41-69.81) | 58.60 (55.53-61.71) | 73.68 (70.57-76.59) | 64.27 (60.67-68.86) | 71.69 (68.65-74.73) | 62.39 (58.56-65.67) |
| Vietnam | 71.11 (69.45-73.05) | 62.16 (59.16-65.03) | 65.27 (63.23-67.12) | 58.12 (55.61-60.67) | 75.49 (73.94-77.73) | 66.09 (63.17-69.01) | 68.53 (66.58-70.74) | 61.21 (58.75-63.77) | 78.10 (76.59-79.12) | 68.44 (65.57-71.08) | 70.87 (69.14-72.67) | 63.30 (60.72-65.77) |
| Oceania | 60.31 (58.44-62.02) | 52.36 (49.47-54.96) | 56.79 (55.08-58.47) | 50.27 (47.85-52.56) | 61.11 (58.80-63.18) | 53.01 (49.94-55.71) | 58.35 (56.00-60.49) | 51.58 (48.75-54.22) | 63.76 (61.41-66.00) | 55.16 (51.88-58.03) | 60.72 (58.24-62.92) | 53.57 (50.64-56.19) |
| American Samoa | 75.02 (73.75-76.44) | 64.48 (61.29-67.60) | 66.93 (65.44-68.49) | 58.64 (56.07-61.15) | 74.38 (72.61-75.88) | 63.93 (60.72-67.22) | 69.25 (67.72-70.70) | 60.50 (57.64-63.08) | 74.43 (71.97-77.03) | 63.94 (60.47-67.29) | 70.37 (67.82-72.73) | 61.39 (58.12-64.51) |
| Federated States of Micronesia | 65.60 (63.23-67.73) | 57.09 (53.99-59.87) | 61.03 (58.77-63.16) | 54.13 (51.41-56.65) | 67.23 (64.39-69.83) | 58.56 (55.40-61.67) | 64.01 (61.74-66.18) | 56.67 (53.87-59.30) | 67.58 (64.54-70.55) | 58.84 (55.44-62.17) | 63.57 (60.73-66.27) | 56.28 (53.25-59.28) |
| Fiji | 68.04 (63.85-72.17) | 58.99 (54.86-62.98) | 62.73 (59.19-66.20) | 55.32 (51.72-58.77) | 67.12 (65.31-69.36) | 58.07 (54.97-61.03) | 62.61 (60.57-64.97) | 55.23 (52.53-57.84) | 67.67 (63.49-71.89) | 58.63 (54.64-62.73) | 63.27 (59.17-67.99) | 56.00 (52.16-60.22) |
| Guam | 76.44 (75.25-77.71) | 66.30 (63.15-69.05) | 70.34 (68.94-71.64) | 62.61 (60.08-64.88) | 76.69 (75.63-77.75) | 66.43 (63.41-69.36) | 70.14 (68.50-71.30) | 62.27 (59.67-64.55) | 76.09 (74.28-78.08) | 65.75 (62.62-68.84) | 69.05 (66.86-71.35) | 61.20 (58.42-63.78) |
| Kiribati | 60.87 (58.96-62.74) | 52.77 (49.98-55.51) | 55.53 (53.76-57.42) | 49.42 (47.04-51.64) | 63.92 (61.83-65.92) | 55.29 (52.20-58.18) | 56.63 (54.56-58.59) | 50.15 (47.47-52.68) | 65.59 (62.39-68.39) | 56.70 (53.26-60.06) | 58.17 (55.44-61.12) | 51.49 (48.39-54.51) |
| Marshall Islands | 67.99 (66.53-69.39) | 58.54 (55.60-61.38) | 61.47 (59.95-62.84) | 54.23 (51.81-56.48) | 65.11 (63.20-67.11) | 56.14 (53.04-59.02) | 61.00 (59.18-62.90) | 53.70 (51.07-56.26) | 67.26 (64.36-70.12) | 57.87 (54.25-61.23) | 62.75 (60.08-65.45) | 55.07 (52.11-58.20) |
| Northern Mariana Islands | 74.74 (72.01-77.56) | 64.95 (61.34-68.42) | 72.67 (70.19-75.43) | 64.25 (61.24-67.37) | 77.62 (75.38-79.86) | 67.24 (63.57-70.35) | 74.91 (72.81-76.87) | 65.99 (62.86-68.76) | 77.37 (74.86-79.95) | 66.90 (63.25-70.24) | 73.88 (71.52-76.20) | 65.02 (62.01-67.86) |
| Papua New Guinea | 57.55 (55.23-59.80) | 49.98 (47.04-52.65) | 54.45 (52.17-56.65) | 48.22 (45.54-50.61) | 58.93 (55.93-61.58) | 51.16 (47.90-54.17) | 56.57 (53.58-59.29) | 50.03 (46.91-52.97) | 62.17 (59.37-64.96) | 53.81 (50.34-56.94) | 59.50 (56.53-62.33) | 52.50 (49.24-55.40) |
| Samoa | 72.32 (70.28-74.30) | 62.82 (59.73-65.93) | 65.84 (63.47-67.86) | 58.20 (55.27-60.85) | 73.62 (71.80-75.57) | 63.79 (60.62-66.81) | 68.98 (67.00-70.99) | 60.94 (58.20-63.63) | 74.00 (71.99-76.12) | 64.09 (60.85-67.24) | 69.90 (67.94-72.24) | 61.73 (58.95-64.39) |
| Solomon Islands | 61.76 (59.54-64.33) | 53.80 (50.92-56.89) | 58.55 (56.42-61.01) | 52.15 (49.75-54.77) | 62.15 (59.76-64.69) | 54.07 (51.07-56.93) | 60.06 (57.82-62.35) | 53.36 (50.66-55.92) | 64.24 (61.49-66.86) | 55.79 (52.73-58.68) | 61.91 (59.31-64.32) | 54.85 (52.12-57.46) |
| Tonga | 70.12 (67.56-72.67) | 60.50 (57.17-63.85) | 65.71 (63.10-68.33) | 58.15 (55.13-61.11) | 72.43 (70.40-74.34) | 62.32 (58.91-65.47) | 66.85 (64.60-69.05) | 59.07 (56.15-61.55) | 73.33 (70.85-75.64) | 63.03 (59.33-66.31) | 67.50 (65.23-70.07) | 59.46 (56.61-62.26) |
| Vanuatu | 63.86 (61.77-66.01) | 55.35 (52.25-58.43) | 59.86 (57.86-61.86) | 53.15 (50.44-55.83) | 63.97 (61.41-66.27) | 55.50 (52.38-58.48) | 60.75 (58.57-62.89) | 54.04 (51.41-56.65) | 65.68 (62.77-68.20) | 56.81 (53.52-60.12) | 62.22 (59.66-64.66) | 55.23 (52.22-58.05) |

(Table 2 continues on next page)

| | 1990, at birth | | | | 2006, at birth | | | | 2016, at birth | | | |
|-------------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| North Africa and Middle East | 68-93 (68-42-69-38) | 57-92 (54-77-60-69) | 65-01 (64-43-65-60) | 56-14 (53-49-58-51) | 73-19 (72-67-73-67) | 61-72 (58-38-64-77) | 69-13 (68-46-69-80) | 59-90 (57-22-62-30) | 75-59 (74-89-76-29) | 63-66 (60-17-66-71) | 70-90 (70-05-71-78) | 61-41 (58-64-63-99) |
| Afghanistan | 51-35 (49-83-52-92) | 42-97 (40-24-45-55) | 51-87 (50-14-53-90) | 44-20 (41-28-46-73) | 54-51 (53-00-55-97) | 46-03 (43-31-48-59) | 53-80 (51-93-55-70) | 46-34 (43-59-49-01) | 59-16 (57-45-60-78) | 49-95 (46-99-52-73) | 56-83 (54-45-59-11) | 49-11 (46-21-51-87) |
| Algeria | 72-56 (71-34-73-76) | 61-07 (57-66-64-19) | 69-44 (68-19-70-72) | 60-04 (57-20-62-71) | 76-85 (75-62-77-68) | 64-85 (61-39-68-08) | 74-42 (73-29-75-58) | 64-42 (61-43-67-15) | 78-41 (77-46-79-34) | 66-11 (62-60-69-36) | 76-43 (75-06-77-66) | 66-04 (62-82-68-90) |
| Bahrain | 70-99 (69-28-72-59) | 59-71 (56-49-63-00) | 68-54 (66-94-70-18) | 59-52 (56-73-62-36) | 74-94 (73-50-76-48) | 62-73 (59-17-66-14) | 72-76 (71-18-74-39) | 62-85 (59-79-65-88) | 77-81 (75-72-80-15) | 65-01 (61-16-68-83) | 75-99 (73-45-78-51) | 65-52 (61-84-68-99) |
| Egypt | 67-29 (66-32-68-21) | 56-84 (53-88-59-56) | 63-32 (62-42-64-29) | 55-04 (52-44-57-31) | 72-89 (72-08-73-70) | 61-63 (58-41-64-61) | 67-66 (66-77-68-54) | 59-00 (56-40-61-49) | 74-98 (73-15-76-99) | 63-54 (60-05-66-80) | 69-44 (67-60-71-52) | 60-64 (57-89-63-40) |
| Iran | 71-07 (68-91-73-37) | 59-86 (56-10-63-19) | 65-24 (62-58-68-12) | 56-37 (53-08-59-74) | 75-20 (72-96-77-42) | 63-59 (59-94-66-96) | 70-75 (68-00-73-35) | 61-37 (57-97-64-66) | 78-38 (76-58-80-58) | 66-03 (62-27-69-57) | 73-78 (71-23-76-22) | 63-72 (60-16-66-85) |
| Iraq | 67-58 (65-61-69-71) | 56-63 (53-29-59-79) | 64-75 (62-05-67-37) | 55-19 (51-77-58-30) | 66-88 (64-23-69-42) | 56-47 (52-82-59-86) | 60-90 (57-38-65-04) | 52-54 (48-76-56-43) | 70-48 (67-37-73-54) | 59-26 (55-41-62-99) | 64-83 (61-47-69-19) | 55-76 (52-01-59-74) |
| Jordan | 72-58 (70-91-74-84) | 60-94 (57-41-64-21) | 71-03 (69-24-72-57) | 61-11 (58-15-63-85) | 74-23 (71-88-76-82) | 62-75 (59-00-66-31) | 73-34 (70-65-75-64) | 63-51 (60-19-66-80) | 77-99 (74-85-81-02) | 65-70 (61-49-69-46) | 74-71 (71-39-78-25) | 64-59 (60-97-68-43) |
| Kuwait | 76-70 (74-82-78-73) | 65-00 (61-29-68-33) | 74-54 (72-38-76-73) | 64-65 (61-43-67-77) | 77-12 (75-72-78-44) | 65-38 (61-84-68-56) | 76-93 (75-01-78-65) | 66-54 (63-23-69-57) | 79-47 (76-63-82-70) | 67-37 (63-50-71-31) | 79-96 (76-38-83-60) | 69-01 (65-03-73-34) |
| Lebanon | 71-92 (70-05-74-03) | 60-87 (57-53-64-06) | 66-38 (64-58-68-79) | 57-36 (54-50-60-39) | 78-72 (77-82-79-96) | 66-52 (62-98-69-83) | 77-22 (75-61-78-85) | 66-63 (63-32-69-56) | 81-42 (79-91-82-87) | 68-66 (64-94-72-23) | 78-79 (77-11-80-32) | 67-94 (64-59-70-98) |
| Libya | 74-68 (73-33-75-82) | 62-95 (59-53-66-07) | 71-59 (70-34-72-98) | 61-97 (59-12-64-75) | 77-01 (75-27-78-13) | 65-03 (61-54-68-16) | 73-87 (72-63-75-29) | 64-02 (60-99-66-73) | 77-59 (76-21-78-74) | 65-47 (61-78-68-74) | 72-62 (71-01-74-65) | 63-00 (59-92-65-82) |
| Morocco | 68-70 (67-60-69-73) | 57-76 (54-46-60-75) | 65-69 (64-85-66-57) | 56-74 (54-04-59-15) | 73-80 (72-47-74-96) | 62-22 (58-80-65-42) | 71-16 (70-12-72-08) | 61-48 (58-51-64-21) | 76-44 (75-14-77-71) | 64-38 (60-74-67-77) | 73-55 (72-48-74-54) | 63-58 (60-60-66-20) |
| Palestine | 71-78 (69-66-73-66) | 61-01 (57-61-64-21) | 70-13 (68-34-72-03) | 60-77 (57-75-63-60) | 74-19 (73-77-74-60) | 63-12 (59-87-65-98) | 69-54 (69-07-69-99) | 60-50 (57-83-62-89) | 73-48 (72-75-74-22) | 62-56 (59-49-65-38) | 70-23 (69-47-71-02) | 61-05 (58-38-63-44) |
| Oman | 71-89 (70-46-73-25) | 60-59 (57-11-63-71) | 69-50 (68-12-70-81) | 59-95 (56-92-62-74) | 78-21 (77-27-79-33) | 66-25 (62-72-69-47) | 74-05 (72-92-75-15) | 64-33 (61-45-67-07) | 79-93 (79-21-80-83) | 67-64 (63-95-70-96) | 75-28 (74-48-76-27) | 65-29 (62-36-68-01) |
| Qatar | 75-25 (73-06-77-85) | 63-35 (59-66-67-15) | 74-82 (72-23-77-11) | 64-23 (60-77-67-49) | 78-98 (77-27-80-91) | 66-20 (62-32-69-81) | 75-74 (73-25-78-54) | 65-28 (61-77-68-76) | 81-82 (78-89-85-16) | 68-54 (64-22-72-79) | 79-00 (75-41-82-65) | 67-94 (63-77-72-07) |
| Saudi Arabia | 73-52 (72-07-74-90) | 61-62 (58-14-64-84) | 70-62 (69-03-72-20) | 61-15 (58-17-63-96) | 74-85 (74-23-75-41) | 63-49 (60-33-66-49) | 73-41 (72-66-74-13) | 64-07 (61-36-66-59) | 78-67 (77-83-79-62) | 66-61 (63-09-69-87) | 75-97 (74-95-77-00) | 66-21 (63-34-68-81) |
| Sudan | 59-53 (58-32-60-64) | 50-03 (47-25-52-75) | 57-60 (56-49-58-74) | 49-47 (46-80-51-85) | 66-70 (65-69-67-70) | 56-23 (53-08-58-98) | 63-67 (62-69-64-72) | 54-90 (52-20-57-27) | 70-29 (68-93-71-52) | 59-15 (55-70-62-35) | 66-37 (64-87-67-67) | 57-23 (54-39-60-00) |
| Syria | 72-81 (71-69-73-94) | 61-54 (58-19-64-52) | 69-12 (68-12-70-07) | 60-03 (57-31-62-60) | 77-06 (76-30-77-70) | 65-44 (61-99-68-51) | 73-22 (72-43-73-99) | 63-91 (61-04-66-43) | 73-62 (69-06-78-25) | 62-58 (57-77-67-20) | 63-29 (56-34-71-29) | 55-64 (49-19-62-44) |
| Tunisia | 74-12 (73-09-75-18) | 62-92 (59-62-66-06) | 69-36 (68-15-70-56) | 60-38 (57-62-62-91) | 78-82 (77-62-80-64) | 67-00 (63-42-70-58) | 73-67 (71-98-75-57) | 64-07 (60-96-67-00) | 80-46 (78-87-82-70) | 68-32 (64-48-72-01) | 74-58 (72-07-77-24) | 65-02 (61-75-68-47) |

(Table 2 continues on next page)

| | 1990, at birth | | | | 2006, at birth | | | | 2016, at birth | | | |
|--------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Turkey | 73.48 (72.41-74.51) | 61.46 (58.05-64.58) | 66.84 (65.71-68.03) | 57.77 (55.03-60.35) | 80.56 (79.58-81.47) | 67.67 (63.94-71.19) | 73.90 (73.02-74.83) | 64.15 (61.44-66.84) | 82.33 (80.42-84.35) | 69.20 (65.37-72.84) | 75.84 (73.66-78.13) | 65.94 (62.76-69.10) |
| United Arab Emirates | 73.80 (71.06-76.69) | 62.23 (58.32-65.78) | 71.73 (68.48-74.76) | 61.87 (57.92-65.75) | 78.57 (77.31-79.40) | 66.33 (62.78-69.58) | 75.06 (73.81-76.25) | 64.89 (61.82-67.73) | 78.64 (76.18-80.63) | 66.53 (62.95-70.18) | 74.52 (72.10-76.83) | 64.62 (61.09-67.66) |
| Yemen | 58.84 (57.63-60.16) | 48.28 (45.15-51.30) | 58.57 (57.40-59.81) | 49.39 (46.58-51.99) | 65.68 (64.61-66.84) | 54.20 (50.83-57.27) | 65.37 (64.18-66.55) | 55.43 (52.41-58.18) | 67.94 (66.60-69.16) | 56.24 (52.77-59.33) | 65.51 (63.70-67.23) | 55.81 (52.76-58.59) |
| South Asia | 59.80 (59.24-60.35) | 50.89 (48.19-53.20) | 58.46 (57.94-58.97) | 50.96 (48.82-52.90) | 66.30 (65.82-66.79) | 56.41 (53.59-58.90) | 63.48 (62.93-64.03) | 55.24 (52.96-57.29) | 70.59 (70.08-71.12) | 60.00 (56.96-62.75) | 67.11 (66.46-67.67) | 58.43 (55.97-60.54) |
| Bangladesh | 58.61 (57.32-59.87) | 49.95 (47.23-52.37) | 57.61 (56.57-58.73) | 50.10 (47.67-52.26) | 70.37 (69.36-71.27) | 59.79 (56.76-62.56) | 66.64 (65.79-67.55) | 58.08 (55.57-60.33) | 75.10 (73.49-76.44) | 64.04 (60.88-67.11) | 70.47 (68.82-72.04) | 61.69 (58.94-64.30) |
| Bhutan | 59.12 (57.08-61.38) | 50.11 (47.16-53.12) | 60.68 (58.50-62.85) | 52.49 (49.55-55.02) | 71.21 (69.51-72.79) | 60.17 (56.70-63.28) | 69.12 (67.36-70.85) | 59.86 (57.04-62.76) | 75.91 (73.78-77.46) | 64.06 (60.43-67.44) | 72.23 (70.06-74.38) | 62.56 (59.32-65.53) |
| India | 59.69 (59.02-60.35) | 50.75 (48.01-53.17) | 58.29 (57.68-58.87) | 50.81 (48.69-52.79) | 66.01 (65.38-66.59) | 56.09 (53.27-58.68) | 63.22 (62.59-63.80) | 54.96 (52.71-57.10) | 70.33 (69.59-71.03) | 59.67 (56.62-62.44) | 66.93 (66.24-67.56) | 58.18 (55.72-60.37) |
| Nepal | 57.71 (56.74-58.64) | 49.63 (47.28-51.88) | 57.69 (56.67-58.72) | 50.14 (47.90-52.23) | 67.17 (66.20-68.21) | 57.74 (54.95-60.24) | 65.78 (64.76-66.74) | 57.35 (54.84-59.72) | 71.91 (71.07-72.81) | 61.80 (58.81-64.42) | 69.74 (68.49-71.04) | 60.92 (58.30-63.39) |
| Pakistan | 62.48 (61.49-63.45) | 53.50 (50.73-55.99) | 62.41 (61.38-63.42) | 54.57 (52.08-56.85) | 64.74 (63.33-66.23) | 55.57 (52.77-58.33) | 63.35 (61.99-64.79) | 55.58 (53.21-57.94) | 68.89 (66.86-71.19) | 59.09 (55.86-62.20) | 66.44 (64.27-68.45) | 58.23 (55.39-60.97) |
| Sub-Saharan Africa | 55.39 (54.92-55.80) | 47.87 (45.54-49.89) | 52.04 (51.48-52.68) | 45.35 (43.39-47.24) | 56.38 (55.83-56.89) | 48.88 (46.67-50.96) | 54.12 (53.55-54.65) | 47.33 (45.35-49.21) | 64.59 (63.91-65.26) | 56.05 (53.55-58.32) | 61.18 (60.30-62.02) | 53.57 (51.23-55.82) |
| Southern sub-Saharan Africa | 67.35 (66.60-68.16) | 58.18 (55.61-60.63) | 60.25 (59.31-61.19) | 52.67 (50.49-54.85) | 51.29 (50.07-52.53) | 48.31 (42.44-46.63) | 48.31 (47.41-49.22) | 42.47 (40.68-44.15) | 64.86 (63.79-65.96) | 55.75 (53.05-58.36) | 58.36 (57.38-59.33) | 50.90 (48.68-53.04) |
| Botswana | 67.60 (65.60-69.84) | 58.83 (55.88-61.79) | 61.23 (58.34-64.21) | 53.70 (50.51-56.85) | 51.04 (45.54-56.94) | 44.50 (39.85-49.30) | 47.99 (44.54-52.03) | 42.16 (38.85-45.79) | 69.21 (64.51-78.50) | 59.74 (54.95-66.73) | 61.69 (58.43-67.01) | 53.88 (50.27-58.28) |
| Lesotho | 64.20 (62.27-66.35) | 55.55 (52.72-58.36) | 56.59 (54.64-58.76) | 49.82 (47.38-52.48) | 45.45 (41.73-49.33) | 39.46 (36.03-42.72) | 41.01 (38.66-43.40) | 36.16 (33.65-38.50) | 53.67 (49.88-58.25) | 46.37 (42.72-50.26) | 47.13 (44.61-49.71) | 41.46 (38.97-43.98) |
| Namibia | 64.69 (63.56-65.91) | 56.44 (53.69-58.78) | 58.32 (57.03-59.62) | 51.36 (49.11-53.52) | 54.71 (50.75-58.90) | 47.83 (44.03-51.74) | 50.34 (47.92-52.74) | 44.40 (41.90-46.92) | 69.31 (65.48-75.19) | 60.14 (55.93-65.17) | 60.39 (57.83-63.37) | 53.00 (50.09-56.01) |
| South Africa | 68.05 (67.23-68.92) | 58.59 (55.83-61.09) | 60.72 (59.85-61.59) | 52.98 (50.78-55.06) | 52.28 (50.85-53.65) | 45.25 (42.99-47.42) | 49.72 (48.70-50.77) | 43.60 (41.64-45.51) | 65.51 (64.17-66.75) | 56.09 (53.35-58.88) | 59.24 (57.90-60.31) | 51.47 (49.14-53.72) |
| Swaziland | 65.42 (63.35-67.49) | 56.99 (54.16-59.75) | 59.06 (56.88-61.39) | 51.93 (49.38-54.50) | 44.30 (40.61-48.56) | 38.50 (35.00-42.26) | 41.59 (38.83-44.65) | 36.49 (33.84-39.30) | 62.16 (57.33-68.37) | 53.38 (48.49-58.49) | 53.25 (50.19-57.12) | 46.38 (43.19-49.67) |
| Zimbabwe | 64.36 (62.06-67.28) | 56.31 (53.45-59.60) | 59.02 (55.45-63.66) | 51.98 (48.42-56.28) | 47.07 (43.92-50.60) | 41.38 (38.39-44.54) | 44.34 (42.25-46.58) | 39.39 (37.15-41.71) | 61.87 (59.24-65.19) | 54.11 (51.15-57.35) | 56.66 (54.38-59.28) | 50.19 (47.52-53.02) |
| Western sub-Saharan Africa | 55.57 (54.69-56.34) | 47.67 (45.11-49.96) | 53.32 (52.28-54.31) | 46.21 (44.00-48.31) | 58.25 (57.26-59.21) | 50.20 (47.85-52.54) | 56.00 (55.03-56.96) | 48.79 (46.59-50.96) | 64.77 (63.72-65.90) | 55.98 (53.26-58.55) | 61.89 (60.59-63.06) | 54.12 (51.54-56.44) |
| Benin | 57.60 (56.50-58.70) | 48.94 (46.09-51.36) | 53.32 (52.04-54.64) | 45.99 (43.44-48.17) | 61.95 (60.46-63.47) | 53.19 (50.16-55.83) | 57.76 (56.51-59.08) | 50.39 (48.08-52.79) | 66.34 (64.76-68.10) | 57.19 (54.17-60.02) | 62.53 (60.91-64.06) | 54.72 (52.08-57.28) |

(Table 2 continues on next page)

| | 1990, at birth | | | | 2006, at birth | | | | 2016, at birth | | | |
|--------------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Burkina Faso | 52-09 (51-11-53-19) | 44-26 (41-75-46-61) | 49-12 (47-86-50-52) | 41-95 (39-58-44-21) | 56-81 (55-85-57-76) | 48-86 (46-30-51-19) | 54-41 (53-23-55-54) | 47-20 (44-77-49-52) | 62-06 (61-04-63-05) | 53-68 (51-08-56-24) | 59-51 (57-95-60-84) | 52-17 (49-79-54-55) |
| Cameroon | 59-63 (58-32-61-10) | 51-05 (48-24-53-59) | 57-28 (55-81-58-81) | 49-57 (47-05-51-91) | 56-56 (54-54-58-87) | 48-79 (46-01-51-46) | 54-00 (52-22-55-70) | 47-10 (44-57-49-48) | 62-02 (59-34-65-18) | 53-81 (50-69-57-12) | 58-34 (56-01-60-59) | 51-26 (48-34-54-11) |
| Cape Verde | 71-55 (70-49-72-79) | 61-66 (58-62-64-50) | 64-09 (62-76-65-53) | 56-21 (53-78-58-70) | 76-48 (74-84-78-29) | 65-95 (62-49-69-20) | 66-27 (64-07-69-43) | 58-27 (55-46-61-31) | 78-50 (77-22-80-25) | 67-83 (64-48-71-09) | 68-79 (66-61-71-11) | 60-61 (57-69-63-48) |
| Chad | 54-43 (53-27-55-67) | 46-62 (44-13-48-84) | 52-10 (50-71-53-32) | 45-17 (42-76-47-24) | 55-55 (53-89-57-37) | 47-63 (45-00-50-25) | 52-59 (50-96-54-22) | 45-68 (43-28-48-08) | 61-36 (59-50-63-20) | 52-71 (49-91-55-50) | 58-32 (56-39-60-28) | 50-74 (48-15-53-29) |
| Côte d'Ivoire | 57-68 (56-58-58-79) | 49-17 (46-50-51-82) | 52-39 (51-01-53-58) | 45-37 (42-99-47-51) | 56-04 (54-36-57-85) | 48-52 (45-85-50-93) | 52-09 (50-45-53-67) | 45-65 (43-25-47-87) | 62-29 (60-54-64-01) | 53-92 (51-07-56-57) | 57-72 (55-83-59-48) | 50-60 (48-00-53-01) |
| The Gambia | 62-30 (60-68-64-02) | 53-36 (50-28-56-14) | 60-88 (59-15-62-62) | 53-05 (50-22-55-72) | 66-26 (64-59-68-06) | 56-75 (53-72-59-73) | 62-80 (61-35-64-29) | 54-78 (51-92-57-28) | 69-22 (67-62-71-33) | 59-42 (56-30-62-46) | 65-46 (63-97-66-94) | 57-15 (54-20-59-62) |
| Ghana | 59-84 (58-36-61-44) | 51-82 (49-16-54-38) | 57-78 (56-18-59-37) | 50-59 (48-06-53-14) | 61-92 (60-12-63-95) | 54-01 (51-22-56-79) | 59-01 (57-79-60-31) | 52-03 (49-63-54-16) | 67-53 (65-97-69-68) | 58-98 (56-19-61-86) | 64-49 (62-75-66-08) | 56-90 (54-34-59-35) |
| Guinea | 51-91 (50-53-53-30) | 44-55 (42-05-46-92) | 52-11 (50-71-53-57) | 45-26 (42-89-47-43) | 57-25 (55-74-58-97) | 49-53 (47-03-51-96) | 55-58 (53-83-57-33) | 48-76 (46-21-51-09) | 61-62 (59-52-64-00) | 53-40 (50-48-56-36) | 59-76 (57-32-62-13) | 52-48 (49-61-55-35) |
| Guinea-Bissau | 51-89 (50-31-53-78) | 44-89 (42-43-47-26) | 46-57 (44-71-48-57) | 40-83 (38-59-43-17) | 56-06 (54-03-58-01) | 48-66 (45-95-51-27) | 51-56 (50-00-53-30) | 45-36 (43-07-47-54) | 61-37 (59-44-63-33) | 53-32 (50-43-56-03) | 56-30 (54-53-58-20) | 49-60 (47-09-52-04) |
| Liberia | 51-76 (50-48-53-07) | 43-77 (41-14-46-17) | 47-99 (46-63-49-49) | 40-93 (38-51-43-17) | 58-82 (57-58-60-31) | 49-98 (47-20-52-57) | 58-52 (57-29-59-75) | 49-98 (47-27-52-50) | 64-82 (63-59-66-07) | 55-40 (52-50-58-00) | 64-00 (62-76-65-16) | 55-19 (52-43-57-82) |
| Mali | 49-39 (48-14-50-66) | 42-49 (40-11-44-73) | 49-41 (48-04-50-70) | 42-56 (40-19-44-83) | 57-72 (56-20-59-37) | 49-75 (47-12-52-25) | 56-46 (54-86-58-11) | 48-81 (46-21-51-40) | 62-74 (60-01-65-49) | 54-29 (51-16-57-53) | 61-02 (58-09-63-77) | 53-18 (49-98-56-44) |
| Mauritania | 60-41 (59-05-61-76) | 52-16 (49-49-54-62) | 59-50 (58-11-61-07) | 51-87 (49-30-54-28) | 65-99 (64-25-68-11) | 57-15 (54-23-59-95) | 66-83 (64-96-68-67) | 58-11 (55-20-60-91) | 70-21 (67-56-73-27) | 60-85 (57-43-64-22) | 70-29 (67-75-73-20) | 61-28 (57-96-64-65) |
| Niger | 48-07 (46-57-49-48) | 41-61 (39-47-43-84) | 46-19 (44-61-47-75) | 40-45 (38-31-42-43) | 57-00 (55-24-58-62) | 49-47 (46-71-51-86) | 55-43 (53-62-57-32) | 48-68 (46-27-51-07) | 62-83 (60-16-65-50) | 54-70 (51-47-57-94) | 60-60 (58-25-62-87) | 53-42 (50-58-56-30) |
| Nigeria | 55-80 (53-93-57-43) | 47-83 (45-05-50-36) | 53-99 (51-98-55-95) | 46-72 (44-09-49-35) | 58-29 (56-34-60-30) | 50-06 (47-39-52-97) | 56-69 (54-87-58-62) | 49-22 (46-60-51-93) | 66-41 (64-30-69-14) | 57-17 (53-86-60-50) | 63-69 (61-16-66-21) | 55-46 (52-16-58-36) |
| São Tomé and Príncipe | 64-55 (62-95-66-14) | 56-13 (53-44-58-78) | 61-94 (60-46-63-55) | 54-06 (51-51-56-55) | 68-04 (66-93-69-14) | 59-38 (56-77-61-81) | 65-63 (64-35-66-88) | 57-37 (54-72-59-76) | 72-09 (70-52-73-93) | 62-86 (59-84-65-70) | 69-04 (66-71-71-46) | 60-40 (57-08-63-43) |
| Senegal | 59-13 (57-93-60-29) | 51-16 (48-64-53-57) | 55-97 (54-99-56-95) | 49-06 (46-73-51-08) | 63-73 (62-67-64-93) | 55-34 (52-58-57-81) | 60-56 (59-49-61-69) | 53-17 (50-79-55-42) | 67-78 (66-59-68-83) | 58-95 (56-18-61-49) | 64-64 (63-32-65-97) | 56-81 (54-17-59-31) |
| Sierra Leone | 52-78 (51-41-54-15) | 45-44 (42-94-47-75) | 48-43 (46-82-50-02) | 42-07 (39-83-44-17) | 54-14 (52-72-55-61) | 46-80 (44-24-49-05) | 51-62 (50-37-52-85) | 44-94 (42-80-47-02) | 60-19 (58-38-62-14) | 52-30 (49-54-55-01) | 58-11 (56-42-59-82) | 50-98 (48-48-53-42) |
| Togo | 58-64 (57-47-59-92) | 50-53 (47-88-52-99) | 56-13 (54-90-57-39) | 49-04 (46-54-51-27) | 59-12 (57-12-61-53) | 51-10 (48-41-53-75) | 54-86 (53-29-56-46) | 48-25 (45-93-50-52) | 65-08 (63-58-66-82) | 56-63 (53-68-59-26) | 60-11 (58-22-61-70) | 53-13 (50-60-55-49) |
| Eastern sub-Saharan Africa | 52-92 (52-39-53-47) | 46-24 (44-24-48-02) | 49-46 (48-81-50-17) | 43-47 (41-58-45-13) | 56-64 (55-91-57-36) | 49-56 (47-39-51-57) | 54-57 (53-86-55-19) | 48-01 (46-07-49-81) | 65-08 (64-13-66-09) | 56-97 (54-43-59-30) | 61-72 (60-63-62-80) | 54-35 (52-04-56-58) |

(Table 2 continues on next page)

| | 1990, at birth | | | | 2006, at birth | | | | 2016, at birth | | | |
|--------------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Burundi | 49.28 (47.21-51.62) | 43.59 (41.19-46.03) | 46.69 (44.34-49.16) | 41.49 (39.09-43.99) | 55.18 (53.18-57.01) | 48.71 (46.26-51.06) | 53.46 (51.54-55.47) | 47.33 (44.70-49.70) | 61.51 (58.70-64.10) | 54.51 (51.44-57.29) | 59.20 (56.03-62.28) | 52.57 (49.11-55.76) |
| Comoros | 59.46 (57.78-61.52) | 51.92 (49.20-54.61) | 57.80 (55.88-59.73) | 50.47 (47.88-52.97) | 65.30 (63.79-67.10) | 57.14 (54.36-59.97) | 63.78 (62.30-65.32) | 55.98 (53.42-58.63) | 68.52 (66.78-70.86) | 60.05 (57.27-63.16) | 66.61 (64.38-68.77) | 58.57 (55.52-61.31) |
| Djibouti | 64.10 (62.77-65.15) | 55.99 (53.27-58.39) | 58.43 (57.21-59.67) | 51.61 (49.41-53.77) | 64.36 (61.78-67.15) | 56.34 (53.15-59.62) | 59.51 (57.39-61.57) | 52.59 (49.86-55.17) | 68.80 (66.11-71.85) | 60.24 (56.93-63.65) | 64.67 (62.50-66.70) | 57.03 (54.20-59.80) |
| Eritrea | 53.74 (52.30-55.22) | 47.10 (44.78-49.34) | 50.96 (49.44-52.49) | 45.01 (42.76-47.10) | 61.51 (59.71-63.42) | 54.15 (51.29-56.75) | 60.05 (58.00-62.07) | 52.98 (50.39-55.63) | 64.53 (62.75-66.62) | 56.84 (54.01-59.55) | 62.88 (60.62-65.07) | 55.54 (52.73-58.26) |
| Ethiopia | 48.89 (47.64-50.20) | 43.00 (40.98-44.98) | 44.37 (43.16-45.76) | 39.31 (37.54-41.04) | 57.30 (55.68-59.03) | 50.56 (48.24-52.87) | 56.81 (55.05-58.31) | 50.17 (47.86-52.43) | 66.53 (64.21-68.96) | 58.67 (55.72-61.51) | 64.74 (62.10-67.59) | 57.20 (54.14-60.45) |
| Kenya | 62.63 (61.99-63.30) | 55.07 (52.71-57.15) | 60.25 (59.41-61.07) | 53.14 (50.93-55.15) | 59.91 (59.18-60.60) | 52.60 (50.41-54.67) | 57.04 (56.37-57.71) | 50.40 (48.34-52.22) | 69.03 (68.24-69.88) | 60.30 (57.58-62.64) | 64.72 (63.89-65.56) | 56.94 (54.49-59.05) |
| Madagascar | 56.76 (55.42-58.23) | 49.55 (47.22-51.91) | 54.43 (53.13-55.73) | 47.74 (45.56-49.87) | 61.40 (59.57-63.44) | 53.79 (50.81-56.51) | 59.41 (57.47-61.41) | 52.19 (49.62-54.76) | 63.88 (60.78-67.84) | 56.22 (52.80-59.92) | 61.51 (58.34-64.83) | 54.29 (51.18-57.82) |
| Malawi | 49.89 (48.02-51.91) | 43.53 (41.08-46.03) | 47.56 (44.19-51.16) | 41.45 (38.16-44.72) | 49.75 (46.64-53.37) | 43.49 (40.30-46.97) | 47.18 (44.96-49.62) | 41.39 (39.05-44.13) | 62.59 (59.81-65.90) | 54.63 (51.37-57.95) | 57.90 (55.27-60.52) | 50.94 (48.22-53.98) |
| Mozambique | 52.80 (51.72-53.94) | 45.66 (43.25-47.89) | 48.51 (47.27-49.70) | 42.25 (40.17-44.09) | 54.89 (51.81-58.00) | 47.39 (44.02-50.59) | 50.33 (48.12-52.56) | 43.91 (41.18-46.35) | 62.88 (60.39-65.36) | 54.32 (51.22-57.53) | 57.05 (54.88-59.18) | 49.78 (46.92-52.53) |
| Rwanda | 50.78 (49.17-52.31) | 44.72 (42.38-46.96) | 47.51 (45.55-49.50) | 42.13 (39.82-44.32) | 61.04 (59.22-62.93) | 53.38 (50.75-55.88) | 57.93 (56.51-59.53) | 50.93 (48.55-53.13) | 69.32 (67.21-71.54) | 60.74 (57.78-63.78) | 65.98 (63.86-67.95) | 58.02 (55.04-60.77) |
| Somalia | 51.72 (50.09-53.38) | 45.34 (42.86-47.64) | 49.60 (47.49-51.75) | 43.81 (41.26-46.38) | 54.55 (52.80-56.31) | 47.93 (45.51-50.47) | 53.54 (51.26-55.80) | 47.33 (44.72-50.09) | 57.74 (55.75-59.45) | 50.75 (48.08-53.18) | 56.64 (54.11-59.07) | 50.10 (47.03-52.87) |
| South Sudan | 53.28 (51.20-55.51) | 44.83 (41.62-47.86) | 49.85 (47.40-52.59) | 42.34 (39.34-45.54) | 57.77 (54.98-60.99) | 49.35 (45.93-52.82) | 55.76 (53.13-58.60) | 47.94 (44.83-51.03) | 60.72 (57.90-63.93) | 52.40 (49.04-55.82) | 58.71 (56.05-61.53) | 50.96 (47.77-54.05) |
| Tanzania | 55.98 (54.76-57.26) | 48.69 (46.28-51.04) | 53.74 (52.07-55.35) | 47.08 (44.67-49.47) | 56.72 (54.77-59.54) | 49.54 (46.90-52.22) | 55.30 (53.82-56.98) | 48.68 (46.36-51.06) | 66.05 (64.28-68.33) | 57.79 (54.94-60.76) | 62.59 (60.50-64.45) | 55.33 (52.61-57.81) |
| Uganda | 52.07 (50.38-53.44) | 44.88 (42.39-47.10) | 46.34 (43.41-48.89) | 40.17 (37.23-43.06) | 55.59 (53.98-57.23) | 48.16 (45.69-50.66) | 51.84 (50.37-53.29) | 45.29 (42.89-47.31) | 64.75 (62.93-67.01) | 56.33 (53.55-59.24) | 59.77 (57.99-61.40) | 52.45 (50.10-54.90) |
| Zambia | 52.33 (50.38-54.59) | 45.94 (43.52-48.33) | 52.21 (49.34-54.78) | 45.87 (42.88-48.71) | 48.21 (45.39-51.35) | 42.34 (39.62-45.29) | 45.23 (43.18-47.31) | 40.12 (37.88-42.23) | 61.95 (58.22-66.84) | 54.14 (50.45-58.71) | 55.58 (52.54-59.34) | 49.10 (46.12-52.44) |
| Central sub-Saharan Africa | 54.41 (53.25-55.53) | 46.46 (43.99-48.74) | 50.83 (49.71-51.99) | 43.71 (41.49-45.86) | 56.35 (55.20-57.49) | 48.42 (45.79-50.67) | 54.80 (53.60-55.95) | 47.29 (44.88-49.53) | 62.80 (61.19-64.43) | 54.20 (51.34-56.76) | 60.62 (58.87-62.28) | 52.58 (49.77-55.26) |
| Angola | 52.16 (49.57-55.16) | 45.20 (42.37-48.22) | 48.27 (45.63-51.05) | 42.20 (39.52-45.23) | 57.13 (54.31-60.11) | 49.77 (46.75-52.71) | 56.77 (53.90-59.62) | 49.56 (46.60-52.46) | 65.37 (61.78-68.70) | 56.85 (53.28-60.26) | 63.94 (60.34-67.08) | 55.76 (52.30-59.13) |
| Central African Republic | 51.04 (48.84-53.26) | 43.80 (40.92-46.39) | 45.38 (43.16-47.67) | 39.24 (36.70-41.72) | 47.44 (44.14-51.09) | 41.04 (37.74-44.37) | 43.21 (40.31-46.22) | 37.65 (34.73-40.63) | 52.59 (48.76-56.64) | 45.60 (41.97-49.45) | 47.91 (44.46-51.58) | 41.91 (38.62-45.46) |
| Congo (Brazzaville) | 55.96 (53.78-58.10) | 48.26 (45.37-51.06) | 52.19 (50.02-54.61) | 45.47 (42.70-48.24) | 56.61 (54.64-58.58) | 48.95 (46.20-51.56) | 58.33 (56.48-60.14) | 50.60 (47.97-53.23) | 62.35 (59.54-65.61) | 53.99 (50.40-57.77) | 63.62 (60.10-67.33) | 55.47 (51.93-59.12) |

(Table 2 continues on next page)

| | 1990, at birth | | | | 2006, at birth | | | | 2016, at birth | | | |
|----------------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Democratic Republic of the Congo | 55.32 (53.74–56.81) | 46.96 (44.23–49.57) | 52.33 (50.71–53.96) | 44.65 (42.09–47.12) | 56.75 (55.28–58.22) | 48.53 (45.70–51.08) | 55.03 (53.48–56.45) | 47.24 (44.61–49.74) | 62.72 (60.80–64.53) | 53.99 (50.96–56.80) | 60.44 (58.33–62.42) | 52.26 (49.34–55.04) |
| Equatorial Guinea | 49.82 (47.16–52.85) | 43.11 (40.10–46.00) | 46.54 (43.66–49.72) | 40.62 (37.58–43.52) | 59.26 (54.70–64.65) | 51.26 (47.01–55.97) | 58.66 (54.71–63.17) | 50.91 (46.94–55.13) | 66.55 (62.28–71.43) | 57.44 (53.04–61.96) | 64.49 (60.25–69.11) | 55.97 (51.95–60.33) |
| Gabon | 62.90 (61.23–64.49) | 54.14 (51.30–56.75) | 56.02 (54.28–57.83) | 49.01 (46.57–51.43) | 61.32 (59.50–63.25) | 52.87 (50.05–55.57) | 59.71 (57.34–61.77) | 52.16 (49.40–54.83) | 68.31 (65.83–71.18) | 58.73 (55.38–62.22) | 64.90 (61.35–68.37) | 56.59 (52.95–60.04) |

Data in parentheses are 95% uncertainty intervals. To download the data in this table, please visit the Global Health Data Exchange. GBD=Global Burden of Disease. HALE=healthy life expectancy. SDI=Socio-demographic index.

Table 2: Global, regional, and GBD location-specific life expectancy and HALE at birth, by sex, in 1990, 2006, and 2016

3.91 million), extensively drug-resistant tuberculosis 369 000 (301 000–445 000), drug-susceptible HIV/AIDS-tuberculosis 11.7 million (8.15 million to 15.5 million), multidrug-resistant HIV/AIDS-tuberculosis without extensive drug resistance 0.98 million (0.60 million to 1.48 million), and extensively drug-resistant HIV/AIDS-tuberculosis 57 300 (34 500–89 400; table 1).

Changes in leading causes of disease burden over time and SDI quintiles

The leading Level 3 causes of global DALYs in 1990 were lower respiratory infections, diarrhoeal diseases, and ischaemic heart disease (figure 2A). By 2016, lower respiratory infections had decreased in relative rank to the third-leading cause, whereas diarrhoeal diseases were fifth and ischaemic heart disease rose to first. The leading 30 causes of DALYs by SDI quintile varied considerably (figure 2B–F). CMNN causes were dominant in the low-SDI quintile, occupying 11 of the top 15 causes in 2016. NCDs were more commonly among the leading causes of DALYs with successively higher levels of SDI, occupying 12 of the top 15 causes in the high-SDI quintile in 2016.

For low SDI, only two NCD causes (congenital birth defects and ischaemic heart disease) ranked in the top ten in 2016, compared with all of the top ten for high SDI. Of the leading 30 causes from 2006 to 2016, HIV/AIDS had the largest decrease in age-standardised DALY rates for low, low-middle, and middle SDI. For high-middle SDI, the largest decrease was for lower respiratory infection, and for high SDI, the largest decrease was for road injuries. At low-middle SDI, 14 of the leading 30 causes in 2016 were NCDs, whereas 13 were CMNN causes and three were injuries, compared with middle SDI, where seven were CMNN causes, 19 were NCDs, and four were injuries. The leading seven causes for low, low-middle, and middle SDI quintiles all decreased in age-standardised DALY rates from 2006 to 2016.

At high-middle and high SDI, the leading three causes of DALYs remained the same from 2006 to 2016 (ischaemic heart disease, low back and neck pain, and cerebrovascular disease). Two of the leading 30 causes for high-middle SDI were CMNN causes in 2016, whereas 24 were NCDs and four were injuries, compared with high SDI, where one was a CMNN cause, 26 were NCDs, and three were injuries. The leading five causes for high-middle SDI decreased in age-standardised DALY rates from both 1990 to 2006 and 2006 to 2016, although this pattern did not hold for high SDI, where just the leading cause decreased over both intervals. Road injuries and falls are the only injuries that appear across all five SDI quintiles; self-harm appears in all quintiles except for low SDI. Ischaemic heart disease and cerebrovascular disease appear in the leading 13 causes across all SDI quintiles. HIV/AIDS appears in the leading ten causes for all quintiles except for high and high-middle SDI, where it did not appear in the leading 30.

Regional and country-specific HALE and DALYs in 2016

In 2016, HALE at birth was highest in Singapore for both females (75.2 years [95% UI 71.9–78.6]) and males (72.0 years [68.8–75.1]) and lowest for females in the Central African Republic (45.6 years [42.0–49.5]) and for males in Lesotho (41.5 [39.0–44.0]; table 2 and 3). In 2016, HALE at birth was greater than 70 years in only 12 locations for males and was greater than this threshold in 49 locations for females. HALE at birth was less than 50 years for both sexes in three locations: the Central African Republic, Lesotho, and Afghanistan. Between 1990 and 2016, HALE at birth increased for males in 160 locations and for females in 167 locations. For males, the largest increase occurred in Ethiopia, rising from 39.3 years (37.5–41.0) in 1990 to 57.2 years (54.1–60.45) in 2016. For females, the largest increase occurred in the Maldives, rising from 53.6 (51.1–56.1) in 1990 to 70.3 (66.3–73.7) in 2016. Over the same time period, HALE at birth decreased in just

| | 1990, at age 65 years | | | | 2006, at age 65 years | | | | 2016, at age 65 years | | | |
|---------------------------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| Global | 15-90 (15-82-15-98) | 11-92 (10-88-12-89) | 13-33 (13-26-13-39) | 10-09 (9-22-10-87) | 17-27 (17-19-17-35) | 12-93 (11-74-13-98) | 14-69 (14-62-14-75) | 11-10 (10-13-11-97) | 18-57 (18-37-18-72) | 13-88 (12-57-15-02) | 15-72 (15-61-15-83) | 11-87 (10-83-12-80) |
| High SDI | 18-57 (18-50-18-63) | 14-06 (12-85-15-16) | 14-79 (14-73-14-84) | 11-22 (10-27-12-08) | 20-72 (20-67-20-78) | 15-69 (14-31-16-93) | 17-22 (17-17-17-28) | 13-00 (11-86-14-03) | 21-67 (21-53-21-81) | 16-38 (14-91-17-68) | 18-37 (18-22-18-50) | 13-86 (12-65-14-98) |
| High-middle SDI | 16-31 (16-07-16-54) | 12-15 (11-02-13-22) | 13-35 (13-16-13-54) | 10-09 (9-20-10-92) | 17-39 (17-15-17-62) | 13-00 (11-78-14-08) | 14-27 (14-06-14-45) | 10-81 (9-86-11-68) | 19-19 (18-36-19-89) | 14-32 (12-85-15-71) | 15-84 (15-39-16-28) | 12-01 (10-90-13-06) |
| Middle SDI | 14-81 (14-63-14-98) | 11-25 (10-32-12-13) | 12-64 (12-50-12-78) | 9-78 (9-00-10-50) | 16-55 (16-43-16-67) | 12-48 (11-38-13-50) | 14-09 (13-98-14-21) | 10-84 (9-96-11-66) | 18-19 (18-02-18-35) | 13-68 (12-48-14-77) | 15-01 (14-86-15-17) | 11-50 (10-55-12-33) |
| Low-middle SDI | 13-08 (12-90-13-28) | 9-47 (8-51-10-37) | 12-48 (12-32-12-64) | 9-15 (8-28-9-96) | 14-35 (14-16-14-54) | 10-40 (9-36-11-34) | 13-24 (13-10-13-39) | 9-74 (8-83-10-58) | 15-50 (15-26-15-73) | 11-24 (10-13-12-27) | 13-95 (13-75-14-14) | 10-27 (9-28-11-13) |
| Low SDI | 11-65 (11-45-11-86) | 8-54 (7-67-9-29) | 11-75 (11-52-11-97) | 8-59 (7-72-9-42) | 12-29 (12-02-12-58) | 9-08 (8-19-9-91) | 12-43 (12-20-12-67) | 9-17 (8-24-9-97) | 13-32 (13-00-13-64) | 9-90 (8-94-10-78) | 13-22 (12-89-13-54) | 9-80 (8-82-10-69) |
| High income | 18-77 (18-71-18-84) | 14-28 (13-06-15-39) | 14-96 (14-90-15-01) | 11-40 (10-45-12-27) | 20-89 (20-83-20-95) | 15-88 (14-51-17-12) | 17-35 (17-30-17-40) | 13-16 (12-02-14-18) | 21-80 (21-66-21-94) | 16-55 (15-09-17-85) | 18-47 (18-32-18-60) | 14-00 (12-79-15-11) |
| High-income North America | 19-09 (19-03-19-16) | 14-49 (13-27-15-62) | 15-17 (15-11-15-24) | 11-46 (10-48-12-36) | 20-05 (19-99-20-12) | 15-06 (13-71-16-29) | 17-33 (17-28-17-40) | 12-91 (11-73-13-98) | 20-74 (20-59-20-89) | 15-46 (14-05-16-73) | 18-19 (18-05-18-34) | 13-48 (12-23-14-63) |
| Canada | 19-69 (19-47-19-89) | 15-30 (14-07-16-43) | 15-50 (15-31-15-70) | 12-03 (11-03-12-93) | 21-08 (20-86-21-27) | 16-28 (15-00-17-49) | 17-92 (17-75-18-13) | 13-82 (12-68-14-86) | 21-96 (21-66-22-27) | 16-91 (15-50-18-21) | 19-12 (18-82-19-44) | 14-71 (13-46-15-88) |
| Greenland | 11-30 (10-69-11-92) | 8-66 (7-82-9-51) | 8-95 (8-51-9-40) | 6-83 (6-17-7-45) | 11-88 (11-27-12-47) | 9-10 (8-25-9-92) | 10-70 (10-19-11-17) | 8-20 (7-47-8-92) | 14-20 (12-77-15-95) | 10-89 (9-51-12-36) | 11-89 (10-54-13-29) | 9-14 (7-93-10-46) |
| USA | 19-04 (18-97-19-11) | 14-41 (13-19-15-55) | 15-14 (15-07-15-20) | 11-41 (10-43-12-30) | 19-94 (19-88-20-01) | 14-93 (13-57-16-16) | 17-27 (17-21-17-33) | 12-80 (11-62-13-88) | 20-60 (20-44-20-76) | 15-30 (13-89-16-59) | 18-09 (17-93-18-25) | 13-34 (12-08-14-49) |
| Australasia | 18-95 (18-79-19-12) | 14-66 (13-46-15-73) | 15-10 (14-95-15-25) | 11-65 (10-69-12-53) | 21-35 (21-21-21-49) | 16-52 (15-14-17-76) | 18-31 (18-17-18-46) | 14-11 (12-94-15-15) | 22-10 (21-65-22-56) | 17-07 (15-62-18-38) | 19-37 (18-91-19-84) | 14-92 (13-66-16-08) |
| Australia | 19-05 (18-88-19-25) | 14-73 (13-54-15-81) | 15-18 (15-01-15-35) | 11-71 (10-75-12-59) | 21-51 (21-34-21-68) | 16-63 (15-26-17-91) | 18-40 (18-24-18-58) | 14-17 (12-98-15-23) | 22-22 (21-67-22-74) | 17-16 (15-72-18-49) | 19-47 (18-93-19-96) | 14-98 (13-69-16-17) |
| New Zealand | 18-43 (18-18-18-66) | 14-29 (13-11-15-34) | 14-71 (14-50-14-92) | 11-36 (10-40-12-22) | 20-52 (20-29-20-73) | 15-95 (14-61-17-11) | 17-83 (17-62-18-03) | 13-81 (12-64-14-85) | 21-48 (20-65-22-40) | 16-64 (15-19-18-03) | 18-90 (18-08-19-77) | 14-60 (13-30-15-87) |
| High-income Asia Pacific | 19-46 (19-31-19-61) | 15-03 (13-80-16-15) | 15-65 (15-51-15-78) | 11-91 (10-90-12-82) | 22-73 (22-57-22-88) | 17-47 (15-99-18-77) | 17-93 (17-80-18-07) | 13-50 (12-30-14-59) | 23-79 (23-27-24-24) | 18-30 (16-64-19-74) | 19-20 (18-75-19-61) | 14-51 (13-22-15-73) |
| Brunei | 16-84 (15-98-17-54) | 12-75 (11-48-13-92) | 14-52 (14-31-14-78) | 10-70 (9-68-11-64) | 18-67 (18-25-19-03) | 14-26 (13-01-15-41) | 15-85 (15-56-16-19) | 11-78 (10-69-12-79) | 18-88 (18-18-20-19) | 14-50 (13-16-15-79) | 15-88 (15-05-17-40) | 11-88 (10-61-13-13) |
| Japan | 19-96 (19-92-20-00) | 15-43 (14-15-16-55) | 16-22 (16-19-16-25) | 12-36 (11-32-13-29) | 23-23 (23-17-23-28) | 17-89 (16-38-19-22) | 18-24 (18-21-18-28) | 13-75 (12-53-14-83) | 24-24 (24-07-24-40) | 18-67 (17-07-20-03) | 19-53 (19-35-19-70) | 14-78 (13-50-15-95) |
| Singapore | 17-64 (16-49-18-82) | 13-74 (12-40-15-01) | 14-65 (13-72-15-61) | 11-26 (10-13-12-42) | 21-50 (20-48-22-49) | 16-78 (15-30-18-19) | 17-71 (16-77-18-66) | 13-67 (12-45-14-94) | 23-33 (21-62-25-20) | 18-16 (16-31-20-02) | 19-70 (17-89-21-49) | 15-13 (13-37-16-82) |
| South Korea | 16-70 (15-82-17-56) | 12-82 (11-56-14-06) | 12-36 (11-66-13-12) | 9-33 (8-35-10-31) | 20-05 (19-28-20-86) | 15-26 (13-84-16-68) | 16-20 (15-45-17-11) | 12-11 (10-94-13-41) | 21-67 (19-44-23-94) | 16-62 (14-54-18-80) | 17-52 (15-45-20-08) | 13-20 (11-32-15-13) |

(Table 3 continues on next page)

| | 1990, at age 65 years | | | | 2006, at age 65 years | | | | 2016, at age 65 years | | | |
|--------------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Western Europe | 18.48 (18.37-18.59) | 13.98 (12.75-15.11) | 14.70 (14.61-14.79) | 11.25 (10.32-12.10) | 20.81 (20.71-20.91) | 15.80 (14.41-17.04) | 17.26 (17.17-17.36) | 13.23 (12.12-14.24) | 21.76 (21.56-21.96) | 16.54 (15.08-17.87) | 18.49 (18.29-18.69) | 14.21 (13.00-15.28) |
| Andorra | 20.84 (19.08-22.79) | 15.59 (13.63-17.52) | 16.54 (15.62-17.99) | 12.54 (11.23-13.98) | 23.19 (21.49-24.44) | 17.38 (15.50-19.13) | 18.33 (17.43-19.29) | 13.94 (12.51-15.26) | 23.06 (20.90-24.50) | 17.31 (15.33-19.16) | 18.46 (17.43-20.09) | 14.08 (12.52-15.58) |
| Austria | 17.88 (17.71-18.06) | 13.60 (12.43-14.67) | 14.36 (14.20-14.51) | 10.97 (9.99-11.81) | 20.48 (20.31-20.63) | 15.61 (14.25-16.81) | 17.16 (17.00-17.32) | 13.07 (11.92-14.08) | 21.45 (20.98-21.95) | 16.35 (14.86-17.69) | 18.41 (18.01-18.90) | 14.08 (12.82-15.30) |
| Belgium | 18.47 (18.16-18.74) | 13.84 (12.59-14.99) | 14.29 (14.00-14.59) | 10.86 (9.84-11.73) | 20.49 (20.24-20.77) | 15.22 (13.79-16.51) | 16.87 (16.60-17.15) | 12.70 (11.54-13.74) | 21.38 (20.63-22.22) | 16.00 (14.41-17.50) | 17.99 (17.22-18.72) | 13.62 (12.27-14.90) |
| Cyprus | 15.67 (15.34-15.99) | 11.88 (10.80-12.88) | 14.03 (13.70-14.36) | 10.77 (9.81-11.61) | 18.99 (18.64-19.34) | 14.41 (13.09-15.59) | 16.52 (16.19-16.84) | 12.64 (11.53-13.62) | 20.17 (19.74-20.58) | 15.38 (14.04-16.67) | 17.44 (17.01-17.92) | 13.39 (12.22-14.49) |
| Denmark | 17.97 (17.41-18.50) | 13.58 (12.33-14.73) | 14.27 (13.81-14.80) | 10.99 (10.00-11.87) | 19.09 (18.53-19.65) | 14.47 (13.09-15.70) | 16.33 (15.86-16.81) | 12.50 (11.38-13.56) | 20.76 (19.77-21.83) | 15.65 (14.08-17.10) | 18.01 (17.04-18.92) | 13.77 (12.45-15.05) |
| Finland | 17.74 (17.50-17.98) | 13.32 (12.13-14.43) | 13.76 (13.56-13.97) | 10.37 (9.43-11.23) | 20.74 (20.51-20.94) | 15.37 (13.87-16.74) | 16.73 (16.52-16.95) | 12.42 (11.26-13.46) | 21.99 (21.44-22.62) | 16.49 (14.87-18.00) | 18.39 (17.82-18.97) | 13.86 (12.61-15.07) |
| France | 20.15 (19.94-20.34) | 15.31 (13.98-16.52) | 15.72 (15.52-15.93) | 12.08 (11.05-13.01) | 22.25 (22.04-22.44) | 17.06 (15.60-18.34) | 17.91 (17.71-18.15) | 13.85 (12.66-14.89) | 23.22 (22.83-23.66) | 17.74 (16.17-19.19) | 19.24 (18.87-19.63) | 14.87 (13.56-16.01) |
| Germany | 17.68 (17.28-18.06) | 13.32 (12.06-14.47) | 14.03 (13.65-14.41) | 10.67 (9.69-11.56) | 20.21 (19.87-20.57) | 15.31 (13.94-16.58) | 16.75 (16.41-17.13) | 12.79 (11.69-13.86) | 21.10 (20.46-21.82) | 15.98 (14.47-17.42) | 17.93 (17.27-18.62) | 13.69 (12.43-14.90) |
| Greece | 18.02 (17.79-18.26) | 13.71 (12.55-14.78) | 15.58 (15.35-15.83) | 11.98 (10.98-12.88) | 20.24 (20.03-20.46) | 15.40 (14.10-16.59) | 17.42 (17.19-17.64) | 13.37 (12.24-14.38) | 20.98 (20.42-21.55) | 16.07 (14.70-17.39) | 18.34 (17.72-18.99) | 14.12 (12.91-15.28) |
| Iceland | 19.13 (18.70-19.55) | 14.48 (13.19-15.67) | 16.16 (15.82-16.53) | 12.37 (11.30-13.32) | 20.76 (20.29-21.27) | 15.73 (14.34-16.95) | 18.31 (17.99-18.66) | 14.02 (12.79-15.10) | 21.35 (20.53-22.14) | 16.24 (14.78-17.62) | 18.98 (18.37-19.56) | 14.60 (13.28-15.89) |
| Ireland | 16.89 (16.46-17.33) | 12.86 (11.69-13.94) | 13.36 (13.00-13.78) | 10.28 (9.37-11.12) | 19.70 (19.27-20.16) | 14.96 (13.66-16.21) | 16.66 (16.26-17.04) | 12.79 (11.69-13.85) | 21.03 (19.99-22.10) | 16.02 (14.42-17.48) | 18.13 (17.14-19.09) | 13.96 (12.61-15.28) |
| Israel | 16.81 (15.99-17.61) | 12.74 (11.45-13.98) | 15.08 (14.39-15.87) | 11.53 (10.39-12.59) | 20.00 (19.21-20.82) | 15.12 (13.63-16.54) | 17.63 (16.86-18.41) | 13.46 (12.15-14.68) | 21.54 (20.02-22.96) | 16.31 (14.53-17.97) | 18.93 (17.36-20.55) | 14.48 (12.89-16.15) |
| Italy | 18.91 (18.73-19.10) | 14.31 (13.04-15.46) | 15.08 (14.91-15.25) | 11.53 (10.55-12.45) | 21.36 (21.18-21.52) | 16.26 (14.82-17.57) | 17.57 (17.39-17.76) | 13.50 (12.38-14.54) | 21.94 (21.44-22.46) | 16.77 (15.29-18.16) | 18.60 (18.05-19.13) | 14.34 (13.08-15.50) |
| Luxembourg | 17.75 (17.43-18.09) | 13.25 (12.00-14.38) | 14.01 (13.75-14.27) | 10.51 (9.54-11.40) | 20.82 (20.44-21.15) | 15.60 (14.14-16.86) | 17.44 (17.14-17.75) | 13.13 (11.92-14.24) | 21.49 (20.63-22.52) | 16.16 (14.62-17.61) | 18.95 (18.21-19.74) | 14.32 (12.93-15.58) |
| Malta | 16.55 (15.73-17.38) | 12.60 (11.35-13.85) | 14.28 (13.61-14.99) | 10.99 (9.95-12.06) | 19.57 (18.79-20.34) | 14.93 (13.49-16.22) | 16.55 (15.89-17.23) | 12.76 (11.64-13.80) | 21.34 (19.94-22.91) | 16.28 (14.58-18.01) | 18.03 (16.76-19.40) | 13.92 (12.45-15.45) |
| Netherlands | 18.96 (18.64-19.33) | 14.26 (12.98-15.45) | 14.43 (14.11-14.73) | 10.98 (10.03-11.85) | 20.17 (19.85-20.47) | 15.21 (13.82-16.48) | 16.73 (16.45-17.05) | 12.72 (11.62-13.74) | 21.38 (20.65-22.06) | 16.12 (14.61-17.54) | 18.11 (17.36-18.78) | 13.79 (12.53-15.00) |
| Norway | 18.81 (18.47-19.16) | 14.30 (13.04-15.45) | 14.80 (14.50-15.15) | 11.24 (10.21-12.17) | 20.62 (20.28-20.92) | 15.74 (14.38-16.96) | 17.42 (17.12-17.70) | 13.20 (12.02-14.27) | 21.57 (20.67-22.52) | 16.49 (14.89-17.88) | 18.68 (17.94-19.52) | 14.21 (12.90-15.60) |
| Portugal | 17.57 (17.36-17.79) | 13.32 (12.17-14.41) | 14.26 (14.04-14.47) | 10.96 (9.99-11.76) | 20.21 (19.95-20.46) | 15.36 (13.97-16.55) | 16.50 (16.25-16.75) | 12.76 (11.70-13.73) | 21.52 (21.01-22.08) | 16.43 (14.99-17.78) | 17.82 (17.32-18.37) | 13.82 (12.60-14.92) |

(Table 3 continues on next page)

| | 1990, at age 65 years | | | | 2006, at age 65 years | | | | 2016, at age 65 years | | | |
|---|-------------------------------|-------------------------------|-------------------------------|----------------------------|-------------------------------|-------------------------------|-------------------------------|----------------------------|-------------------------------|-------------------------------|-------------------------------|------------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Spain | 19.26 (19.12–19.40) | 14.71 (13.40–15.84) | 15.59 (15.44–15.74) | 12.07 (11.09–12.98) | 21.58 (21.44–21.70) | 16.53 (15.08–17.79) | 17.57 (17.43–17.71) | 13.65 (12.54–14.65) | 22.82 (22.47–23.17) | 17.57 (16.08–18.94) | 19.17 (18.76–19.56) | 14.99 (13.74–16.09) |
| Sweden | 19.05 (18.78–19.30) | 14.38 (13.12–15.57) | 15.34 (15.10–15.60) | 11.77 (10.74–12.66) | 20.58 (20.36–20.83) | 15.53 (14.13–16.77) | 17.58 (17.35–17.81) | 13.36 (12.20–14.42) | 21.37 (20.36–22.32) | 16.05 (14.52–17.62) | 18.64 (17.74–19.51) | 14.12 (12.67–15.40) |
| Switzerland | 19.57 (19.09–20.04) | 14.64 (13.24–15.95) | 15.38 (14.93–15.84) | 11.70 (10.61–12.72) | 21.55 (21.10–21.98) | 16.14 (14.60–17.55) | 18.22 (17.74–18.69) | 13.82 (12.60–15.03) | 22.51 (20.50–24.40) | 17.02 (14.97–18.88) | 19.47 (17.46–21.31) | 14.90 (12.94–16.63) |
| UK | 17.80 (17.71–17.89) | 13.43 (12.25–14.50) | 13.98 (13.90–14.05) | 10.69 (9.78–11.49) | 19.88 (19.79–19.97) | 15.01 (13.68–16.19) | 17.10 (17.02–17.18) | 13.08 (11.97–14.10) | 20.88 (20.73–21.04) | 15.77 (14.37–17.07) | 18.27 (18.13–18.40) | 14.00 (12.83–15.05) |
| England | 17.94 (17.87–18.00) | 13.51 (12.31–14.61) | 14.11 (14.05–14.17) | 10.78 (9.86–11.60) | 20.03 (19.97–20.10) | 15.09 (13.74–16.30) | 17.24 (17.18–17.30) | 13.17 (12.07–14.18) | 21.05 (20.94–21.16) | 15.87 (14.44–17.17) | 18.44 (18.34–18.54) | 14.11 (12.91–15.18) |
| Northern Ireland | 17.39 (16.77–18.11) | 13.25 (12.04–14.43) | 13.52 (12.95–14.16) | 10.41 (9.47–11.31) | 19.69 (19.11–20.35) | 15.00 (13.66–16.35) | 16.70 (16.11–17.30) | 12.86 (11.72–13.93) | 20.59 (19.45–21.77) | 15.71 (14.13–17.25) | 17.75 (16.64–18.93) | 13.68 (12.32–15.06) |
| Scotland | 16.75 (16.13–17.38) | 12.77 (11.63–13.89) | 12.99 (12.46–13.53) | 10.02 (9.10–10.89) | 18.69 (18.06–19.34) | 14.30 (13.06–15.50) | 16.00 (15.44–16.53) | 12.38 (11.31–13.40) | 19.71 (18.66–20.75) | 15.02 (13.60–16.58) | 17.16 (16.23–18.09) | 13.24 (11.94–14.55) |
| Wales | 17.67 (17.07–18.28) | 13.40 (12.18–14.59) | 13.68 (13.14–14.18) | 10.51 (9.55–11.40) | 19.67 (19.19–20.26) | 15.00 (13.70–16.27) | 16.92 (16.34–17.45) | 13.03 (11.95–14.11) | 20.43 (19.42–21.41) | 15.58 (14.14–17.10) | 17.70 (16.71–18.66) | 13.65 (12.32–14.97) |
| Southern Latin America | 17.74 (17.50–17.99) | 13.79 (12.63–14.82) | 14.03 (13.79–14.27) | 10.84 (9.94–11.67) | 19.34 (19.11–19.60) | 15.03 (13.83–16.15) | 15.30 (15.05–15.55) | 11.83 (10.86–12.74) | 20.24 (19.54–20.92) | 15.72 (14.32–16.97) | 16.14 (15.49–16.77) | 12.50 (11.38–13.58) |
| Argentina | 17.76 (17.46–18.08) | 13.83 (12.70–14.89) | 13.90 (13.61–14.19) | 10.75 (9.84–11.59) | 19.01 (18.70–19.33) | 14.82 (13.65–15.91) | 14.81 (14.50–15.12) | 11.47 (10.52–12.36) | 19.73 (19.11–20.39) | 15.37 (14.05–16.56) | 15.55 (14.94–16.19) | 12.06 (11.04–13.05) |
| Chile | 17.69 (17.18–18.26) | 13.64 (12.46–14.77) | 14.65 (14.15–15.19) | 11.25 (10.19–12.25) | 20.40 (19.90–20.91) | 15.72 (14.40–16.95) | 17.08 (16.61–17.62) | 13.14 (11.99–14.23) | 21.63 (19.39–23.86) | 16.71 (14.58–18.88) | 17.91 (15.85–20.06) | 13.84 (11.93–15.86) |
| Uruguay | 17.70 (17.44–17.95) | 13.78 (12.64–14.77) | 13.69 (13.46–13.92) | 10.65 (9.75–11.44) | 19.50 (19.24–19.74) | 15.17 (13.95–16.22) | 14.76 (14.54–15.01) | 11.46 (10.49–12.30) | 20.44 (19.85–21.02) | 15.87 (14.53–17.10) | 15.54 (15.03–16.08) | 12.09 (11.04–13.07) |
| Central Europe, eastern Europe, and central Asia | 15.81 (15.50–16.12) | 11.59 (10.43–12.70) | 12.43 (12.17–12.66) | 9.08 (8.14–9.92) | 16.22 (15.89–16.52) | 11.99 (10.77–13.05) | 12.41 (12.12–12.70) | 9.12 (8.21–9.96) | 17.80 (16.51–18.92) | 13.16 (11.60–14.68) | 14.07 (13.21–14.96) | 10.33 (9.14–11.60) |
| Eastern Europe | 15.85 (15.36–16.35) | 11.64 (10.50–12.82) | 12.20 (11.76–12.66) | 8.92 (8.01–9.79) | 15.81 (15.30–16.30) | 11.74 (10.53–12.79) | 11.57 (11.10–12.05) | 8.53 (7.66–9.39) | 17.37 (15.46–19.28) | 12.90 (10.99–14.77) | 13.26 (11.85–14.92) | 9.79 (8.39–11.34) |
| Belarus | 16.52 (16.15–16.86) | 12.23 (11.04–13.34) | 12.94 (12.58–13.28) | 9.56 (8.60–10.48) | 16.27 (15.89–16.65) | 12.10 (10.89–13.16) | 11.53 (11.20–11.88) | 8.54 (7.67–9.35) | 18.15 (16.70–19.71) | 13.45 (11.83–15.17) | 13.25 (12.01–14.60) | 9.79 (8.48–11.19) |
| Estonia | 16.11 (15.79–16.44) | 11.79 (10.58–12.88) | 12.21 (11.91–12.52) | 8.91 (7.98–9.77) | 18.25 (17.93–18.60) | 13.55 (12.24–14.75) | 13.28 (12.96–13.57) | 9.78 (8.80–10.71) | 20.14 (19.43–21.38) | 14.90 (13.28–16.52) | 15.65 (14.85–16.60) | 11.50 (10.22–12.74) |
| Latvia | 15.93 (15.55–16.30) | 11.58 (10.34–12.70) | 12.23 (11.83–12.64) | 8.87 (7.93–9.75) | 16.87 (16.49–17.26) | 12.49 (11.28–13.60) | 12.38 (12.03–12.74) | 9.09 (8.16–9.96) | 19.04 (18.02–20.10) | 14.02 (12.56–15.56) | 14.24 (13.32–15.35) | 10.44 (9.21–11.68) |
| Lithuania | 17.10 (16.81–17.39) | 12.49 (11.23–13.64) | 13.47 (13.18–13.78) | 9.82 (8.81–10.74) | 17.97 (17.68–18.24) | 13.14 (11.83–14.34) | 13.26 (12.99–13.56) | 9.69 (8.69–10.62) | 19.63 (19.02–20.27) | 14.42 (12.96–15.79) | 14.70 (14.11–15.31) | 10.68 (9.43–11.81) |
| Moldova | 14.43 (13.82–15.08) | 10.67 (9.61–11.75) | 12.09 (11.51–12.71) | 8.94 (7.98–9.83) | 14.59 (13.99–15.27) | 11.00 (9.86–12.05) | 11.89 (11.28–12.52) | 8.95 (8.01–9.88) | 16.35 (15.32–17.49) | 12.29 (10.95–13.68) | 13.35 (12.40–14.45) | 10.01 (8.85–11.26) |

(Table 3 continues on next page)

| | 1990, at age 65 years | | | | 2006, at age 65 years | | | | 2016, at age 65 years | | | |
|--------------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Russia | 15.82 (15.10-16.62) | 11.60 (10.43-12.83) | 12.01 (11.35-12.78) | 8.74 (7.77-9.73) | 15.80 (15.04-16.52) | 11.70 (10.45-12.88) | 11.46 (10.77-12.20) | 8.41 (7.47-9.38) | 17.27 (14.52-20.14) | 12.84 (10.56-15.28) | 13.18 (11.27-15.67) | 9.72 (7.95-11.73) |
| Ukraine | 15.79 (15.35-16.24) | 11.65 (10.50-12.72) | 12.41 (12.02-12.80) | 9.15 (8.19-10.04) | 15.60 (15.19-16.00) | 11.65 (10.49-12.71) | 11.64 (11.27-12.04) | 8.66 (7.81-9.49) | 17.23 (15.13-19.48) | 12.93 (11.04-15.00) | 13.26 (11.61-15.48) | 9.92 (8.44-11.63) |
| Central Europe | 15.72 (15.58-15.85) | 11.42 (10.24-12.46) | 12.64 (12.52-12.77) | 9.16 (8.18-10.03) | 17.49 (17.36-17.64) | 12.79 (11.50-13.94) | 14.03 (13.90-14.16) | 10.19 (9.12-11.12) | 18.92 (18.59-19.25) | 13.84 (12.43-15.11) | 15.38 (15.08-15.69) | 11.16 (10.02-12.19) |
| Albania | 18.27 (18.06-18.51) | 13.70 (12.40-14.86) | 14.91 (14.71-15.13) | 11.14 (10.05-12.10) | 17.66 (17.41-17.91) | 13.33 (12.10-14.47) | 14.96 (14.76-15.17) | 11.26 (10.18-12.19) | 19.59 (18.61-20.63) | 14.70 (13.25-16.22) | 16.11 (14.93-17.27) | 12.07 (10.68-13.40) |
| Bosnia and Herzegovina | 15.47 (14.56-16.56) | 11.53 (10.22-12.78) | 12.83 (12.00-13.72) | 9.57 (8.51-10.72) | 17.32 (16.38-18.35) | 12.84 (11.52-14.23) | 14.66 (13.73-15.57) | 10.78 (9.57-11.90) | 18.56 (17.17-19.95) | 13.69 (12.11-15.33) | 15.58 (14.44-16.83) | 11.39 (10.05-12.76) |
| Bulgaria | 15.20 (14.94-15.45) | 11.24 (10.16-12.23) | 12.81 (12.58-13.03) | 9.34 (8.39-10.22) | 16.22 (15.99-16.46) | 12.07 (10.92-13.12) | 13.17 (12.93-13.41) | 9.71 (8.75-10.59) | 17.60 (16.23-18.93) | 13.07 (11.52-14.58) | 14.28 (13.12-15.58) | 10.50 (9.26-11.91) |
| Croatia | 15.96 (15.35-16.56) | 11.68 (10.40-12.87) | 12.72 (12.18-13.36) | 9.26 (8.25-10.23) | 17.50 (16.97-18.04) | 12.88 (11.53-14.10) | 13.90 (13.41-14.44) | 10.18 (9.10-11.21) | 18.57 (17.49-19.78) | 13.66 (12.08-15.13) | 15.15 (14.20-16.27) | 11.04 (9.76-12.45) |
| Czech Republic | 15.33 (15.16-15.49) | 10.84 (9.61-11.93) | 11.84 (11.68-12.01) | 8.44 (7.49-9.26) | 18.09 (17.89-18.26) | 12.92 (11.51-14.19) | 14.69 (14.51-14.89) | 10.43 (9.25-11.46) | 19.66 (19.21-20.11) | 14.10 (12.46-15.53) | 16.34 (15.88-16.79) | 11.62 (10.26-12.83) |
| Hungary | 15.43 (15.02-15.87) | 10.85 (9.57-12.01) | 12.08 (11.72-12.46) | 8.58 (7.59-9.51) | 17.48 (17.05-17.92) | 12.56 (11.22-13.85) | 13.54 (13.12-13.98) | 9.70 (8.60-10.69) | 18.34 (17.45-19.21) | 13.25 (11.82-14.70) | 14.74 (13.88-15.67) | 10.61 (9.36-11.91) |
| Macedonia | 14.55 (14.03-15.08) | 10.86 (9.76-11.88) | 12.61 (12.18-13.04) | 9.36 (8.42-10.25) | 14.93 (14.49-15.43) | 11.14 (10.05-12.19) | 12.71 (12.32-13.13) | 9.44 (8.50-10.33) | 16.26 (15.79-16.82) | 12.08 (10.86-13.19) | 13.67 (13.22-14.15) | 10.13 (9.08-11.06) |
| Montenegro | 18.05 (17.25-18.70) | 13.38 (12.05-14.73) | 15.39 (14.78-16.01) | 11.28 (10.01-12.38) | 17.80 (17.45-18.18) | 13.23 (11.83-14.45) | 14.76 (14.34-15.23) | 10.88 (9.78-11.93) | 18.50 (17.59-19.69) | 13.73 (12.17-15.34) | 15.72 (14.87-16.56) | 11.56 (10.30-12.82) |
| Poland | 16.07 (15.76-16.37) | 11.70 (10.48-12.78) | 12.40 (12.15-12.70) | 9.02 (8.08-9.89) | 18.64 (18.37-18.94) | 13.56 (12.18-14.84) | 14.52 (14.22-14.83) | 10.51 (9.40-11.53) | 20.10 (19.32-20.89) | 14.69 (13.06-16.19) | 16.03 (15.32-16.80) | 11.61 (10.31-12.85) |
| Romania | 15.37 (15.00-15.76) | 11.32 (10.16-12.37) | 13.17 (12.78-13.52) | 9.57 (8.57-10.52) | 16.86 (16.48-17.27) | 12.44 (11.23-13.61) | 13.89 (13.54-14.25) | 10.13 (9.06-11.13) | 18.11 (17.30-18.94) | 13.40 (12.02-14.73) | 14.78 (13.93-15.63) | 10.79 (9.53-12.02) |
| Serbia | 16.12 (15.50-16.90) | 11.85 (10.61-13.13) | 13.48 (12.93-14.25) | 9.88 (8.82-10.96) | 15.81 (15.34-16.26) | 11.67 (10.49-12.76) | 13.42 (13.05-13.85) | 9.84 (8.80-10.81) | 17.87 (17.46-18.22) | 13.12 (11.77-14.30) | 14.95 (14.58-15.45) | 10.92 (9.77-11.97) |
| Slovakia | 15.65 (15.33-15.99) | 11.39 (10.22-12.44) | 12.14 (11.83-12.46) | 8.74 (7.76-9.58) | 17.30 (16.98-17.65) | 12.75 (11.47-13.88) | 13.46 (13.14-13.76) | 9.76 (8.70-10.71) | 18.79 (17.84-19.81) | 13.84 (12.34-15.36) | 15.07 (14.16-16.02) | 10.90 (9.55-12.23) |
| Slovenia | 16.98 (16.32-17.66) | 12.11 (10.74-13.47) | 13.24 (12.67-13.90) | 9.42 (8.27-10.46) | 19.67 (19.06-20.38) | 14.01 (12.47-15.47) | 15.53 (14.86-16.18) | 10.97 (9.67-12.25) | 21.28 (20.30-22.41) | 15.12 (13.28-16.79) | 17.51 (16.48-18.62) | 12.44 (10.88-14.05) |
| Central Asia | 15.86 (15.55-16.14) | 11.82 (10.69-12.85) | 12.81 (12.53-13.07) | 9.64 (8.74-10.45) | 15.03 (14.74-15.32) | 11.29 (10.24-12.21) | 12.04 (11.76-12.30) | 9.13 (8.31-9.91) | 16.82 (16.31-17.33) | 12.58 (11.34-13.68) | 13.51 (13.06-14.02) | 10.21 (9.17-11.16) |
| Armenia | 16.36 (15.90-16.84) | 12.21 (11.01-13.31) | 13.96 (13.48-14.47) | 10.43 (9.38-11.39) | 16.46 (15.89-16.98) | 12.35 (11.12-13.46) | 13.65 (13.18-14.16) | 10.26 (9.26-11.21) | 18.31 (17.42-19.33) | 13.70 (12.15-15.07) | 14.76 (14.01-15.62) | 11.09 (9.93-12.20) |
| Azerbaijan | 15.87 (15.41-16.47) | 11.96 (10.82-13.07) | 12.45 (12.04-12.87) | 9.48 (8.59-10.34) | 14.90 (14.14-15.75) | 11.31 (10.09-12.42) | 12.22 (11.59-12.93) | 9.37 (8.51-10.26) | 17.26 (15.65-18.62) | 12.98 (11.35-14.41) | 13.75 (12.53-15.42) | 10.48 (9.13-12.11) |

(Table 3 continues on next page)

| | 1990, at age 65 years | | | | 2006, at age 65 years | | | | 2016, at age 65 years | | | |
|------------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Georgia | 16-28 (15-61-16-95) | 12-30 (11-04-13-42) | 13-14 (12-46-13-87) | 9-94 (8-92-10-88) | 17-77 (17-33-18-20) | 13-44 (12-20-14-58) | 13-26 (12-81-13-78) | 10-12 (9-16-11-03) | 18-50 (17-13-20-32) | 13-91 (12-39-15-46) | 13-78 (12-62-15-14) | 10-40 (9-08-11-74) |
| Kazakhstan | 15-94 (15-11-16-76) | 11-75 (10-44-12-93) | 12-21 (11-48-13-01) | 9-09 (8-03-10-08) | 14-72 (13-98-15-51) | 10-92 (9-69-12-05) | 10-91 (10-23-11-64) | 8-13 (7-20-9-07) | 16-95 (15-49-18-45) | 12-54 (11-03-14-06) | 13-26 (11-89-14-72) | 9-86 (8-52-11-34) |
| Kyrgyzstan | 15-81 (15-45-16-19) | 11-73 (10-58-12-80) | 12-46 (12-14-12-82) | 9-33 (8-46-10-13) | 14-41 (14-08-14-76) | 10-85 (9-85-11-76) | 11-52 (11-18-11-83) | 8-75 (7-93-9-49) | 16-64 (15-84-17-41) | 12-47 (11-20-13-72) | 13-55 (12-87-14-30) | 10-27 (9-21-11-28) |
| Mongolia | 14-23 (13-32-15-22) | 10-73 (9-49-11-90) | 12-20 (11-48-12-78) | 9-26 (8-31-10-19) | 13-75 (13-13-14-34) | 10-36 (9-33-11-37) | 11-11 (10-62-11-75) | 8-40 (7-58-9-23) | 15-61 (14-55-16-74) | 11-69 (10-34-13-01) | 12-29 (11-45-13-25) | 9-20 (8-10-10-27) |
| Tajikistan | 15-87 (15-14-16-68) | 11-88 (10-66-13-07) | 13-64 (12-79-14-49) | 10-28 (9-21-11-35) | 14-52 (13-56-15-40) | 11-01 (9-92-12-18) | 13-00 (12-29-13-80) | 9-92 (8-89-10-94) | 16-92 (15-66-17-89) | 12-78 (11-34-14-08) | 14-90 (13-58-15-85) | 11-34 (10-05-12-50) |
| Turkmenistan | 15-09 (14-84-15-36) | 11-39 (10-31-12-35) | 12-42 (12-23-12-61) | 9-49 (8-65-10-24) | 15-60 (15-30-15-91) | 11-86 (10-77-12-87) | 12-73 (12-48-12-97) | 9-79 (8-93-10-57) | 16-95 (16-34-17-52) | 12-85 (11-60-14-03) | 13-73 (13-31-14-15) | 10-54 (9-62-11-42) |
| Uzbekistan | 15-72 (15-18-16-23) | 11-69 (10-51-12-80) | 13-19 (12-71-13-72) | 9-97 (9-00-10-89) | 14-18 (13-65-14-77) | 10-68 (9-59-11-72) | 11-93 (11-30-12-50) | 9-11 (8-24-10-01) | 15-76 (14-90-16-65) | 11-84 (10-59-13-07) | 13-13 (12-43-13-80) | 10-02 (8-97-10-98) |
| Latin America and Caribbean | 17-13 (17-03-17-22) | 12-97 (11-81-13-99) | 14-69 (14-60-14-78) | 11-18 (10-21-12-06) | 18-73 (18-62-18-83) | 14-17 (12-89-15-31) | 16-05 (15-95-16-14) | 12-21 (11-13-13-19) | 19-49 (19-29-19-68) | 14-74 (13-39-15-94) | 16-78 (16-61-16-95) | 12-78 (11-65-13-77) |
| Central Latin America | 17-96 (17-84-18-09) | 13-61 (12-38-14-68) | 15-76 (15-64-15-89) | 11-94 (10-84-12-89) | 19-16 (19-01-19-32) | 14-60 (13-31-15-75) | 16-71 (16-55-16-86) | 12-72 (11-58-13-73) | 19-75 (19-46-20-07) | 15-03 (13-67-16-25) | 17-30 (17-00-17-58) | 13-17 (12-00-14-22) |
| Colombia | 17-85 (17-57-18-11) | 13-69 (12-44-14-78) | 15-41 (15-17-15-66) | 11-86 (10-81-12-78) | 18-65 (18-41-18-89) | 14-40 (13-23-15-45) | 16-03 (15-78-16-27) | 12-44 (11-41-13-35) | 20-24 (19-48-21-04) | 15-61 (14-17-16-91) | 17-58 (16-85-18-33) | 13-60 (12-36-14-77) |
| Costa Rica | 18-86 (18-53-19-21) | 14-42 (13-13-15-58) | 16-38 (16-04-16-70) | 12-66 (11-59-13-65) | 21-26 (20-90-21-58) | 16-21 (14-75-17-52) | 18-52 (18-19-18-85) | 14-24 (13-01-15-37) | 22-29 (21-64-22-94) | 16-99 (15-40-18-49) | 19-50 (18-88-20-18) | 15-00 (13-61-16-28) |
| El Salvador | 17-51 (17-14-17-91) | 13-43 (12-21-14-52) | 15-26 (14-91-15-61) | 11-67 (10-65-12-61) | 18-74 (18-39-19-10) | 14-36 (13-07-15-50) | 17-54 (17-20-17-89) | 13-35 (12-13-14-50) | 19-56 (18-61-20-48) | 14-99 (13-49-16-35) | 17-95 (17-06-18-90) | 13-68 (12-29-14-98) |
| Guatemala | 15-45 (14-53-16-45) | 11-75 (10-45-13-01) | 14-76 (13-91-15-90) | 11-23 (10-11-12-51) | 18-05 (16-81-19-26) | 13-60 (12-08-15-08) | 16-33 (15-32-17-47) | 12-40 (11-06-13-85) | 18-76 (16-58-21-20) | 14-24 (12-19-16-30) | 16-86 (15-14-18-96) | 12-84 (11-08-14-83) |
| Honduras | 14-00 (13-26-14-84) | 10-79 (9-76-11-81) | 15-71 (14-10-16-95) | 12-11 (10-76-13-48) | 15-24 (13-70-17-73) | 11-69 (10-10-13-62) | 15-29 (13-25-17-63) | 11-85 (9-93-13-82) | 16-03 (14-34-18-60) | 12-29 (10-60-14-36) | 15-73 (13-50-17-94) | 12-16 (10-19-14-03) |
| Mexico | 18-45 (18-33-18-59) | 13-87 (12-60-15-00) | 16-04 (15-91-16-18) | 12-00 (10-87-13-00) | 19-42 (19-24-19-59) | 14-71 (13-38-15-92) | 16-91 (16-77-17-05) | 12-74 (11-55-13-76) | 19-48 (19-23-19-75) | 14-72 (13-35-15-93) | 17-20 (16-99-17-42) | 12-97 (11-74-14-04) |
| Nicaragua | 20-99 (20-64-21-35) | 15-97 (14-58-17-26) | 18-21 (17-86-18-57) | 13-94 (12-68-15-09) | 20-55 (20-18-20-95) | 15-73 (14-32-17-00) | 18-29 (17-93-18-63) | 14-08 (12-86-15-23) | 20-98 (19-63-22-42) | 16-05 (14-41-17-73) | 18-38 (17-12-19-75) | 14-16 (12-53-15-71) |
| Panama | 18-86 (18-27-19-48) | 14-45 (13-09-15-71) | 16-24 (15-62-16-85) | 12-49 (11-36-13-54) | 20-69 (20-12-21-25) | 15-85 (14-35-17-10) | 18-01 (17-40-18-61) | 13-82 (12-58-14-95) | 21-73 (20-64-22-85) | 16-60 (14-95-18-18) | 18-76 (17-55-19-96) | 14-32 (12-83-15-78) |
| Venezuela | 17-29 (16-76-17-88) | 13-21 (11-94-14-34) | 15-08 (14-50-15-59) | 11-56 (10-48-12-58) | 19-62 (19-01-20-20) | 15-00 (13-59-16-26) | 16-42 (15-91-16-98) | 12-62 (11-48-13-67) | 20-19 (18-68-21-62) | 15-47 (13-72-17-22) | 16-71 (15-27-17-97) | 12-85 (11-36-14-26) |
| Andean Latin America | 17-23 (16-84-17-65) | 13-14 (11-88-14-26) | 15-71 (15-35-16-06) | 11-97 (10-91-12-93) | 19-66 (19-22-20-09) | 15-03 (13-66-16-22) | 17-61 (17-23-18-01) | 13-45 (12-21-14-55) | 20-12 (19-15-21-07) | 15-39 (13-86-16-86) | 18-36 (17-53-19-23) | 14-04 (12-63-15-41) |

(Table 3 continues on next page)

| | 1990, at age 65 years | | | | 2006, at age 65 years | | | | 2016, at age 65 years | | | |
|----------------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Bolivia | 13.86 (12.86-14.74) | 10.46 (9.25-11.63) | 13.76 (13.09-14.45) | 10.31 (9.25-11.31) | 16.68 (15.58-17.39) | 12.62 (11.19-13.88) | 15.79 (15.11-16.68) | 11.92 (10.69-13.14) | 16.93 (14.96-18.57) | 12.88 (10.99-14.55) | 16.60 (15.49-18.03) | 12.55 (11.04-14.20) |
| Ecuador | 18.27 (17.92-18.63) | 13.99 (12.74-15.13) | 16.17 (15.84-16.50) | 12.33 (11.20-13.34) | 20.21 (19.87-20.58) | 15.46 (14.06-16.68) | 17.90 (17.50-18.28) | 13.57 (12.33-14.73) | 20.84 (20.21-21.44) | 15.93 (14.44-17.28) | 18.67 (18.02-19.29) | 14.19 (12.88-15.46) |
| Peru | 18.20 (17.64-18.79) | 13.91 (12.58-15.12) | 16.24 (15.61-16.83) | 12.45 (11.36-13.53) | 20.54 (19.91-21.30) | 15.75 (14.26-17.02) | 18.13 (17.46-18.81) | 13.96 (12.60-15.15) | 21.06 (19.44-22.77) | 16.17 (14.35-18.06) | 18.88 (17.36-20.52) | 14.55 (12.85-16.20) |
| Caribbean | 16.54 (16.30-16.77) | 12.69 (11.53-13.70) | 15.12 (14.93-15.31) | 11.66 (10.65-12.59) | 17.99 (17.59-18.41) | 13.73 (12.50-14.87) | 16.16 (15.91-16.45) | 12.40 (11.31-13.40) | 18.59 (18.05-19.17) | 14.19 (12.92-15.40) | 16.50 (16.08-16.90) | 12.66 (11.51-13.71) |
| Antigua and Barbuda | 17.09 (16.33-17.88) | 13.17 (11.91-14.43) | 14.14 (13.51-14.78) | 10.84 (9.86-11.83) | 19.02 (18.29-19.74) | 14.52 (13.08-15.89) | 16.29 (15.63-17.05) | 12.44 (11.23-13.63) | 19.74 (18.72-20.85) | 15.10 (13.53-16.61) | 16.79 (15.84-17.78) | 12.81 (11.51-14.18) |
| The Bahamas | 16.94 (16.60-17.27) | 13.01 (11.83-14.05) | 13.96 (13.64-14.31) | 10.79 (9.87-11.67) | 17.81 (17.35-18.34) | 13.66 (12.44-14.78) | 15.53 (15.21-15.85) | 11.99 (10.96-12.92) | 17.58 (16.57-18.78) | 13.50 (12.09-14.92) | 15.74 (15.06-16.52) | 12.16 (11.02-13.37) |
| Barbados | 17.20 (16.69-17.72) | 13.19 (11.94-14.35) | 14.19 (13.68-14.60) | 10.95 (9.96-11.88) | 19.40 (18.87-19.90) | 14.74 (13.35-15.99) | 16.36 (15.86-16.88) | 12.53 (11.41-13.56) | 19.37 (18.61-20.22) | 14.70 (13.25-16.03) | 16.74 (15.99-17.50) | 12.85 (11.63-14.06) |
| Belize | 17.21 (16.26-18.14) | 13.19 (11.80-14.46) | 15.55 (14.28-16.17) | 12.02 (10.79-13.10) | 16.04 (15.65-16.40) | 12.21 (11.10-13.28) | 13.46 (12.58-14.25) | 10.35 (9.26-11.43) | 16.61 (15.62-17.69) | 12.67 (11.31-14.03) | 14.05 (13.07-15.19) | 10.82 (9.66-11.98) |
| Bermuda | 13.93 (13.33-14.58) | 10.77 (9.76-11.75) | 12.47 (11.92-13.01) | 9.61 (8.67-10.44) | 16.90 (16.17-17.60) | 13.00 (11.83-14.16) | 14.73 (14.17-15.26) | 11.35 (10.36-12.31) | 20.42 (19.08-22.13) | 15.70 (14.10-17.46) | 16.31 (15.32-17.25) | 12.57 (11.28-13.80) |
| Cuba | 18.18 (17.83-18.54) | 13.94 (12.70-15.07) | 15.72 (15.35-16.04) | 12.19 (11.13-13.19) | 19.45 (19.10-19.79) | 14.87 (13.50-16.07) | 16.88 (16.52-17.24) | 13.07 (11.93-14.09) | 20.23 (19.53-21.00) | 15.44 (13.98-16.78) | 17.31 (16.65-18.00) | 13.37 (12.10-14.57) |
| Dominica | 14.93 (14.41-15.55) | 11.56 (10.49-12.53) | 13.63 (13.14-14.30) | 10.54 (9.61-11.48) | 16.65 (15.84-17.31) | 12.83 (11.63-13.89) | 14.38 (13.82-15.01) | 11.06 (10.03-12.02) | 17.26 (16.02-18.20) | 13.30 (11.90-14.66) | 14.63 (13.65-15.80) | 11.25 (9.98-12.51) |
| Dominican Republic | 17.36 (16.88-18.08) | 13.47 (12.25-14.65) | 16.05 (15.48-16.60) | 12.50 (11.38-13.60) | 19.45 (18.41-20.68) | 15.02 (13.38-16.68) | 16.81 (16.34-17.89) | 12.98 (11.82-14.26) | 19.65 (18.48-21.24) | 15.19 (13.66-17.00) | 17.21 (16.33-18.48) | 13.33 (12.00-14.63) |
| Grenada | 15.64 (14.97-16.19) | 11.90 (10.74-12.91) | 13.71 (12.95-14.40) | 10.50 (9.44-11.58) | 16.05 (15.32-16.71) | 12.17 (10.94-13.21) | 13.79 (13.16-14.36) | 10.54 (9.50-11.48) | 16.29 (15.19-17.62) | 12.37 (11.08-13.68) | 13.96 (13.11-14.93) | 10.67 (9.53-11.79) |
| Guyana | 14.71 (14.21-15.34) | 11.22 (10.18-12.19) | 11.79 (11.24-12.38) | 9.03 (8.16-9.91) | 14.36 (13.76-14.92) | 10.87 (9.76-11.87) | 12.34 (11.84-12.84) | 9.35 (8.43-10.20) | 15.17 (14.32-16.19) | 11.49 (10.28-12.70) | 12.98 (12.28-13.74) | 9.88 (8.86-10.85) |
| Haiti | 10.84 (10.21-11.40) | 8.24 (7.41-9.09) | 12.30 (11.78-12.82) | 9.34 (8.43-10.24) | 12.27 (11.18-13.67) | 9.35 (8.20-10.54) | 13.25 (12.48-14.47) | 10.10 (8.98-11.30) | 13.04 (11.71-14.55) | 9.95 (8.74-11.38) | 13.75 (12.36-15.23) | 10.52 (9.14-11.91) |
| Jamaica | 17.83 (17.01-18.62) | 13.63 (12.25-14.93) | 16.83 (16.27-17.47) | 12.99 (11.83-14.11) | 18.14 (17.52-18.87) | 13.71 (12.37-15.01) | 16.79 (16.09-17.44) | 12.85 (11.59-13.95) | 17.93 (16.46-19.36) | 13.55 (11.91-15.24) | 16.28 (14.77-17.58) | 12.45 (10.95-13.86) |
| Puerto Rico | 18.89 (18.50-19.30) | 14.53 (13.27-15.68) | 15.88 (15.48-16.28) | 12.17 (11.08-13.17) | 20.52 (20.09-20.90) | 15.59 (14.12-16.90) | 17.38 (16.94-17.76) | 13.14 (11.88-14.26) | 21.13 (20.34-21.98) | 16.10 (14.54-17.53) | 17.57 (16.86-18.31) | 13.29 (12.00-14.48) |
| Saint Lucia | 17.07 (16.72-17.43) | 12.89 (11.64-14.01) | 14.99 (14.66-15.35) | 11.40 (10.43-12.39) | 18.95 (18.51-19.41) | 14.43 (13.08-15.70) | 16.32 (15.93-16.73) | 12.49 (11.38-13.53) | 20.02 (19.49-20.59) | 15.26 (13.84-16.59) | 17.16 (16.60-17.66) | 13.15 (11.94-14.26) |
| Saint Vincent and the Grenadines | 15.68 (15.08-16.14) | 11.96 (10.81-12.98) | 13.84 (13.14-15.13) | 10.63 (9.53-11.84) | 16.03 (15.63-16.49) | 12.14 (11.00-13.20) | 14.02 (13.42-14.51) | 10.66 (9.60-11.67) | 16.39 (15.68-17.33) | 12.45 (11.18-13.74) | 14.02 (13.13-14.60) | 10.64 (9.49-11.70) |

(Table 3 continues on next page)

| | 1990, at age 65 years | | | | 2006, at age 65 years | | | | 2016, at age 65 years | | | |
|---|--------------------------------|--------------------------------|--------------------------------|------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|--------------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Suriname | 16.97 (16.61-17.36) | 12.97 (11.75-14.08) | 15.17 (14.98-15.38) | 11.64 (10.64-12.56) | 16.22 (15.66-16.64) | 12.32 (11.14-13.38) | 14.05 (13.80-14.37) | 10.74 (9.75-11.63) | 17.14 (16.40-17.75) | 12.99 (11.71-14.22) | 14.49 (13.84-15.34) | 11.05 (9.99-12.13) |
| Trinidad and Tobago | 15.44 (15.10-15.81) | 11.69 (10.61-12.67) | 12.43 (12.14-12.73) | 9.44 (8.56-10.27) | 17.67 (17.25-18.11) | 13.29 (12.00-14.44) | 13.95 (13.63-14.31) | 10.40 (9.36-11.37) | 18.72 (18.05-19.41) | 14.08 (12.68-15.41) | 14.40 (13.83-15.02) | 10.71 (9.59-11.83) |
| Virgin Islands | 17.05 (16.58-18.07) | 13.21 (12.03-14.43) | 13.96 (13.56-14.49) | 10.80 (9.85-11.68) | 17.93 (17.52-18.31) | 13.79 (12.54-14.86) | 14.50 (13.96-14.99) | 11.12 (10.09-12.08) | 18.47 (17.86-19.07) | 14.17 (12.89-15.45) | 14.61 (13.76-15.71) | 11.18 (9.97-12.37) |
| Tropical Latin America | 16.41 (16.23-16.57) | 12.34 (11.22-13.34) | 13.16 (13.01-13.32) | 10.04 (9.15-10.87) | 18.28 (18.11-18.43) | 13.67 (12.37-14.83) | 15.00 (14.84-15.14) | 11.35 (10.35-12.29) | 19.31 (19.02-19.58) | 14.45 (13.09-15.67) | 15.99 (15.74-16.23) | 12.14 (11.05-13.14) |
| Brazil | 16.36 (16.18-16.52) | 12.30 (11.18-13.29) | 13.08 (12.93-13.24) | 9.97 (9.09-10.80) | 18.29 (18.12-18.45) | 13.68 (12.37-14.84) | 14.96 (14.81-15.11) | 11.32 (10.32-12.26) | 19.34 (19.05-19.61) | 14.48 (13.11-15.69) | 16.00 (15.75-16.24) | 12.14 (11.06-13.15) |
| Paraguay | 18.47 (17.94-18.88) | 13.98 (12.70-15.17) | 16.60 (16.23-17.00) | 12.70 (11.60-13.80) | 17.79 (17.10-18.51) | 13.43 (12.05-14.74) | 16.26 (16.01-16.45) | 12.38 (11.28-13.37) | 17.90 (17.02-18.91) | 13.45 (11.97-14.85) | 15.75 (14.68-16.54) | 12.03 (10.72-13.30) |
| Southeast Asia, east Asia, and Oceania | 14.53 (14.35-14.71) | 11.16 (10.28-12.02) | 12.44 (12.30-12.58) | 9.76 (9.02-10.43) | 16.47 (16.33-16.61) | 12.53 (11.46-13.49) | 14.06 (13.93-14.19) | 10.95 (10.09-11.75) | 18.48 (18.30-18.65) | 13.99 (12.81-15.07) | 15.09 (14.93-15.25) | 11.69 (10.79-12.52) |
| East Asia | 14.51 (14.29-14.73) | 11.23 (10.37-12.07) | 12.20 (12.03-12.37) | 9.68 (8.98-10.32) | 16.76 (16.60-16.93) | 12.83 (11.76-13.80) | 14.06 (13.92-14.20) | 11.05 (10.23-11.84) | 18.97 (18.76-19.18) | 14.43 (13.22-15.51) | 15.17 (15.00-15.35) | 11.85 (10.95-12.67) |
| China | 14.45 (14.22-14.68) | 11.19 (10.34-12.02) | 12.14 (11.97-12.31) | 9.64 (8.94-10.28) | 16.73 (16.57-16.90) | 12.81 (11.75-13.78) | 14.01 (13.86-14.15) | 11.02 (10.20-11.81) | 19.01 (18.79-19.22) | 14.45 (13.24-15.53) | 15.14 (14.97-15.32) | 11.83 (10.93-12.65) |
| North Korea | 16.18 (15.40-17.33) | 12.48 (11.32-13.76) | 14.00 (13.20-15.35) | 11.05 (9.93-12.35) | 15.59 (14.98-16.45) | 11.94 (10.80-13.08) | 13.41 (13.04-13.90) | 10.56 (9.72-11.32) | 15.93 (15.36-16.53) | 12.26 (11.20-13.38) | 13.49 (13.10-13.97) | 10.60 (9.77-11.46) |
| Taiwan (Province of China) | 16.63 (16.27-16.99) | 12.55 (11.40-13.63) | 14.61 (14.28-14.95) | 11.31 (10.35-12.19) | 19.85 (19.50-20.20) | 14.95 (13.63-16.14) | 17.05 (16.70-17.43) | 13.14 (12.09-14.12) | 20.99 (19.97-22.16) | 15.88 (14.32-17.50) | 17.87 (16.87-19.00) | 13.79 (12.50-15.10) |
| Southeast Asia | 14.63 (14.34-14.93) | 10.95 (9.93-11.91) | 13.39 (13.16-13.58) | 10.08 (9.14-10.88) | 15.69 (15.50-16.02) | 11.73 (10.66-12.70) | 14.13 (13.90-14.38) | 10.62 (9.71-11.47) | 17.13 (16.85-17.41) | 12.79 (11.63-13.88) | 14.84 (14.52-15.17) | 11.12 (10.12-12.02) |
| Cambodia | 11.93 (11.51-12.45) | 8.69 (7.75-9.61) | 11.75 (11.15-12.30) | 8.65 (7.71-9.57) | 13.52 (13.24-13.86) | 9.80 (8.78-10.72) | 12.67 (12.01-13.24) | 9.25 (8.30-10.18) | 15.07 (14.62-15.57) | 11.06 (9.97-12.11) | 13.07 (12.57-13.72) | 9.67 (8.71-10.55) |
| Indonesia | 13.72 (13.37-14.03) | 10.32 (9.32-11.21) | 13.37 (13.21-13.56) | 10.08 (9.19-10.90) | 14.55 (14.29-14.79) | 10.90 (9.87-11.81) | 14.29 (13.98-14.65) | 10.73 (9.75-11.62) | 15.81 (15.48-16.15) | 11.76 (10.69-12.76) | 14.60 (14.01-15.27) | 10.89 (9.83-11.92) |
| Laos | 11.73 (11.39-12.17) | 8.67 (7.78-9.51) | 11.77 (11.57-11.95) | 8.73 (7.87-9.50) | 13.68 (13.19-14.14) | 10.13 (9.09-11.06) | 12.67 (12.50-12.86) | 9.38 (8.48-10.23) | 15.17 (14.61-15.85) | 11.23 (10.07-12.33) | 13.73 (13.34-14.22) | 10.19 (9.23-11.12) |
| Malaysia | 14.92 (14.72-15.13) | 11.19 (10.19-12.12) | 13.52 (13.33-13.71) | 10.25 (9.33-11.06) | 15.32 (15.15-15.48) | 11.54 (10.47-12.46) | 14.82 (14.64-15.01) | 11.18 (10.22-12.06) | 16.88 (16.51-17.23) | 12.69 (11.48-13.75) | 15.29 (14.84-15.80) | 11.52 (10.48-12.49) |
| Maldives | 13.01 (12.41-13.75) | 9.59 (8.57-10.60) | 13.63 (13.03-14.33) | 10.18 (9.17-11.15) | 16.96 (16.25-17.74) | 12.59 (11.28-13.86) | 16.18 (15.50-16.86) | 12.14 (10.94-13.28) | 19.43 (17.36-21.50) | 14.45 (12.44-16.33) | 17.22 (15.48-19.16) | 12.97 (11.30-14.71) |
| Mauritius | 15.09 (14.74-15.43) | 11.24 (10.20-12.24) | 11.01 (10.75-11.27) | 8.24 (7.50-8.94) | 16.74 (16.36-17.09) | 12.41 (11.14-13.51) | 13.47 (13.17-13.81) | 9.99 (9.08-10.83) | 18.00 (16.79-19.13) | 13.27 (11.79-14.82) | 14.82 (13.88-15.88) | 10.92 (9.63-12.24) |
| Myanmar | 12.18 (11.76-12.67) | 9.03 (8.11-9.93) | 11.62 (10.81-12.30) | 8.65 (7.71-9.53) | 13.84 (13.40-14.39) | 10.27 (9.21-11.25) | 12.70 (11.00-13.42) | 9.44 (8.16-10.46) | 15.81 (15.37-16.33) | 11.77 (10.65-12.85) | 13.35 (12.79-13.87) | 9.93 (8.97-10.86) |

(Table 3 continues on next page)

| | 1990, at age 65 years | | | | 2006, at age 65 years | | | | 2016, at age 65 years | | | |
|--------------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Philippines | 16.15 (15.77-16.56) | 12.05 (10.91-13.14) | 13.10 (12.72-13.52) | 9.72 (8.81-10.60) | 15.31 (14.86-15.73) | 11.38 (10.31-12.43) | 11.96 (11.59-12.37) | 8.84 (8.01-9.64) | 15.97 (14.91-17.14) | 11.92 (10.64-13.33) | 12.57 (11.61-13.58) | 9.33 (8.33-10.43) |
| Sri Lanka | 17.52 (17.00-18.08) | 13.19 (11.99-14.37) | 14.96 (14.42-15.50) | 11.28 (10.21-12.32) | 18.32 (17.49-19.02) | 13.74 (12.46-14.96) | 14.95 (14.25-15.60) | 11.21 (10.13-12.29) | 19.70 (17.93-21.75) | 14.78 (12.86-16.74) | 16.19 (14.38-18.07) | 12.16 (10.51-13.86) |
| Seychelles | 16.08 (15.70-16.49) | 12.20 (11.15-13.23) | 12.46 (12.07-12.83) | 9.44 (8.59-10.20) | 17.04 (16.59-17.71) | 12.84 (11.62-14.03) | 13.77 (13.52-14.07) | 10.34 (9.39-11.21) | 17.73 (17.02-18.45) | 13.37 (12.11-14.63) | 14.42 (13.57-15.40) | 10.83 (9.78-11.97) |
| Thailand | 17.07 (16.70-17.44) | 12.73 (11.52-13.82) | 14.71 (14.35-15.08) | 11.06 (10.03-11.97) | 18.61 (18.26-18.95) | 13.94 (12.66-15.12) | 16.71 (16.32-17.12) | 12.60 (11.48-13.67) | 20.33 (19.48-21.10) | 15.20 (13.72-16.69) | 18.29 (17.42-19.15) | 13.72 (12.39-15.05) |
| Timor-Leste | 12.77 (11.35-14.28) | 9.51 (8.13-10.90) | 14.36 (13.39-15.51) | 10.59 (9.34-11.75) | 15.27 (13.91-16.46) | 11.29 (9.92-12.61) | 15.14 (14.16-16.34) | 11.09 (9.79-12.44) | 16.51 (14.75-18.09) | 12.30 (10.69-14.01) | 16.13 (14.64-17.71) | 11.85 (10.29-13.35) |
| Vietnam | 15.25 (14.17-16.51) | 11.45 (10.09-12.89) | 13.48 (12.46-14.35) | 10.26 (9.20-11.37) | 16.67 (15.90-18.27) | 12.50 (11.23-14.01) | 13.60 (12.77-14.67) | 10.35 (9.39-11.47) | 18.08 (17.34-18.54) | 13.64 (12.34-14.82) | 14.45 (13.89-15.11) | 11.02 (10.01-11.98) |
| Oceania | 11.30 (10.70-11.96) | 8.31 (7.36-9.20) | 10.56 (10.16-11.10) | 7.90 (7.10-8.67) | 11.38 (10.70-12.09) | 8.32 (7.36-9.27) | 10.91 (10.42-11.50) | 8.13 (7.29-8.97) | 12.01 (11.22-12.77) | 8.76 (7.75-9.74) | 11.45 (10.87-12.11) | 8.51 (7.63-9.39) |
| American Samoa | 16.43 (15.88-17.15) | 11.91 (10.65-13.22) | 12.91 (12.44-13.49) | 9.45 (8.46-10.44) | 16.12 (15.15-16.94) | 11.64 (10.29-13.04) | 14.02 (13.37-14.63) | 10.21 (9.09-11.29) | 15.98 (14.58-17.47) | 11.53 (9.99-12.99) | 14.37 (13.33-15.54) | 10.46 (9.16-11.80) |
| Federated States of Micronesia | 12.64 (11.69-13.56) | 9.39 (8.27-10.49) | 11.23 (10.67-12.11) | 8.45 (7.58-9.41) | 12.76 (11.43-14.06) | 9.42 (8.17-10.72) | 9.14 (8.10-10.13) | 12.75 (11.38-14.23) | 9.42 (8.06-10.85) | 12.04 (10.95-13.00) | 9.00 (7.87-10.11) | |
| Fiji | 13.45 (11.49-15.47) | 9.85 (8.21-11.54) | 11.66 (10.65-12.89) | 8.62 (7.53-9.83) | 13.13 (12.17-14.29) | 9.52 (8.37-10.74) | 11.20 (10.47-12.23) | 8.24 (7.26-9.23) | 13.47 (11.50-15.69) | 9.82 (8.23-11.61) | 11.68 (10.19-13.64) | 8.66 (7.25-10.35) |
| Guam | 17.14 (16.61-17.86) | 12.76 (11.46-13.97) | 14.25 (13.65-14.71) | 10.87 (9.84-11.79) | 17.31 (16.78-17.86) | 12.80 (11.54-14.04) | 14.27 (13.59-14.70) | 10.78 (9.75-11.72) | 17.05 (16.15-18.10) | 12.54 (11.21-13.92) | 13.84 (12.96-14.80) | 10.41 (9.31-11.48) |
| Kiribati | 11.64 (11.08-12.19) | 8.48 (7.53-9.40) | 9.98 (9.56-10.37) | 7.43 (6.67-8.17) | 12.46 (11.70-13.23) | 9.03 (7.98-10.11) | 10.19 (9.74-10.71) | 7.48 (6.64-8.27) | 12.82 (11.82-13.94) | 9.30 (8.09-10.54) | 10.57 (9.87-11.34) | 7.77 (6.83-8.72) |
| Marshall Islands | 13.63 (13.11-14.14) | 9.82 (8.76-10.81) | 11.30 (10.95-11.79) | 8.31 (7.48-9.12) | 12.03 (11.25-12.89) | 8.65 (7.56-9.71) | 11.28 (10.73-11.95) | 8.21 (7.31-9.09) | 12.74 (11.41-14.15) | 9.15 (7.80-10.49) | 11.80 (10.93-12.79) | 8.57 (7.50-9.72) |
| Northern Mariana Islands | 16.02 (14.53-17.64) | 11.88 (10.32-13.46) | 15.25 (14.18-16.44) | 11.44 (10.16-12.73) | 17.57 (16.22-18.95) | 12.96 (11.34-14.42) | 16.23 (15.29-17.11) | 12.10 (10.78-13.31) | 17.39 (15.85-19.01) | 12.82 (11.17-14.47) | 15.83 (14.79-16.83) | 11.78 (10.52-13.01) |
| Papua New Guinea | 10.42 (9.66-11.25) | 7.67 (6.73-8.53) | 9.85 (9.29-10.61) | 7.38 (6.59-8.20) | 10.53 (9.67-11.44) | 7.71 (6.72-8.70) | 10.39 (9.69-11.28) | 7.75 (6.86-8.70) | 11.28 (10.33-12.24) | 8.23 (7.15-9.27) | 11.06 (10.33-12.01) | 8.23 (7.31-9.19) |
| Samoa | 15.24 (14.27-16.22) | 11.28 (10.09-12.57) | 12.69 (11.89-13.41) | 9.52 (8.44-10.48) | 15.81 (14.88-16.91) | 11.63 (10.30-13.03) | 13.77 (12.99-14.79) | 10.30 (9.14-11.42) | 15.90 (14.75-17.11) | 11.69 (10.38-13.14) | 14.12 (13.40-15.41) | 10.56 (9.48-11.80) |
| Solomon Islands | 10.92 (10.08-11.94) | 8.07 (7.13-9.17) | 10.58 (10.00-11.27) | 7.98 (7.24-8.80) | 10.86 (9.92-11.88) | 7.96 (6.91-9.01) | 11.01 (10.37-11.63) | 8.25 (7.38-9.11) | 11.46 (10.32-12.63) | 8.37 (7.28-9.47) | 11.54 (10.65-12.35) | 8.62 (7.64-9.58) |
| Tonga | 14.22 (13.03-15.53) | 10.35 (9.05-11.70) | 12.64 (11.67-13.59) | 9.53 (8.42-10.67) | 15.32 (14.40-16.39) | 11.13 (9.84-12.46) | 13.09 (12.29-13.88) | 9.79 (8.74-10.85) | 15.71 (14.39-16.99) | 11.41 (9.94-12.84) | 13.34 (12.52-14.34) | 9.94 (8.88-11.04) |
| Vanuatu | 11.74 (10.88-12.67) | 8.66 (7.59-9.81) | 10.89 (10.39-11.41) | 8.21 (7.39-9.03) | 11.80 (10.72-12.79) | 8.69 (7.56-9.81) | 11.19 (10.57-11.88) | 8.45 (7.62-9.32) | 12.30 (11.03-13.39) | 9.02 (7.80-10.23) | 11.65 (10.89-12.46) | 8.76 (7.81-9.74) |

(Table 3 continues on next page)

| | 1990, at age 65 years | | | | 2006, at age 65 years | | | | 2016, at age 65 years | | | |
|-------------------------------------|-------------------------------|-------------------------------|-------------------------------|------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| North Africa and Middle East | 16.10 (15.87–16.32) | 11.38 (10.13–12.52) | 14.38 (14.15–14.64) | 10.49 (9.38–11.50) | 16.75 (16.42–17.03) | 11.92 (10.59–13.15) | 15.12 (14.79–15.41) | 11.13 (10.01–12.13) | 17.73 (17.30–18.17) | 12.64 (11.24–13.95) | 15.68 (15.25–16.08) | 11.55 (10.38–12.65) |
| Afghanistan | 10.00 (9.46–10.63) | 6.86 (5.96–7.76) | 10.76 (10.11–11.63) | 7.65 (6.62–8.68) | 9.82 (9.30–10.39) | 6.79 (5.91–7.66) | 10.29 (9.73–11.04) | 7.38 (6.51–8.28) | 10.74 (10.16–11.32) | 7.43 (6.48–8.34) | 10.88 (10.37–11.49) | 7.79 (6.90–8.67) |
| Algeria | 16.89 (16.13–17.53) | 11.95 (10.57–13.27) | 15.33 (14.80–15.99) | 11.17 (10.00–12.32) | 18.22 (17.40–18.59) | 12.98 (11.48–14.36) | 17.13 (16.56–17.81) | 12.58 (11.24–13.85) | 18.60 (18.10–18.98) | 13.26 (11.78–14.59) | 17.83 (17.03–18.63) | 13.10 (11.67–14.42) |
| Bahrain | 13.09 (12.02–14.15) | 9.13 (7.90–10.42) | 11.68 (10.77–12.67) | 8.43 (7.39–9.58) | 14.63 (13.63–15.74) | 10.10 (8.72–11.43) | 13.74 (12.77–14.84) | 9.84 (8.58–11.14) | 16.61 (15.16–18.32) | 11.50 (9.81–13.22) | 15.48 (13.90–17.23) | 11.17 (9.46–12.88) |
| Egypt | 15.50 (15.07–15.93) | 11.06 (9.86–12.20) | 12.95 (12.55–13.39) | 9.54 (8.51–10.44) | 16.19 (15.69–16.69) | 11.47 (10.14–12.78) | 13.56 (13.10–14.01) | 9.97 (8.91–10.95) | 16.72 (15.54–18.03) | 11.92 (10.41–13.43) | 13.96 (12.98–15.12) | 10.31 (9.14–11.49) |
| Iran | 17.07 (16.17–18.28) | 12.17 (10.65–13.67) | 14.78 (13.74–16.12) | 10.85 (9.56–12.25) | 16.26 (14.70–17.72) | 11.68 (10.13–13.19) | 15.30 (13.83–16.63) | 11.37 (9.85–12.86) | 18.06 (16.88–19.65) | 12.95 (11.35–14.60) | 16.33 (14.88–17.75) | 12.06 (10.57–13.45) |
| Iraq | 13.72 (12.79–14.82) | 9.56 (8.35–10.78) | 13.49 (12.09–14.54) | 9.65 (8.27–10.90) | 13.76 (12.77–14.59) | 9.65 (8.39–10.83) | 12.55 (11.48–13.61) | 9.06 (7.88–10.27) | 14.88 (13.44–16.39) | 10.43 (8.94–11.94) | 13.18 (12.10–15.66) | 9.47 (8.25–11.13) |
| Jordan | 15.78 (14.71–17.27) | 11.01 (9.60–12.52) | 15.55 (14.57–16.18) | 11.19 (9.90–12.38) | 15.72 (14.19–17.55) | 11.12 (9.45–12.80) | 15.81 (14.23–17.25) | 11.54 (9.99–13.12) | 18.37 (16.45–20.42) | 13.04 (11.13–14.79) | 16.92 (14.71–19.20) | 12.39 (10.66–14.42) |
| Kuwait | 16.73 (15.39–18.20) | 12.00 (10.35–13.56) | 16.73 (15.43–18.06) | 12.32 (10.88–13.78) | 15.91 (14.87–16.92) | 11.36 (9.91–12.70) | 18.03 (16.86–19.13) | 13.21 (11.72–14.69) | 17.30 (14.97–19.97) | 12.47 (10.50–14.68) | 19.83 (17.44–22.40) | 14.59 (12.52–16.91) |
| Lebanon | 15.94 (14.78–17.39) | 11.49 (9.98–12.96) | 13.80 (12.96–15.50) | 10.21 (8.93–11.92) | 18.57 (18.21–19.43) | 13.42 (11.99–14.84) | 18.19 (17.02–19.31) | 13.50 (11.95–14.99) | 20.08 (18.94–21.24) | 14.52 (12.82–16.24) | 18.59 (17.30–19.72) | 13.74 (12.16–15.26) |
| Libya | 17.46 (16.60–17.98) | 12.39 (10.97–13.74) | 15.95 (15.37–16.74) | 11.67 (10.46–12.91) | 17.84 (16.53–18.44) | 12.67 (11.04–14.13) | 16.49 (15.92–17.19) | 12.07 (10.79–13.28) | 18.09 (17.32–18.66) | 12.86 (11.32–14.24) | 15.41 (14.67–16.77) | 11.32 (9.98–12.71) |
| Morocco | 15.46 (14.74–16.09) | 11.00 (9.75–12.24) | 14.26 (13.88–14.63) | 10.45 (9.37–11.45) | 16.61 (15.64–17.43) | 11.88 (10.50–13.23) | 15.77 (15.40–16.10) | 11.61 (10.40–12.67) | 17.53 (16.58–18.42) | 12.54 (10.96–14.05) | 16.50 (16.03–16.94) | 12.16 (10.91–13.32) |
| Palestine | 15.00 (13.73–16.28) | 10.89 (9.51–12.34) | 14.34 (13.52–15.41) | 10.64 (9.47–11.82) | 14.74 (14.53–14.93) | 10.71 (9.56–11.73) | 12.89 (12.77–13.01) | 9.58 (8.66–10.41) | 13.60 (13.20–14.08) | 9.83 (8.77–10.82) | 12.89 (12.58–13.26) | 9.51 (8.54–10.36) |
| Oman | 16.33 (15.44–17.07) | 11.56 (10.15–12.91) | 15.16 (14.75–15.63) | 10.98 (9.77–12.09) | 17.98 (17.43–18.72) | 12.83 (11.35–14.25) | 15.39 (14.77–16.00) | 11.27 (10.08–12.40) | 18.96 (18.60–19.48) | 13.60 (12.06–14.99) | 16.04 (15.69–16.52) | 11.78 (10.58–12.93) |
| Qatar | 16.65 (15.53–18.29) | 11.67 (10.08–13.46) | 16.98 (15.63–18.37) | 12.19 (10.65–13.70) | 18.50 (17.66–19.71) | 12.87 (11.25–14.40) | 16.80 (14.98–18.63) | 12.09 (10.20–13.90) | 20.26 (18.41–22.92) | 14.26 (12.18–16.59) | 18.65 (16.07–21.28) | 13.53 (11.23–15.85) |
| Saudi Arabia | 16.00 (15.23–16.82) | 11.33 (10.04–12.68) | 15.74 (15.05–16.52) | 11.54 (10.23–12.80) | 14.17 (13.81–14.55) | 10.19 (9.07–11.21) | 15.17 (14.82–15.53) | 11.28 (10.17–12.29) | 16.76 (16.12–17.49) | 12.11 (10.82–13.42) | 16.20 (15.60–16.83) | 12.09 (10.88–13.25) |
| Sudan | 13.16 (12.63–13.67) | 9.33 (8.29–10.38) | 13.11 (12.83–13.41) | 9.53 (8.51–10.50) | 14.42 (13.86–15.03) | 10.28 (9.12–11.33) | 13.82 (13.50–14.28) | 10.13 (9.06–11.08) | 15.94 (15.13–16.53) | 11.37 (10.07–12.58) | 14.66 (14.23–15.09) | 10.76 (9.61–11.79) |
| Syria | 16.36 (15.64–17.09) | 11.73 (10.38–13.00) | 15.01 (14.70–15.35) | 11.12 (10.03–12.15) | 17.39 (16.84–17.82) | 12.62 (11.25–13.85) | 15.80 (15.31–16.25) | 11.83 (10.64–12.92) | 18.11 (17.61–18.87) | 13.16 (11.79–14.53) | 14.98 (14.34–15.88) | 11.26 (10.11–12.48) |
| Tunisia | 17.09 (16.46–17.78) | 12.33 (10.90–13.71) | 14.63 (13.97–15.32) | 10.82 (9.65–11.91) | 18.77 (18.22–20.04) | 13.60 (12.08–15.20) | 16.29 (15.52–17.38) | 12.03 (10.72–13.37) | 19.38 (18.34–21.04) | 14.09 (12.37–16.01) | 16.07 (14.63–17.96) | 11.98 (10.39–13.69) |

(Table 3 continues on next page)

| | 1990, at age 65 years | | | | 2006, at age 65 years | | | | 2016, at age 65 years | | | |
|--------------------------------|--------------------------------------|------------------------------------|--------------------------------------|------------------------------------|--------------------------------------|-------------------------------------|--------------------------------------|------------------------------------|--------------------------------------|-------------------------------------|--------------------------------------|-------------------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Turkey | 18.63 (18.02–19.22) | 13.03 (11.49–14.46) | 15.76 (15.30–16.22) | 11.42 (10.19–12.57) | 20.50 (19.87–21.15) | 14.55 (12.80–16.16) | 16.83 (16.41–17.34) | 12.40 (11.17–13.55) | 21.09 (19.63–22.58) | 15.03 (13.16–16.95) | 17.32 (15.87–18.72) | 12.82 (11.25–14.43) |
| United Arab Emirates | 15.95 (14.69–17.58) | 11.24 (9.71–12.78) | 15.28 (13.37–17.00) | 11.05 (9.25–12.83) | 18.38 (17.63–18.78) | 13.05 (11.59–14.44) | 16.29 (15.71–16.82) | 11.84 (10.58–13.03) | 18.38 (17.08–19.44) | 13.15 (11.56–14.77) | 16.00 (14.78–17.04) | 11.73 (10.23–13.06) |
| Yemen | 12.62 (11.92–13.47) | 8.60 (7.42–9.84) | 13.08 (12.62–13.58) | 9.23 (8.14–10.29) | 13.36 (12.97–13.99) | 9.18 (8.00–10.29) | 13.98 (13.56–14.42) | 9.96 (8.80–11.02) | 14.13 (13.58–14.54) | 9.78 (8.54–10.85) | 14.54 (13.60–15.37) | 10.45 (9.18–11.66) |
| South Asia | 12.72 (12.50–12.93) | 9.03 (8.05–9.92) | 12.13 (11.95–12.32) | 8.79 (7.91–9.59) | 14.25 (14.04–14.46) | 10.15 (9.10–11.11) | 13.15 (12.97–13.33) | 9.55 (8.65–10.39) | 15.20 (14.95–15.46) | 10.84 (9.72–11.89) | 13.80 (13.58–14.01) | 10.03 (9.06–10.87) |
| Bangladesh | 12.85 (12.14–13.57) | 9.29 (8.24–10.32) | 12.78 (12.25–13.38) | 9.44 (8.44–10.39) | 16.29 (15.70–16.71) | 11.78 (10.52–12.93) | 14.43 (14.00–14.92) | 10.73 (9.70–11.65) | 17.59 (16.60–18.34) | 12.76 (11.42–14.11) | 15.35 (14.45–16.11) | 11.44 (10.25–12.56) |
| Bhutan | 13.21 (11.99–14.44) | 9.44 (8.18–10.72) | 14.26 (13.14–15.32) | 10.39 (9.15–11.56) | 17.01 (16.06–17.93) | 12.20 (10.74–13.53) | 16.00 (15.06–16.91) | 11.74 (10.46–13.07) | 18.12 (16.85–19.04) | 13.01 (11.50–14.54) | 16.25 (15.21–17.40) | 11.92 (10.56–13.23) |
| India | 12.57 (12.31–12.83) | 8.88 (7.89–9.77) | 11.82 (11.62–12.02) | 8.52 (7.68–9.30) | 14.11 (13.88–14.36) | 10.02 (8.97–10.99) | 12.97 (12.76–13.17) | 9.36 (8.48–10.19) | 15.03 (14.75–15.31) | 10.68 (9.54–11.73) | 13.59 (13.37–13.82) | 9.82 (8.87–10.67) |
| Nepal | 12.41 (11.89–12.95) | 9.01 (8.08–9.96) | 12.65 (12.25–13.08) | 9.23 (8.33–10.10) | 14.02 (13.54–14.63) | 10.20 (9.12–11.26) | 13.94 (13.53–14.31) | 10.26 (9.27–11.25) | 15.17 (14.78–15.68) | 11.04 (9.94–12.08) | 14.74 (14.00–15.59) | 10.89 (9.75–12.07) |
| Pakistan | 14.20 (13.66–14.72) | 10.32 (9.18–11.37) | 14.12 (13.60–14.64) | 10.42 (9.35–11.46) | 13.81 (13.02–14.71) | 10.02 (8.89–11.22) | 13.36 (12.92–13.88) | 9.86 (8.89–10.81) | 14.78 (13.59–16.30) | 10.73 (9.35–12.19) | 14.06 (13.20–14.86) | 10.45 (9.31–11.57) |
| Sub-Saharan Africa | 12.74 (12.55–12.91) | 9.47 (8.54–10.30) | 12.58 (12.37–12.79) | 9.28 (8.40–10.14) | 13.08 (12.84–13.32) | 9.77 (8.86–10.62) | 12.68 (12.50–12.86) | 9.41 (8.49–10.24) | 14.39 (14.11–14.69) | 10.82 (9.81–11.76) | 13.67 (13.39–13.96) | 10.19 (9.19–11.10) |
| Southern sub-Saharan Africa | 17.26 (16.87–17.77) | 12.84 (11.61–13.96) | 14.86 (14.48–15.50) | 11.01 (9.97–12.04) | 14.05 (13.62–14.53) | 10.38 (9.39–11.41) | 11.54 (11.35–11.75) | 8.46 (7.63–9.24) | 16.73 (16.29–17.22) | 12.38 (11.16–13.54) | 13.40 (13.08–13.78) | 9.82 (8.86–10.75) |
| Botswana | 14.16 (13.28–15.74) | 10.64 (9.49–12.02) | 12.46 (11.16–14.04) | 9.30 (8.03–10.69) | 13.22 (9.32–16.70) | 9.97 (6.91–12.50) | 10.59 (8.92–12.66) | 7.74 (6.34–9.56) | 15.80 (12.73–22.97) | 11.93 (9.25–16.98) | 12.81 (11.28–15.95) | 9.42 (7.95–11.65) |
| Lesotho | 13.46 (12.52–14.91) | 10.05 (8.86–11.33) | 11.00 (10.18–12.26) | 8.22 (7.27–9.43) | 11.03 (9.65–12.89) | 8.14 (6.84–9.65) | 8.55 (8.05–9.14) | 6.24 (5.52–6.91) | 12.12 (10.11–15.73) | 8.99 (7.28–11.33) | 9.41 (8.57–10.38) | 6.87 (6.00–7.78) |
| Namibia | 13.16 (12.77–13.76) | 9.91 (8.96–10.81) | 11.31 (10.78–12.16) | 8.46 (7.58–9.42) | 13.82 (11.53–16.20) | 10.43 (8.36–12.57) | 10.66 (9.92–11.95) | 7.90 (6.93–8.97) | 16.82 (14.39–21.51) | 12.83 (10.52–16.49) | 12.69 (11.71–14.15) | 9.48 (8.32–10.77) |
| South Africa | 18.40 (18.01–18.79) | 13.63 (12.33–14.85) | 15.44 (15.09–15.77) | 11.40 (10.33–12.40) | 14.77 (14.58–14.97) | 10.89 (9.86–11.85) | 12.30 (12.15–12.47) | 8.98 (8.13–9.81) | 17.65 (17.26–18.05) | 13.01 (11.70–14.24) | 14.15 (13.83–14.47) | 10.30 (9.29–11.29) |
| Swaziland | 13.21 (12.25–14.33) | 9.88 (8.84–11.02) | 11.47 (10.53–12.85) | 8.51 (7.48–9.70) | 11.34 (8.96–12.71) | 8.32 (6.53–9.66) | 9.12 (8.24–10.46) | 6.51 (5.58–7.59) | 14.69 (11.75–19.10) | 10.94 (8.33–14.11) | 11.06 (9.92–13.44) | 8.00 (6.83–9.62) |
| Zimbabwe | 14.08 (13.05–16.47) | 10.73 (9.46–12.72) | 14.17 (12.92–18.34) | 10.74 (9.24–13.75) | 11.44 (9.75–13.85) | 8.70 (7.21–10.70) | 9.68 (9.09–10.45) | 7.25 (6.43–8.12) | 13.23 (11.75–15.66) | 10.08 (8.79–11.82) | 11.58 (10.54–13.13) | 8.77 (7.63–10.19) |
| Western sub-Saharan Africa | 13.26 (12.87–13.57) | 9.75 (8.69–10.69) | 13.41 (12.98–13.82) | 9.81 (8.75–10.80) | 13.96 (13.52–14.50) | 10.35 (9.33–11.37) | 13.66 (13.28–14.12) | 10.09 (9.09–11.09) | 14.99 (14.49–15.59) | 11.19 (10.09–12.24) | 14.55 (14.02–15.07) | 10.82 (9.73–11.84) |
| Benin | 13.82 (13.26–14.30) | 9.99 (8.81–11.00) | 12.69 (12.26–13.18) | 9.14 (8.10–10.11) | 14.25 (13.38–15.11) | 10.46 (9.29–11.58) | 12.69 (12.42–13.14) | 9.30 (8.36–10.19) | 14.29 (13.38–15.41) | 10.59 (9.32–11.79) | 13.34 (12.96–13.76) | 9.84 (8.79–10.78) |

(Table 3 continues on next page)

| | 1990, at age 65 years | | | | 2006, at age 65 years | | | | 2016, at age 65 years | | | |
|--------------------------------|------------------------|------------------------|------------------------|-----------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Burkina Faso | 12.40 (11.91–13.17) | 8.98 (7.91–10.05) | 12.23 (11.74–13.15) | 8.77 (7.71–9.88) | 12.97 (12.44–13.46) | 9.62 (8.60–10.58) | 12.76 (12.43–13.05) | 9.39 (8.42–10.30) | 13.30 (12.88–13.77) | 9.97 (9.00–10.91) | 13.17 (12.89–13.50) | 9.83 (8.88–10.73) |
| Cameroon | 13.39 (12.57–14.32) | 9.84 (8.63–10.89) | 13.20 (12.62–14.04) | 9.66 (8.58–10.75) | 13.02 (11.61–14.67) | 9.67 (8.30–11.14) | 12.33 (11.78–12.90) | 9.08 (8.12–9.98) | 14.03 (12.41–16.13) | 10.55 (8.94–12.39) | 12.92 (12.13–13.78) | 9.61 (8.58–10.65) |
| Cape Verde | 16.82 (16.36–17.50) | 12.57 (11.34–13.76) | 13.64 (13.30–14.20) | 10.22 (9.26–11.18) | 18.52 (17.54–19.66) | 13.89 (12.35–15.40) | 13.37 (12.54–15.40) | 10.05 (8.97–11.50) | 18.81 (18.11–20.06) | 14.20 (12.74–15.70) | 13.99 (13.16–15.03) | 10.55 (9.38–11.70) |
| Chad | 13.10 (12.56–13.70) | 9.58 (8.49–10.56) | 12.80 (12.43–13.23) | 9.33 (8.30–10.25) | 13.59 (12.63–14.50) | 9.91 (8.71–11.12) | 12.52 (11.98–13.13) | 9.12 (8.10–10.07) | 14.31 (13.21–15.24) | 10.53 (9.28–11.82) | 13.14 (12.60–13.96) | 9.64 (8.60–10.67) |
| Côte d'Ivoire | 12.50 (11.87–13.22) | 8.95 (7.86–10.06) | 11.99 (11.62–12.35) | 8.59 (7.60–9.49) | 12.16 (11.28–13.45) | 9.02 (7.95–10.20) | 11.80 (11.06–12.26) | 8.66 (7.65–9.57) | 12.99 (12.23–14.00) | 9.68 (8.59–10.82) | 12.58 (11.69–13.06) | 9.29 (8.14–10.21) |
| The Gambia | 13.49 (12.43–14.56) | 9.92 (8.68–11.15) | 13.98 (13.17–14.71) | 10.38 (9.17–11.49) | 14.45 (13.37–15.71) | 10.63 (9.35–12.04) | 13.74 (13.29–14.36) | 10.19 (9.10–11.18) | 14.90 (14.03–16.33) | 11.07 (9.80–12.57) | 14.50 (13.75–15.08) | 10.81 (9.61–11.81) |
| Ghana | 12.72 (11.81–13.84) | 9.50 (8.43–10.65) | 13.03 (12.47–13.83) | 9.68 (8.64–10.76) | 13.55 (12.53–15.05) | 10.28 (9.00–11.72) | 12.93 (12.69–13.35) | 9.70 (8.75–10.56) | 14.31 (13.63–15.75) | 10.88 (9.79–12.28) | 13.93 (13.19–14.59) | 10.44 (9.34–11.45) |
| Guinea | 12.55 (11.79–13.20) | 9.24 (8.22–10.27) | 13.69 (13.08–14.38) | 10.07 (8.99–11.13) | 12.77 (11.85–13.85) | 9.53 (8.39–10.57) | 12.78 (12.21–13.55) | 9.54 (8.47–10.54) | 13.00 (12.02–14.44) | 9.77 (8.61–11.03) | 13.10 (12.52–13.93) | 9.80 (8.72–10.84) |
| Guinea-Bissau | 11.17 (10.40–12.25) | 8.28 (7.27–9.32) | 10.81 (10.12–11.46) | 7.95 (7.04–8.81) | 11.67 (10.81–13.00) | 8.68 (7.64–9.85) | 10.83 (10.08–11.85) | 8.02 (7.10–9.02) | 12.43 (11.57–13.29) | 9.32 (8.21–10.33) | 11.07 (10.36–12.17) | 8.24 (7.28–9.25) |
| Liberia | 12.87 (12.19–13.55) | 9.19 (8.12–10.24) | 13.32 (12.76–13.88) | 9.44 (8.35–10.51) | 12.85 (11.97–14.00) | 9.26 (8.08–10.42) | 13.39 (12.82–14.01) | 9.62 (8.49–10.70) | 13.39 (12.92–14.05) | 9.80 (8.69–10.83) | 13.87 (13.47–14.35) | 10.11 (9.03–11.10) |
| Mali | 11.41 (10.78–12.08) | 8.45 (7.48–9.39) | 12.64 (12.18–13.24) | 9.22 (8.21–10.21) | 13.50 (12.59–14.36) | 10.03 (8.88–11.14) | 14.23 (13.49–14.98) | 10.45 (9.31–11.53) | 14.28 (12.68–15.69) | 10.70 (9.24–12.11) | 14.69 (13.35–16.01) | 10.91 (9.54–12.34) |
| Mauritania | 12.41 (11.70–13.18) | 9.15 (8.13–10.19) | 13.14 (12.66–13.85) | 9.59 (8.50–10.66) | 14.11 (13.03–15.46) | 10.45 (9.16–11.71) | 15.56 (14.62–16.62) | 11.41 (10.07–12.76) | 15.12 (13.82–17.09) | 11.31 (9.75–13.01) | 16.12 (14.81–17.86) | 11.90 (10.35–13.61) |
| Niger | 13.13 (12.55–13.58) | 9.80 (8.78–10.74) | 12.73 (12.10–13.48) | 9.45 (8.40–10.48) | 13.86 (12.92–14.49) | 10.38 (9.23–11.42) | 13.34 (12.71–14.10) | 9.97 (8.96–10.96) | 14.14 (12.56–15.46) | 10.68 (9.22–12.03) | 13.47 (12.69–14.49) | 10.11 (9.00–11.35) |
| Nigeria | 13.88 (12.99–14.54) | 10.20 (9.05–11.27) | 14.32 (13.34–15.44) | 10.51 (9.27–11.90) | 15.00 (14.02–16.49) | 11.09 (9.84–12.49) | 14.95 (13.99–16.27) | 11.05 (9.81–12.48) | 16.78 (15.59–18.56) | 12.49 (10.97–14.15) | 16.32 (14.96–17.83) | 12.14 (10.60–13.64) |
| São Tomé and Príncipe | 14.67 (13.71–15.41) | 11.04 (9.88–12.18) | 14.05 (13.33–14.80) | 10.41 (9.27–11.53) | 13.91 (13.52–14.37) | 10.51 (9.54–11.41) | 13.97 (13.51–14.51) | 10.38 (9.32–11.34) | 15.21 (14.52–16.23) | 11.55 (10.38–12.75) | 14.59 (13.22–15.99) | 10.89 (9.37–12.43) |
| Senegal | 13.25 (12.57–13.91) | 9.89 (8.85–10.95) | 12.63 (12.28–13.06) | 9.37 (8.38–10.26) | 13.28 (12.72–14.02) | 9.96 (8.93–11.01) | 13.02 (12.82–13.40) | 9.69 (8.68–10.58) | 13.86 (13.33–14.26) | 10.42 (9.40–11.37) | 13.62 (13.00–14.31) | 10.17 (9.08–11.22) |
| Sierra Leone | 13.64 (13.11–14.12) | 10.11 (9.03–11.09) | 12.33 (11.78–12.96) | 9.01 (7.96–9.99) | 12.39 (11.59–13.22) | 9.17 (8.13–10.23) | 12.03 (11.70–12.50) | 8.79 (7.85–9.68) | 12.93 (11.99–14.20) | 9.70 (8.58–10.91) | 12.86 (12.42–13.49) | 9.52 (8.46–10.54) |
| Togo | 13.00 (12.26–13.76) | 9.63 (8.59–10.69) | 12.81 (12.34–13.30) | 9.47 (8.41–10.42) | 13.60 (12.35–15.28) | 10.15 (8.83–11.60) | 12.22 (11.58–12.65) | 9.10 (8.16–10.03) | 13.81 (13.12–15.13) | 10.44 (9.27–11.72) | 12.82 (11.89–13.45) | 9.64 (8.58–10.61) |
| Eastern sub-Saharan Africa | 11.12 (10.88–11.37) | 8.41 (7.63–9.10) | 11.42 (11.16–11.72) | 8.54 (7.69–9.31) | 12.35 (11.95–12.78) | 9.37 (8.48–10.18) | 12.29 (12.03–12.52) | 9.23 (8.34–10.02) | 13.67 (13.20–14.20) | 10.43 (9.47–11.34) | 13.18 (12.71–13.64) | 9.94 (8.96–10.81) |

(Table 3 continues on next page)

| | 1990, at age 65 years | | | | 2006, at age 65 years | | | | 2016, at age 65 years | | | |
|--------------------------------|------------------------|------------------------|------------------------|-----------------------|------------------------|-----------------------|------------------------|-----------------------|------------------------|------------------------|------------------------|-----------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Burundi | 9.43 (8.73–10.38) | 7.25 (6.51–8.09) | 9.51 (8.67–10.67) | 7.21 (6.37–8.26) | 11.02 (10.25–11.74) | 8.49 (7.62–9.38) | 11.46 (10.48–12.32) | 8.73 (7.60–9.77) | 12.09 (11.16–13.02) | 9.39 (8.35–10.40) | 12.29 (11.00–13.34) | 9.42 (8.29–10.56) |
| Comoros | 12.25 (11.44–13.47) | 9.26 (8.20–10.46) | 12.63 (12.08–13.35) | 9.35 (8.37–10.29) | 13.82 (12.96–14.96) | 10.53 (9.39–11.83) | 13.71 (13.27–14.51) | 10.31 (9.29–11.38) | 14.31 (13.57–15.90) | 10.96 (9.83–12.39) | 14.28 (13.44–15.15) | 10.79 (9.64–11.90) |
| Djibouti | 14.87 (14.23–15.29) | 11.27 (10.14–12.31) | 12.89 (12.58–13.36) | 9.75 (8.81–10.66) | 14.72 (12.98–16.44) | 11.21 (9.45–12.88) | 13.05 (12.38–13.73) | 9.86 (8.78–10.82) | 15.34 (13.92–17.48) | 11.73 (10.11–13.71) | 13.91 (12.86–15.01) | 10.52 (9.34–11.78) |
| Eritrea | 10.29 (9.80–10.84) | 7.80 (7.00–8.61) | 10.64 (9.97–11.34) | 7.99 (7.13–8.88) | 12.03 (11.31–12.92) | 9.19 (8.18–10.21) | 12.61 (11.58–13.31) | 9.50 (8.41–10.50) | 12.49 (11.73–13.45) | 9.61 (8.63–10.69) | 12.75 (11.68–13.73) | 9.67 (8.49–10.76) |
| Ethiopia | 9.54 (9.06–10.09) | 7.25 (6.56–7.97) | 9.78 (9.24–10.50) | 7.35 (6.56–8.14) | 11.41 (10.61–12.18) | 8.73 (7.79–9.69) | 12.47 (11.67–12.84) | 9.43 (8.46–10.31) | 13.26 (12.20–14.32) | 10.18 (9.02–11.38) | 13.32 (12.03–14.75) | 10.12 (8.80–11.52) |
| Kenya | 13.41 (13.09–13.74) | 10.31 (9.40–11.15) | 13.74 (13.38–14.10) | 10.44 (9.47–11.32) | 14.05 (13.65–14.44) | 10.79 (9.84–11.66) | 13.14 (12.83–13.42) | 9.97 (9.05–10.78) | 15.37 (14.93–15.85) | 11.79 (10.73–12.79) | 14.14 (13.81–14.51) | 10.69 (9.71–11.58) |
| Madagascar | 11.92 (11.31–12.61) | 9.05 (8.19–10.01) | 12.13 (11.79–12.43) | 9.11 (8.22–9.93) | 12.57 (11.72–13.83) | 9.61 (8.49–10.80) | 12.78 (12.19–13.40) | 9.66 (8.70–10.60) | 13.10 (11.71–15.54) | 10.11 (8.76–12.00) | 13.05 (11.51–14.31) | 9.93 (8.66–11.13) |
| Malawi | 12.06 (10.88–13.26) | 9.11 (7.97–10.40) | 13.32 (11.80–16.19) | 9.89 (8.35–12.03) | 12.63 (10.45–14.93) | 9.63 (7.73–11.73) | 11.51 (10.27–13.33) | 8.57 (7.39–10.06) | 13.97 (12.43–16.42) | 10.67 (9.21–12.67) | 12.90 (11.40–14.06) | 9.70 (8.43–10.99) |
| Mozambique | 12.71 (12.07–13.26) | 9.51 (8.52–10.49) | 11.86 (11.50–12.31) | 8.76 (7.87–9.60) | 14.52 (12.48–16.02) | 10.81 (8.97–12.42) | 11.94 (10.91–12.86) | 8.76 (7.56–9.86) | 14.76 (13.22–16.48) | 11.05 (9.57–12.65) | 13.01 (11.80–14.04) | 9.57 (8.27–10.72) |
| Rwanda | 10.17 (9.56–10.82) | 7.74 (6.92–8.55) | 10.69 (9.78–11.65) | 8.06 (7.08–9.05) | 14.05 (12.83–15.21) | 10.69 (9.41–11.91) | 13.03 (12.64–13.77) | 9.82 (8.87–10.78) | 15.00 (13.91–16.44) | 11.48 (10.13–12.98) | 14.27 (13.24–15.18) | 10.77 (9.57–11.89) |
| Somalia | 10.21 (9.54–10.86) | 7.76 (6.89–8.58) | 11.58 (10.94–12.19) | 8.72 (7.78–9.63) | 10.34 (9.69–11.08) | 7.87 (7.03–8.86) | 11.59 (10.64–12.32) | 8.77 (7.76–9.78) | 10.85 (10.08–11.56) | 8.29 (7.35–9.17) | 11.93 (10.81–12.75) | 9.03 (7.87–10.03) |
| South Sudan | 12.45 (11.38–13.56) | 8.77 (7.43–10.01) | 12.70 (11.77–13.88) | 8.89 (7.68–10.13) | 12.72 (11.39–14.54) | 9.21 (7.81–10.90) | 12.72 (11.88–13.89) | 9.13 (7.99–10.42) | 12.86 (11.61–14.69) | 9.50 (8.22–11.12) | 12.94 (12.02–13.95) | 9.44 (8.33–10.58) |
| Tanzania | 12.46 (11.77–13.34) | 9.43 (8.40–10.53) | 12.65 (12.07–13.54) | 9.49 (8.47–10.54) | 13.15 (11.96–15.19) | 10.01 (8.70–11.77) | 12.80 (12.36–13.65) | 9.66 (8.66–10.64) | 14.10 (13.34–15.98) | 10.80 (9.62–12.29) | 13.64 (12.43–14.45) | 10.38 (9.17–11.42) |
| Uganda | 10.95 (10.45–11.64) | 8.14 (7.30–9.03) | 11.31 (10.58–12.11) | 8.27 (7.27–9.25) | 12.01 (11.33–12.82) | 9.01 (8.03–10.01) | 11.81 (10.83–12.44) | 8.76 (7.74–9.66) | 13.61 (12.73–15.05) | 10.35 (9.21–11.69) | 12.79 (11.72–13.52) | 9.58 (8.43–10.61) |
| Zambia | 11.59 (10.67–13.24) | 8.88 (7.83–10.24) | 14.29 (13.12–15.19) | 10.74 (9.49–11.99) | 10.41 (9.17–12.23) | 7.93 (6.83–9.42) | 9.47 (8.91–10.43) | 7.12 (6.34–7.99) | 13.01 (11.11–16.03) | 9.95 (8.29–12.45) | 10.99 (9.92–12.90) | 8.27 (7.19–9.81) |
| Central sub-Saharan Africa | 11.66 (11.23–12.15) | 8.47 (7.56–9.35) | 11.32 (11.00–11.66) | 8.14 (7.24–8.98) | 11.65 (11.24–12.11) | 8.56 (7.67–9.40) | 12.10 (11.81–12.39) | 8.80 (7.89–9.69) | 12.76 (12.32–13.22) | 9.46 (8.51–10.35) | 12.88 (12.42–13.25) | 9.46 (8.45–10.39) |
| Angola | 11.07 (9.98–12.74) | 8.23 (7.17–9.58) | 10.78 (9.56–11.90) | 7.98 (6.87–9.08) | 11.57 (10.28–13.28) | 8.67 (7.46–10.05) | 12.53 (11.28–13.71) | 9.30 (8.12–10.51) | 13.33 (11.70–15.06) | 9.99 (8.59–11.47) | 13.62 (11.94–15.00) | 10.11 (8.69–11.47) |
| Central African Republic | 10.33 (9.69–11.07) | 7.60 (6.71–8.45) | 9.17 (8.60–9.84) | 6.70 (5.91–7.52) | 10.17 (8.73–12.21) | 7.56 (6.31–9.11) | 8.94 (8.24–10.36) | 6.58 (5.71–7.76) | 10.65 (9.27–12.61) | 7.97 (6.73–9.62) | 9.38 (8.61–10.77) | 6.95 (6.09–8.16) |
| Congo (Brazzaville) | 10.23 (9.47–11.12) | 7.54 (6.62–8.50) | 9.93 (9.32–10.88) | 7.27 (6.36–8.22) | 11.08 (10.24–12.10) | 8.21 (7.19–9.20) | 12.97 (12.28–13.52) | 9.54 (8.50–10.51) | 12.36 (11.03–13.94) | 9.22 (7.89–10.70) | 13.89 (12.35–15.53) | 10.32 (8.97–11.90) |

(Table 3 continues on next page)

| | 1990, at age 65 years | | | | 2006, at age 65 years | | | | 2016, at age 65 years | | | |
|----------------------------------|------------------------|----------------------|------------------------|---------------------|------------------------|-----------------------|------------------------|-----------------------|------------------------|-----------------------|------------------------|-----------------------|
| | Females | | Males | | Females | | Males | | Females | | Males | |
| | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE | Life expectancy | HALE |
| (Continued from previous page) | | | | | | | | | | | | |
| Democratic Republic of the Congo | 12.12 (11.58–12.81) | 8.73 (7.67–9.72) | 12.06 (11.75–12.46) | 8.55 (7.57–9.47) | 11.83 (11.37–12.42) | 8.65 (7.71–9.57) | 12.25 (11.94–12.49) | 8.84 (7.89–9.74) | 12.78 (12.31–13.29) | 9.45 (8.42–10.39) | 12.92 (12.42–13.33) | 9.46 (8.43–10.38) |
| Equatorial Guinea | 9.99 (9.12–11.11) | 7.42 (6.43–8.39) | 9.97 (8.98–11.29) | 7.37 (6.34–8.47) | 13.95 (11.33–16.90) | 10.37 (8.18–12.78) | 14.22 (12.47–16.55) | 10.46 (8.83–12.43) | 15.71 (13.12–19.29) | 11.71 (9.45–14.15) | 15.35 (13.29–18.00) | 11.33 (9.53–13.46) |
| Gabon | 12.61 (11.83–13.42) | 9.26 (8.22–10.27) | 10.76 (10.09–11.80) | 7.93 (7.01–8.93) | 12.21 (11.38–13.06) | 8.98 (7.94–10.04) | 13.15 (11.99–13.91) | 9.72 (8.51–10.79) | 14.23 (13.07–16.01) | 10.51 (9.24–11.89) | 14.30 (12.61–16.05) | 10.62 (9.01–12.20) |

Data in parentheses are 95% uncertainty intervals. To download the data in this table, please visit the Global Health Data Exchange. GBD=Global Burden of Disease. HALE=healthy life expectancy. SDI=Socio-demographic index.

Table 3: Global, regional, and GBD location-specific life expectancy and HALE at age 65 years, by sex, in 1990, 2006, and 2016

two countries for females (Lesotho and South Africa) and three for males (Lesotho, South Africa, and Swaziland). In 2016, HALE at age 65 years was highest in Singapore for males (15.1 years [13.4–16.8]) and Japan for females (18.7 years [17.1–20.0]), whereas it was lowest for males in Lesotho (6.9 years [6.0–7.8]) and females in Afghanistan (7.4 years [6.5–8.3]).

The leading Level 3 causes of DALYs in 2016 varied by country, but with regional commonalities, described on the basis of GBD regions¹² in figure 3. Across the 46 countries within sub-Saharan Africa, the leading Level 3 cause of DALYs for females was either HIV/AIDS (17 countries), malaria (13 countries), or diarrhoeal diseases (nine countries). Cerebrovascular disease, ischaemic heart disease, and low back and neck pain were the most common leading Level 3 causes of all-age DALYs for females in all GBD regions except for Oceania, Andean Latin America and central Latin America, and all regions of sub-Saharan Africa. Among the 34 countries in the high-income super-region, low back and neck pain was the leading cause of DALYs for females in all but three countries. Exceptions within the region were Greece and the USA, for which ischaemic heart disease was the leading cause of DALYs, and Greenland, where self-harm was the highest cause of DALYs for females (and for males). For males, these regional commonalities also occurred, but with some notable differences in the leading causes by location compared with females. Leading causes of DALYs for males in sub-Saharan Africa were broadly similar to those for females: HIV/AIDS (16 countries), malaria (11 countries), or lower respiratory infections (nine countries). Cerebrovascular disease or ischaemic heart disease was the leading cause of all-age DALYs for males in almost all GBD regions except for Andean Latin America, central Latin America, tropical Latin America, high-income Asia Pacific, and all regions of sub-Saharan Africa. Exceptions within the regions included Greenland, where self-harm was the leading cause of DALYs for males in 2016, Peru, where lower respiratory infection was the

leading cause, and Nicaragua, where chronic kidney disease was the leading cause. The leading cause of DALYs for both males and females in Syria and Iraq was conflict and terrorism, reflecting the ongoing conflict.

Epidemiological transition

The shift in the global burden of DALYs from CMNN to NCDs is a clear indicator of the epidemiological transition.²¹ The ratios of DALYs from NCDs to those from CMNN diseases in 1990, 1995, 2000, 2006, 2010, and 2016 are shown by SDI and at the global, GBD regional, and GBD super-regional scales for both males and females in figure 4. In 2016, 17 (of 21) GBD regions and six (of seven) GBD super-regions had a ratio of more than 1 for either men or women, representing the shift to more DALYs from NCDs than from CMNN diseases, compared with 13 regions and five super-regions in 1990. In 2016, females had more DALYs due to NCDs relative to CMNN causes than did males, with higher ratios in 14 regions. This figure also shows how these ratios shift with increasing SDI, with 18.5 as the 2016 ratio for high SDI for men and 20.5 for women and 0.52 for men in low SDI and 0.57 for women. Several regions showed large shifts toward more NCD DALYs than CMNN DALYs, with 12 regions doubling in ratio from 1990 to 2016 for females and 11 regions doubling for males over the same time period.

Expected age-standardised YLL rates decreased substantially with increasing SDI, most notably because of decreases in CMNN causes, such as diarrhoea, lower respiratory infections, and other common infectious diseases; NTDs and malaria; neonatal disorders; nutritional deficiencies; and HIV/AIDS and tuberculosis (figure 5A). Several causes had their largest expected age-standardised rates at intermediate levels of SDI, such as diabetes, urogenital, blood, and endocrine diseases; cardiovascular diseases; and chronic respiratory diseases. Causes that increased in expected age-standardised YLL rates with increasing SDI included neoplasms and neurological disorders. Much less variation was estimated





Figure 1: Trends of total DALYs (A) and age-standardised DALY rates (B) from 1990 to 2016 for GBD Level 1 cause groups by SDI quintile
 Shaded areas show 95% uncertainty intervals. DALYs=disability-adjusted life-years. GBD=Global Burden of Disease. SDI=Socio-demographic Index.


for expected age-standardised YLD rates with SDI than for YLL rates. Although the relationship between disability and SDI was generally constant for most Level 2 causes, nutritional deficiencies, NTDs, and malaria resulted in greater than expected levels of disability at lower SDI.

Compared with YLLs and YLDs expressed as all-age rates, the consequences of differences in population age structure become apparent for a number of causes (figure 5B). Cardiovascular diseases, in particular, result in greater YLLs at higher SDI in terms of all-age rates,

reflecting the older age of populations at higher levels of SDI. By contrast, the pattern observed when standardising for age structure—YLLs accruing at intermediate SDI—highlights health loss from diseases that typically manifest at older ages undistorted by the lower overall age structure of the populations. The expected all-age YLD rates generally increased with rising SDI, particularly for mental and substance use disorders and musculoskeletal disorders, although variation with SDI remained substantially lower than for YLLs.



(Figure 2 continues on next page)



(Figure 2 continues on next page)




Figure 2: Leading 30 Level 3 causes of total DALYs for 1990, 2006, and 2016, with percentage change in number of DALYs and all-age and age-standardised DALY rates, overall and by SDI quintile
 Overall (A). Low SDI (B). Low-middle SDI (C). Middle SDI (D). High-middle SDI (E). High SDI (F). Causes are connected by lines between time periods; solid lines are increases and dashed lines are decreases. For the time period of 1990-2006 and 2006-16, three measures of change are shown: percentage change in the number of DALYs, percentage change in the all-age DALY rate, and percentage change in the age-standardised DALY rate. Statistically significant changes are shown in bold. COPD=chronic obstructive pulmonary disease. DALYs=disability-adjusted life-years. STDs=sexually transmitted diseases.

The proportion of DALYs due to YLDs versus YLLs has shifted distinctly since 1990, and these changes are shown in figure 6. More all-age DALYs as a result of YLDs rather than YLLs indicates a reduction in premature death, with individuals living longer with disability than previously. Globally, 95.9% of countries increased in the proportion of all-age DALYs due to YLDs from 1990 to 2016. In 2016, Kuwait (0.64), Qatar (0.63), and Bahrain (0.60) had the highest ratios of DALYs due to YLDs, whereas Chad (0.14), Niger (0.14), and the Central African Republic (0.13) had the lowest. This finding is a change from 1990, when the highest three proportions were for Andorra (0.49), Kuwait (0.48), and Iceland (0.47), and the lowest three were for Burundi (0.08), Malawi (0.08), and Niger (0.06).

Observed versus expected total and cause-specific burden

As shown in figure 7, 83 countries had greater than expected age-standardised DALY rates on the basis of SDI in 2016 (a ratio of observed to expected age-standardised DALY rate of greater than 1), generally within central Europe, eastern Europe, and central Asia (20 of 29 countries) or in the super-region of sub-Saharan Africa (24 of 46 countries). Among the 34 locations in the five regions within the high-income GBD super-region, three had greater than expected DALYs relative to their SDI. The five locations with the lowest level of age-standardised DALYs relative to the level expected on the basis of SDI were Nicaragua, Costa Rica, the Maldives, Peru, and Israel. Countries within western Europe (eg, France, Norway, and Iceland), western sub-Saharan Africa (eg, The Gambia, Senegal, and Liberia), eastern sub-Saharan Africa (eg, Ethiopia, Rwanda, and Burundi), east Asia (eg, China), and a subset of countries in south and southeast Asia (eg, Bhutan and Bangladesh) all had ratios of observed to expected DALY rates of less than 1 in 2016. The five locations with the highest age-standardised DALY rates relative to the rates expected on the basis of SDI were Lesotho, Swaziland, South Africa, Fiji, and Botswana, with Lesotho's ratio of observed to expected measured at 2.43 in 2016.

Worldwide, ischaemic heart disease and stroke remained the two leading causes of DALYs, and expected levels were close to those observed at a global scale, with ratios of 0.84 for ischaemic heart disease and 1.0 for stroke (figure 8). The ratio of observed to expected DALYs ranged from 0.24 in Qatar to 3.33 in Ukraine for ischaemic heart disease and from 0.32 in Oman to 2.98 in Bulgaria for stroke. This ratio was less than 1 for 114 of 167 global locations for ischaemic heart disease and 91 of 145 for stroke. Low back and neck pain resulted in fewer than expected DALYs globally, with a ratio of 0.95 globally and 74 locations showing a ratio of less than 1. Of the ten leading causes for each location, observed levels of DALYs were frequently greater than expected on the basis of SDI for causes such as diabetes mellitus (ratio of observed

to expected DALYs of greater than 1 in 95 global locations), sense organ diseases (79 locations), and skin and subcutaneous diseases (72 locations).

Causes that resulted in greater than expected DALYs for both sexes combined showed strong regional patterns. In the GBD high-income super-region, of the leading ten causes of DALYs, COPD had greater than expected DALYs in 14 of 34 locations and less than expected in none. Observed DALYs were frequently greater than expected for low back and neck pain (27 locations), although the ratio did not exceed 1.66 for any single location. Although drug use disorders were only within the leading ten causes for one location in the high-income super-region, the ratio of observed to expected DALYs in the USA was 4.37. In addition to being the most common leading causes of DALYs, ischaemic heart disease and stroke were sources of more DALYs than expected for most locations in the central Europe, eastern Europe, and central Asia super-region. Here, the highest ratio of observed to expected DALYs was 3.33 for ischaemic heart disease and 2.98 for stroke. Mental and substance use disorders were also notable in the region for the high levels of observed DALYs relative to those expected, particularly in eastern Europe, where the ratio for the Level 4 cause of alcohol use disorders was 5.38 and ranged from 3.33 in Moldova to 5.7 in Russia. For locations in the super-region of Latin America and Caribbean, the burden from interpersonal violence was greater than expected, with a ratio of 3.37 for the super-region and ranging from 1.68 in Ecuador to 7.22 in Venezuela. Diabetes was also a source of greater than expected burden in this super-region, with DALYs exceeding expected levels in 19 of 37 locations. Additionally, diabetes contributed to a higher burden of DALYs in many locations in the super-region of southeast Asia, east Asia, and Oceania, with ratios of observed to expected levels ranging from 0.68 in the Maldives to 10.48 in Fiji.

In south Asia, neonatal encephalopathy due to birth asphyxia and trauma caused greater than expected DALYs for two countries, with ratios of 1.01 in Nepal and 2.72 in Pakistan. Ongoing conflicts in north Africa and the Middle East were reflected in higher than expected DALYs in seven of the 21 locations in the region. Patterns of observed DALYs compared with those expected in sub-Saharan Africa were dominated by the ongoing effect of HIV/AIDS, particularly in southern sub-Saharan Africa. Observed levels of DALYs exceeded those expected in 37 of 46 locations, ranging from a ratio of 8.10 in Burundi to 316.34 in Lesotho.

The five locations with the largest magnitude of improvement—a decrease in the difference between observed and expected all-cause age-standardised DALY rates from 1990 to 2016—included Greenland, the Maldives, Bermuda, Ethiopia, and Liberia. The five locations that showed the smallest decrease in the difference between observed and expected DALY rates




Figure 3: Leading Level 3 causes of all-age DALY rates by location for females (A) and males (B) in 2016

ATG=Antigua and Barbuda. DALY=disability-adjusted life-year. FSM=Federated States of Micronesia. LCA=Saint Lucia. TLS=Timor-Leste. TTO=Trinidad and Tobago. VCT=Saint Vincent and the Grenadines. VUT=Vanuatu. WSM=Samoa.




Figure 4: Ratios of all-age DALY counts from NCDs to those from CMNN diseases by SDI quintile, GBD super-region, and region, for males and females separately, for 1990, 1995, 2000, 2006, 2010, and 2016

Each point represents the ratio of DALYs from NCDs to those from CMNN diseases for each GBD region in a given year by sex. Ratios of more than 1.00 represent a greater number of DALYs from NCDs than from CMNN diseases. CMNN=communicable, maternal, neonatal, and nutritional. DALY=disability-adjusted life-year. GBD=Global Burden of Disease. NCDs=non-communicable diseases. SDI=Socio-demographic Index.

over the period 1990–2016 were Lesotho, Swaziland, Syria, South Africa, and Botswana.

Discussion
General findings

Although the world has become healthier since 1990, this progress has been highly heterogeneous. From 1990 to 2016, global HALE at birth increased. The total number of years of functional health lost (life expectancy minus HALE) increased from 1990 to 2016, indicating an absolute expansion of morbidity. YLL rates have

fallen more rapidly than YLD rates so that non-fatal health loss as a proportion of DALYs has grown from 1990 to 2016. This progress has been driven by marked reductions in DALYs due to CMNN causes and, to a lesser extent, by declines in age-standardised NCD and injury DALYs. Population growth in the past 27 years (from 5.27 billion to 7.39 billion) has resulted in a significant increase in total DALYs due to NCDs, coupled with a high total DALY burden in 2016, thus leading to an overall increasing toll on health systems.^{22,23} GBD 2016 continues to unequivocally show an absolute




Figure 5: Expected relationship between age-standardised YLL and YLD rates (per 100 000 people) and SDI (A) and all-age YLL and YLD rates (per 100 000 people) and SDI (B) for 21 Level 2 causes These stacked curves represent the average relationship between SDI and YLL and YLD rates for each cause observed across all geographies over the time period 1990–2016. In each figure, the y axis goes from lowest SDI to highest SDI. The left side shows rates for YLLs on a reflected axis and the right side shows rates for YLDs; higher rates are further from the midline in each direction. Differences in patterns between (A) and (B) is the effect of shifts in population age structure in relation to SDI. DALY=disability-adjusted life-year. SDI=Socio-demographic Index. YLDs=years lived with disability. YLLs=years of life lost.




Figure 6: Proportion of all-age DALY counts due to years lived with disability for all ages and both sexes combined for 1990 (A) and 2016 (B)
 Locations are colour-coded in terms of the magnitude of the ratio. ATG=Antigua and Barbuda. DALY=disability-adjusted life-year. FSM=Federated States of Micronesia. LCA=Saint Lucia. TLS=Timor-Leste. TTO=Trinidad and Tobago. VCT=Saint Vincent and the Grenadines.




Figure 7: Ratio of observed to expected age-standardised DALY rates (per 100 000) on the basis of SDI alone for both sexes combined in 2016

Ratios are colour-coded in terms of the magnitude of differences between observed and expected age-standardised DALY rates. Blues indicate much lower observed DALY levels than expected on the basis of SDI, whereas reds reflect observed DALYs that far exceed expected levels given SDI. ATG=Antigua and Barbuda. DALY=disability-adjusted life-year. FSM=Federated States of Micronesia. LCA=Saint Lucia. SDI=Socio-demographic Index. TLS=Timor-Leste. TTO=Trinidad and Tobago. VCT=Saint Vincent and the Grenadines.

expansion of morbidity⁴—that is, as life expectancy increases across a population, the absolute amount of functional health loss increases—but GBD 2016 also shows a relative compression of morbidity—that is, the proportion of total life spent ill per person decreases with increasing SDI.²⁴ Substantial opportunities remain for innovation in human resources for health, development assistance for health, and targeted health policies to alleviate causes of old-age disability and transform the experience of later life for billions of people.²⁵

From 1990 to 2016, life expectancy, HALE, and years lived with functional health loss all increased as populations moved through the SDI quintiles. These quintiles reflect in a geographical cross section of the world what is seen through time during the epidemiological transition. The rapid decrease in burden from CMNN diseases from low to high SDI quintile was most substantial. NCDs and injuries were also seen to decline, but at much more modest rates. Declines in fertility, increases in education, and increases in income per person (the triad of inputs into the SDI composite), were tightly coupled with population-based HALE outcomes.²⁶ This finding suggests that the wide agendas of these three social determinants of health have roles within discussions of overall health system strengthening.

Further dissection of the results clearly shows that a substantial impact has been made on the global burden of YLLs. A much more modest effect, however, has been made on the burden of large causes of YLDs at all levels of SDI. The contrast in this ratio is marked and illustrates another striking view of the epidemiological transition when looking at the proportion of DALYs due to YLDs globally. Although continued progress on reduction of YLLs prematurely is important, how to ameliorate the primary causes and underlying risks of YLDs globally should be reflected on (low back and neck pain, sense organ diseases, and skin diseases) to decrease the time that ageing populations live with functional health loss across the SDI spectrum. This persisting burden of morbidity highlights the importance, both in terms of relief of suffering and prevention of health-care costs, of placement of emphasis on prevention of disabling illness and mitigation of the ill effects of the disabling conditions that do occur.

Cross-cutting themes

The concept of the grand convergence—that levels of under-5 and maternal mortality, as well as selected infectious diseases, would converge across SDI quintiles within a generation—was promoted in the *Lancet* Commission on investing in health²⁷ and discussed in the

Global Health Metrics

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|----------------------------------|------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| Global | IHD (0.84) | Stroke (1.0) | LRI (1.67) | Back+Neck (0.95) | Diarrhoea (4.94) | Road Inj (0.81) | Sense (1.13) | COPD (1.27) | NN Preterm (1.15) | HIV (11.36) |
| High-income | IHD (0.61) | Back+Neck (1.09) | Stroke (0.7) | Lung C (1.0) | Alzheimer's (1.51) | Skin (1.1) | Sense (1.1) | Diabetes (1.26) | Depression (1.15) | COPD (1.4) |
| High-income North America | IHD (0.72) | Back+Neck (0.87) | Lung C (1.11) | Drugs (4.14) | Diabetes (1.64) | COPD (2.05) | Skin (1.21) | Stroke (0.65) | Depression (1.26) | Alzheimer's (1.11) |
| Canada | IHD (0.55) | Back+Neck (0.94) | Lung C (1.08) | Skin (1.18) | Diabetes (1.26) | Sense (0.93) | Stroke (0.54) | Migraine (1.0) | Oth MSK (2.36) | Depression (0.97) |
| Greenland | Self Harm (6.21) | Lung C (2.71) | IHD (0.62) | Stroke (1.0) | Back+Neck (0.84) | Depression (1.72) | COPD (1.75) | Skin (1.17) | Alcohol (3.85) | Falls (1.35) |
| USA | IHD (0.74) | Back+Neck (0.86) | Drugs (4.37) | Lung C (1.11) | COPD (2.16) | Diabetes (1.68) | Skin (1.21) | Stroke (0.66) | Depression (1.29) | Road Inj (1.11) |
| Australasia | Back+Neck (1.01) | IHD (0.45) | Skin (1.04) | Depression (1.25) | Migraine (1.05) | Sense (0.88) | Oth MSK (2.45) | Stroke (0.49) | Lung C (0.68) | Alzheimer's (0.93) |
| Australia | Back+Neck (0.99) | IHD (0.44) | Skin (1.03) | Depression (1.27) | Migraine (1.06) | Oth MSK (2.57) | Sense (0.87) | Stroke (0.48) | Lung C (0.67) | Alzheimer's (0.93) |
| New Zealand | Back+Neck (1.19) | IHD (0.54) | Skin (1.1) | Stroke (0.52) | Sense (0.91) | Depression (1.15) | Lung C (0.77) | Migraine (1.0) | COPD (1.19) | Alzheimer's (0.99) |
| High-income Asia Pacific | Back+Neck (1.17) | Alzheimer's (2.28) | Stroke (0.93) | IHD (0.4) | Sense (1.38) | Skin (1.0) | Self Harm (1.38) | Lung C (0.83) | Diabetes (1.26) | Depression (1.04) |
| Brunei | IHD (0.43) | Diabetes (2.22) | Back+Neck (0.75) | Skin (1.09) | Road Inj (1.13) | Stroke (0.54) | Migraine (0.9) | Congenital (1.48) | Falls (0.82) | Sense (0.58) |
| Japan | Back+Neck (1.27) | Alzheimer's (2.72) | IHD (0.45) | Stroke (0.96) | Sense (1.55) | Skin (1.0) | Lung C (0.85) | Self Harm (2.05) | LRI (2.05) | Diabetes (1.11) |
| Singapore | IHD (0.46) | Back+Neck (0.81) | Skin (1.02) | Sense (0.95) | LRI (2.27) | Stroke (0.46) | Alzheimer's (0.97) | Migraine (0.86) | Depression (0.96) | Lung C (0.58) |
| South Korea | Back+Neck (0.95) | Stroke (0.88) | Self Harm (1.77) | Diabetes (1.68) | IHD (0.28) | Alzheimer's (1.32) | Sense (1.0) | Skin (0.99) | Lung C (0.79) | Depression (1.16) |
| Western Europe | Back+Neck (1.24) | IHD (0.63) | Alzheimer's (1.65) | Sense (1.22) | Stroke (0.68) | Lung C (1.04) | Skin (1.06) | Migraine (1.11) | Depression (1.12) | Falls (1.08) |
| Andorra | Back+Neck (1.57) | IHD (0.62) | Alzheimer's (1.65) | Skin (1.13) | Sense (1.08) | Stroke (0.61) | Migraine (1.14) | Lung C (0.73) | Depression (1.06) | Falls (1.03) |
| Austria | IHD (0.8) | Back+Neck (1.11) | Alzheimer's (1.48) | Sense (1.12) | Lung C (0.89) | Migraine (1.22) | Skin (0.98) | Falls (1.25) | Stroke (0.53) | Diabetes (1.01) |
| Belgium | Back+Neck (1.19) | IHD (0.58) | Lung C (1.21) | Alzheimer's (1.64) | Stroke (0.69) | Sense (1.15) | Skin (1.08) | COPD (1.67) | Falls (1.29) | Migraine (1.11) |
| Cyprus | IHD (0.7) | Back+Neck (1.16) | Diabetes (1.67) | Skin (1.06) | Migraine (1.2) | Stroke (0.55) | Sense (0.95) | Alzheimer's (1.2) | Road Inj (0.96) | Lung C (0.64) |
| Denmark | Back+Neck (1.66) | IHD (0.52) | Lung C (1.2) | Stroke (0.73) | COPD (2.15) | Skin (1.16) | Migraine (1.06) | Alzheimer's (0.99) | Diabetes (1.1) | Sense (0.78) |
| Finland | IHD (0.82) | Back+Neck (1.24) | Stroke (0.93) | Alzheimer's (1.77) | Skin (1.14) | Falls (1.48) | Depression (1.33) | Sense (0.98) | Migraine (1.09) | Lung C (0.69) |
| France | Back+Neck (1.11) | IHD (0.42) | Lung C (1.1) | Skin (1.13) | Sense (1.12) | Alzheimer's (1.5) | Stroke (0.56) | Depression (1.23) | Migraine (1.1) | Falls (1.27) |
| Germany | IHD (0.91) | Back+Neck (1.37) | Sense (1.42) | Stroke (0.76) | Lung C (1.11) | Alzheimer's (1.52) | Skin (1.0) | Diabetes (1.28) | COPD (1.5) | Migraine (1.04) |
| Greece | IHD (1.16) | Back+Neck (1.3) | Stroke (1.22) | Lung C (1.12) | Alzheimer's (1.97) | Sense (1.28) | COPD (1.7) | Migraine (1.12) | Skin (0.93) | Depression (1.22) |
| Iceland | Back+Neck (1.11) | IHD (0.52) | Skin (1.19) | Lung C (0.82) | Migraine (1.1) | Sense (0.87) | Alzheimer's (1.05) | Stroke (0.49) | Depression (0.88) | Falls (0.83) |
| Ireland | IHD (0.55) | Back+Neck (1.02) | Skin (1.1) | Migraine (1.12) | Lung C (0.76) | Sense (0.88) | Depression (1.11) | Stroke (0.46) | Alzheimer's (0.97) | COPD (1.18) |
| Israel | Back+Neck (1.1) | IHD (0.34) | Skin (1.11) | Sense (0.97) | Migraine (1.07) | Diabetes (1.08) | Depression (0.97) | Alzheimer's (1.15) | Stroke (0.36) | Lung C (0.65) |
| Italy | Back+Neck (1.29) | IHD (0.65) | Alzheimer's (2.4) | Sense (1.43) | Stroke (0.7) | Lung C (1.08) | Migraine (1.22) | Diabetes (1.31) | Skin (0.96) | Depression (1.04) |
| Luxembourg | Back+Neck (1.1) | IHD (0.48) | Skin (1.12) | Migraine (1.23) | Stroke (0.6) | Lung C (0.79) | Sense (0.93) | Oth Cardio (1.79) | Falls (0.96) | Depression (1.0) |
| Malta | IHD (0.88) | Back+Neck (1.28) | Stroke (0.65) | Skin (1.14) | Sense (1.13) | Migraine (1.15) | Alzheimer's (1.37) | Diabetes (1.16) | Lung C (0.8) | Falls (1.09) |
| Netherlands | Back+Neck (1.25) | IHD (0.46) | Lung C (1.23) | Skin (1.15) | Stroke (0.65) | Sense (0.99) | COPD (1.74) | Migraine (1.14) | Alzheimer's (1.14) | Depression (1.09) |
| Norway | Back+Neck (1.01) | IHD (0.52) | Skin (1.14) | Alzheimer's (1.27) | Stroke (0.57) | Lung C (0.76) | Migraine (0.99) | Sense (0.83) | COPD (1.48) | Falls (1.02) |
| Portugal | Back+Neck (1.45) | IHD (0.57) | Stroke (1.17) | Alzheimer's (2.21) | Sense (1.3) | Diabetes (1.4) | Depression (1.41) | Migraine (1.19) | Skin (0.99) | Lung C (1.04) |
| Spain | Back+Neck (1.09) | IHD (0.47) | Alzheimer's (2.04) | Sense (1.26) | Lung C (0.97) | Migraine (1.21) | Stroke (0.55) | Skin (0.95) | Depression (1.04) | COPD (1.17) |
| Sweden | Back+Neck (1.4) | IHD (0.72) | Stroke (0.69) | Skin (1.22) | Alzheimer's (1.51) | Depression (1.33) | Sense (0.92) | Diabetes (1.21) | Lung C (0.71) | COPD (1.26) |
| Switzerland | Back+Neck (1.29) | IHD (0.52) | Alzheimer's (1.43) | Skin (1.1) | Sense (1.02) | Migraine (1.14) | Lung C (0.77) | Stroke (0.51) | Falls (1.17) | Depression (1.03) |

(Figure 8 continues on next page)

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|---|---------------------|---------------------|---------------------|---------------------|-----------------------|-----------------------|-----------------------|-----------------------|----------------------|-----------------------|
| UK | Back+Neck (1-27) | IHD (0-58) | Lung C (1-08) | Skin (1-2) | Stroke (0-68) | Sense (1-12) | Alzheimer's (1-47) | COPD (1-58) | Migraine (1-05) | Depression (1-13) |
| England | Back+Neck (1-28) | IHD (0-55) | Skin (1-21) | Lung C (1-02) | Stroke (0-64) | Sense (1-14) | Alzheimer's (1-44) | COPD (1-54) | Migraine (1-03) | Depression (1-13) |
| Northern Ireland | IHD (0-66) | Back+Neck (1-12) | Lung C (1-05) | Stroke (0-67) | Skin (1-14) | Alzheimer's (1-33) | Migraine (1-12) | Sense (0-95) | COPD (1-46) | Depression (1-13) |
| Scotland | IHD (0-79) | Back+Neck (1-26) | Stroke (0-94) | Lung C (1-61) | COPD (1-84) | Skin (1-16) | Alzheimer's (1-7) | Sense (1-02) | Migraine (1-18) | Depression (1-19) |
| Wales | IHD (0-76) | Back+Neck (1-24) | Lung C (1-32) | Stroke (0-79) | Alzheimer's (1-93) | Skin (1-15) | COPD (1-74) | Sense (1-04) | Migraine (1-11) | Depression (1-13) |
| Southern Latin America | IHD (0-57) | Back+Neck (1-15) | Stroke (0-67) | Skin (1-07) | LRI (1-59) | Sense (0-98) | Road Inj (0-71) | Diabetes (1-01) | Migraine (0-98) | Oth MSK (1-98) |
| Argentina | IHD (0-66) | Back+Neck (1-12) | LRI (1-91) | Stroke (0-65) | Skin (1-07) | Sense (0-95) | Road Inj (0-73) | Diabetes (1-07) | Migraine (0-97) | Oth MSK (1-91) |
| Chile | Back+Neck (1-22) | IHD (0-36) | Stroke (0-65) | Skin (1-07) | Sense (1-01) | Depression (1-11) | Oth MSK (2-18) | Migraine (0-98) | Road Inj (0-66) | Anxiety (1-39) |
| Uruguay | IHD (0-6) | Back+Neck (1-18) | Stroke (1-0) | Sense (1-17) | Lung C (1-5) | Skin (1-08) | Road Inj (0-76) | COPD (1-23) | Self Harm (1-12) | Migraine (0-99) |
| Central Europe, eastern Europe, and central Asia | IHD (1-94) | Stroke (2-02) | Back+Neck (1-12) | Sense (1-36) | Falls (1-86) | Self Harm (1-47) | Road Inj (1-02) | LRI (2-04) | Lung C (1-01) | Alcohol (3-49) |
| Eastern Europe | IHD (2-47) | Stroke (2-56) | Back+Neck (1-14) | Self Harm (2-03) | Alcohol (5-38) | Sense (1-41) | Falls (1-89) | CMP (5-86) | Road Inj (1-24) | Lung C (0-93) |
| Belarus | IHD (2-78) | Stroke (1-94) | Back+Neck (1-18) | Falls (2-41) | Sense (1-34) | Self Harm (1-67) | Alcohol (4-39) | Road Inj (1-04) | Depression (1-28) | Migraine (1-15) |
| Estonia | IHD (1-54) | Back+Neck (1-15) | Stroke (1-18) | HTN HD (6-72) | Sense (1-34) | Falls (1-78) | Alcohol (4-8) | Lung C (0-91) | Depression (1-25) | Migraine (1-1) |
| Latvia | IHD (2-1) | Stroke (2-44) | Back+Neck (1-24) | Sense (1-57) | Falls (2-03) | CMP (5-83) | Lung C (1-04) | Self Harm (1-38) | Alcohol (3-79) | Migraine (1-12) |
| Lithuania | IHD (2-12) | Stroke (1-78) | Back+Neck (1-2) | Self Harm (2-3) | Falls (2-28) | Sense (1-48) | Lung C (0-95) | Depression (1-38) | Alcohol (3-76) | Migraine (1-06) |
| Moldova | IHD (2-32) | Stroke (1-88) | Back+Neck (1-47) | Sense (1-28) | Migraine (1-26) | Alcohol (3-33) | Falls (1-84) | LRI (1-15) | Cirr Alc (5-2) | Self Harm (1-17) |
| Russia | IHD (2-24) | Stroke (2-73) | Back+Neck (1-06) | Self Harm (2-27) | CMP (7-66) | Alcohol (5-7) | Sense (1-41) | Road Inj (1-35) | Falls (1-88) | LRI (2-06) |
| Ukraine | IHD (3-33) | Stroke (2-27) | Back+Neck (1-43) | Sense (1-46) | Alcohol (4-8) | Falls (1-95) | Self Harm (1-47) | Depression (1-36) | Road Inj (0-91) | Migraine (1-18) |
| Central Europe | IHD (1-38) | Stroke (1-69) | Back+Neck (1-23) | Falls (2-18) | Lung C (1-4) | Sense (1-5) | Diabetes (1-22) | Migraine (1-07) | Colorect C (1-42) | Road Inj (0-87) |
| Albania | IHD (1-24) | Stroke (1-65) | Back+Neck (1-48) | Sense (1-38) | Falls (1-68) | Lung C (1-35) | Migraine (1-11) | Oth Cardio (3-04) | Skin (0-76) | Road Inj (0-52) |
| Bosnia | IHD (1-39) | Stroke (1-91) | Back+Neck (1-5) | Diabetes (2-31) | Lung C (2-41) | Sense (1-66) | Falls (1-92) | Migraine (1-14) | Colorect C (2-22) | Skin (0-73) |
| Bulgaria | IHD (2-29) | Stroke (2-98) | Back+Neck (1-25) | Sense (1-62) | Falls (2-01) | HTN HD (6-4) | Lung C (1-23) | Oth Cardio (3-32) | Diabetes (1-44) | COPD (1-47) |
| Croatia | IHD (1-46) | Stroke (1-86) | Back+Neck (1-29) | Lung C (1-5) | Falls (2-24) | Sense (1-53) | Colorect C (1-8) | Diabetes (1-28) | Migraine (1-06) | Alzheimer's (1-24) |
| Czech Republic | IHD (1-26) | Back+Neck (1-24) | Stroke (1-04) | Falls (2-43) | Sense (1-36) | Lung C (1-03) | Diabetes (1-49) | Migraine (1-05) | Colorect C (1-36) | Oth Cardio (1-97) |
| Hungary | IHD (1-65) | Stroke (1-55) | Back+Neck (1-34) | Lung C (2-08) | Falls (2-49) | Sense (1-44) | COPD (1-82) | Colorect C (2-04) | Diabetes (1-38) | Self Harm (1-27) |
| Macedonia | Stroke (2-52) | IHD (1-13) | CMP (11-79) | Back+Neck (1-17) | Diabetes (1-93) | Lung C (1-49) | Sense (1-31) | Falls (1-65) | Migraine (1-1) | Skin (0-75) |
| Montenegro | IHD (1-23) | Stroke (2-42) | Back+Neck (1-21) | Lung C (1-68) | Sense (1-34) | Falls (1-59) | Diabetes (1-34) | Migraine (1-06) | Road Inj (0-77) | Skin (0-77) |
| Poland | IHD (1-19) | Back+Neck (1-12) | Stroke (1-13) | Falls (2-2) | Lung C (1-38) | Sense (1-57) | Migraine (1-08) | Road Inj (1-01) | Diabetes (1-08) | Self Harm (1-1) |
| Romania | IHD (1-62) | Stroke (2-64) | Back+Neck (1-28) | Falls (2-22) | Sense (1-51) | Lung C (1-27) | Migraine (1-07) | HTN HD (4-23) | LRI (1-91) | Road Inj (0-89) |
| Serbia | IHD (1-28) | Stroke (2-12) | Back+Neck (1-44) | CMP (10-46) | Lung C (2-21) | Sense (1-58) | Diabetes (1-68) | Falls (1-92) | Migraine (1-09) | Colorect C (2-09) |
| Slovakia | IHD (1-44) | Back+Neck (1-22) | Stroke (1-17) | Falls (2-11) | Sense (1-23) | Lung C (0-93) | Migraine (1-08) | Colorect C (1-44) | Skin (0-79) | Road Inj (0-86) |
| Slovenia | IHD (0-73) | Back+Neck (1-28) | Falls (3-07) | Stroke (0-89) | Sense (1-4) | Lung C (1-04) | Migraine (1-05) | Alzheimer's (1-11) | Self Harm (1-15) | Colorect C (1-28) |
| Central Asia | IHD (1-45) | LRI (3-07) | Stroke (1-18) | Back+Neck (0-9) | NN Enceph (3-69) | Sense (1-03) | Road Inj (0-74) | Congenital (1-17) | NN Preterm (1-33) | Migraine (0-98) |
| Armenia | IHD (1-6) | Back+Neck (1-13) | Stroke (0-97) | Diabetes (1-79) | Sense (1-48) | Lung C (1-35) | Migraine (1-04) | Congenital (1-08) | Skin (0-75) | Falls (1-19) |
| Azerbaijan | IHD (1-67) | LRI (3-69) | Stroke (1-14) | Back+Neck (0-93) | NN Enceph (6-42) | Sense (1-1) | Congenital (1-42) | NN Preterm (1-75) | Diabetes (1-13) | Migraine (1-01) |
| Georgia | IHD (2-01) | Stroke (2-37) | Back+Neck (1-13) | Sense (1-63) | Road Inj (0-9) | Diabetes (1-28) | Lung C (1-16) | HTN HD (3-6) | Migraine (1-01) | Falls (1-41) |
| Kazakhstan | IHD (1-47) | Stroke (1-42) | Back+Neck (0-93) | Self Harm (1-79) | Road Inj (1-02) | Sense (1-07) | Congenital (1-37) | LRI (1-4) | COPD (1-09) | Falls (1-37) |

(Figure 8 continues on next page)

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|------------------------------------|--------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|
| Kyrgyzstan | IHD (1-24) | Stroke (1-09) | LRI (1-27) | NN Preterm (1-39) | NN Enceph (2-38) | Back+Neck (1-03) | Congenital (1-04) | Road Inj (0-67) | Sense (0-98) | Iron (2-0) |
| Mongolia | IHD (1-15) | Stroke (1-86) | Liver C (10-82) | LRI (1-72) | Road Inj (1-0) | NN Enceph (3-65) | Back+Neck (0-93) | Self Harm (1-86) | Alcohol (4-1) | Congenital (1-13) |
| Tajikistan | LRI (1-89) | IHD (0-9) | NN Preterm (1-12) | Diarrhoea (1-64) | NN Enceph (1-69) | Stroke (0-71) | Back+Neck (1-03) | Congenital (0-86) | Sense (0-88) | Skin (0-82) |
| Turkmenistan | IHD (1-36) | LRI (4-21) | Stroke (1-3) | Congenital (2-66) | NN Preterm (2-44) | Back+Neck (0-78) | NN Enceph (4-86) | Sense (0-88) | Migraine (0-97) | Skin (0-79) |
| Uzbekistan | IHD (1-56) | LRI (3-33) | Stroke (0-93) | HTN HD (5-73) | Back+Neck (0-9) | NN Enceph (3-53) | Road Inj (0-75) | Iron (3-48) | Diabetes (1-05) | Sense (0-96) |
| Latin America and Caribbean | IHD (0-55) | Violence (3-37) | Diabetes (1-47) | Back+Neck (0-83) | Road Inj (0-86) | Sense (1-1) | Stroke (0-52) | Skin (0-98) | LRI (1-14) | Congenital (0-98) |
| Central Latin America | IHD (0-48) | Violence (3-71) | Diabetes (1-84) | CKD (2-59) | Road Inj (0-77) | Back+Neck (0-67) | Sense (1-06) | Skin (0-92) | Congenital (1-02) | Migraine (0-97) |
| Colombia | Violence (4-91) | IHD (0-47) | Back+Neck (0-8) | Sense (1-18) | Skin (0-96) | Road Inj (0-64) | Migraine (1-04) | Diabetes (0-85) | Congenital (0-74) | Stroke (0-33) |
| Costa Rica | IHD (0-43) | Sense (1-28) | Back+Neck (0-73) | Road Inj (0-7) | Skin (0-9) | Migraine (1-02) | CKD (1-67) | Diabetes (0-8) | Depression (0-9) | Congenital (0-77) |
| El Salvador | Violence (6-04) | IHD (0-64) | CKD (3-35) | Diabetes (1-48) | Road Inj (0-79) | Sense (1-24) | Back+Neck (0-79) | LRI (0-74) | Alcohol (3-17) | Skin (0-94) |
| Guatemala | LRI (1-13) | Violence (3-31) | Diabetes (1-44) | Diarrhoea (1-07) | IHD (0-39) | NN Preterm (0-58) | Congenital (0-76) | Road Inj (0-55) | CKD (1-54) | Skin (1-02) |
| Honduras | IHD (0-86) | Violence (3-01) | Stroke (0-58) | NN Preterm (0-57) | CKD (1-6) | Skin (0-97) | Sense (0-96) | Back+Neck (0-81) | Road Inj (0-46) | Diabetes (0-81) |
| Mexico | Diabetes (2-56) | IHD (0-44) | CKD (3-59) | Violence (2-68) | Road Inj (0-79) | Back+Neck (0-62) | Sense (1-04) | Congenital (1-24) | Skin (0-9) | Migraine (0-94) |
| Nicaragua | IHD (2-22) | CKD (0-42) | Diabetes (1-03) | Sense (1-04) | Back+Neck (0-81) | Skin (0-96) | Congenital (0-65) | Migraine (1-08) | Road Inj (0-44) | LRI (0-39) |
| Panama | IHD (0-37) | Diabetes (1-42) | Sense (1-18) | Violence (3-01) | Back+Neck (0-64) | Road Inj (0-79) | Skin (0-98) | Congenital (1-25) | Stroke (0-44) | Migraine (0-95) |
| Venezuela | Violence (7-22) | IHD (0-62) | Road Inj (1-28) | Diabetes (1-52) | Sense (1-01) | Back+Neck (0-58) | Skin (0-91) | Congenital (1-15) | Stroke (0-46) | NN Preterm (1-25) |
| Andean Latin America | LRI (1-51) | Back+Neck (0-92) | IHD (0-35) | Sense (1-12) | Skin (1-08) | Road Inj (0-68) | Congenital (0-95) | NN Preterm (0-87) | Diabetes (0-83) | Migraine (0-96) |
| Bolivia | LRI (1-64) | NN Preterm (1-19) | IHD (0-55) | Congenital (1-18) | Back+Neck (0-99) | Road Inj (0-7) | Skin (1-12) | Stroke (0-53) | Sense (1-09) | Diabetes (0-96) |
| Ecuador | Road Inj (0-97) | Back+Neck (0-89) | IHD (0-34) | LRI (1-05) | Sense (1-08) | NN Preterm (1-03) | Skin (1-04) | Diabetes (1-06) | Congenital (0-95) | Violence (1-68) |
| Peru | LRI (1-6) | Back+Neck (0-91) | Sense (1-15) | Skin (1-08) | IHD (0-29) | Congenital (0-85) | Road Inj (0-53) | Migraine (0-99) | Diabetes (0-66) | NN Preterm (0-62) |
| Caribbean | IHD (0-81) | Stroke (0-92) | Diabetes (1-73) | HIV (17-88) | LRI (1-49) | Back+Neck (1-49) | Sense (0-75) | Road Inj (0-73) | Skin (1-0) | Diarrhoea (3-89) |
| Antigua | Diabetes (2-43) | IHD (0-47) | Stroke (0-67) | Back+Neck (0-65) | Skin (1-01) | Sense (0-97) | Migraine (1-0) | CKD (2-04) | LRI (1-29) | Oth MSK (1-6) |
| Bahamas | IHD (0-65) | Diabetes (2-23) | HIV (27-02) | Stroke (0-83) | Violence (4-5) | Back+Neck (0-65) | Skin (1-06) | HTN HD (4-72) | Road Inj (0-87) | Sense (1-0) |
| Barbados | Diabetes (3-61) | IHD (0-59) | Stroke (1-03) | Back+Neck (0-73) | Sense (1-28) | Skin (1-23) | LRI (1-97) | Oth MSK (2-32) | CKD (2-32) | Migraine (0-99) |
| Belize | Diabetes (1-6) | IHD (0-47) | Violence (2-13) | HIV (12-73) | Road Inj (0-72) | LRI (0-76) | Skin (1-1) | NN Preterm (0-74) | Back+Neck (0-69) | Stroke (0-43) |
| Bermuda | IHD (0-53) | Diabetes (1-42) | Back+Neck (0-58) | Skin (0-99) | Sense (0-95) | Stroke (0-48) | Migraine (0-96) | Lung C (0-71) | Oth MSK (1-79) | Depression (0-88) |
| Cuba | IHD (0-9) | Stroke (0-92) | Sense (1-47) | Back+Neck (0-73) | Lung C (1-43) | Diabetes (1-22) | Depression (1-27) | Skin (0-92) | Migraine (1-02) | LRI (1-37) |
| Dominica | Diabetes (2-75) | IHD (0-58) | Stroke (0-77) | CKD (2-86) | Skin (1-15) | Back+Neck (0-7) | Road Inj (0-8) | Sense (1-07) | LRI (1-32) | Congenital (1-14) |
| Dominican Republic | IHD (0-68) | HIV (26-69) | NN Preterm (1-61) | Road Inj (0-83) | Stroke (0-61) | Back+Neck (0-73) | NN Sepsis (6-98) | Skin (1-01) | Sense (0-98) | Congenital (1-03) |
| Grenada | IHD (0-71) | Diabetes (2-68) | Stroke (1-02) | LRI (1-82) | CKD (2-51) | Skin (1-11) | Back+Neck (0-68) | Road Inj (0-7) | Sense (0-96) | Oth MSK (2-04) |
| Guyana | IHD (1-17) | Diabetes (2-96) | Stroke (1-34) | Self Harm (2-58) | HIV (16-87) | Road Inj (0-83) | NN Preterm (1-1) | Violence (2-07) | LRI (0-97) | Skin (1-06) |
| Haiti | IHD (1-25) | HIV (30-88) | LRI (0-5) | Diarrhoea (0-74) | Stroke (1-3) | NN Enceph (1-12) | Congenital (1-1) | Road Inj (0-97) | Diabetes (1-78) | Oth NN (1-58) |
| Jamaica | Diabetes (3-24) | Stroke (1-06) | IHD (0-48) | Violence (3-15) | NN Preterm (1-66) | Back+Neck (0-73) | Skin (1-1) | Sense (1-08) | Congenital (1-08) | CKD (1-92) |
| Puerto Rico | Diabetes (3-62) | IHD (0-62) | Back+Neck (0-61) | Sense (1-21) | Violence (6-05) | Skin (1-04) | Stroke (0-56) | CKD (2-7) | LRI (1-95) | Migraine (0-97) |
| St Lucia | Diabetes (2-42) | IHD (0-47) | Stroke (0-82) | Back+Neck (0-77) | Sense (1-12) | Skin (1-04) | Violence (2-39) | Road Inj (0-69) | CKD (1-97) | Migraine (1-04) |
| St Vincent | Diabetes (3-19) | IHD (0-87) | Stroke (1-0) | Violence (2-38) | Back+Neck (0-79) | Skin (1-1) | LRI (0-92) | HIV (10-97) | Sense (1-03) | CKD (1-92) |
| Suriname | IHD (0-78) | Stroke (1-17) | Diabetes (2-15) | NN Preterm (1-96) | Self Harm (1-85) | Road Inj (0-89) | Congenital (1-3) | LRI (1-31) | Skin (1-12) | CKD (2-35) |

(Figure 8 continues on next page)

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|---|------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| Trinidad and Tobago | Diabetes (5-75) | IHD (1-0) | Stroke (0-98) | Violence (4-03) | Back+Neck (0-71) | Road Inj (0-93) | Sense (1-08) | CKD (2-62) | Skin (0-96) | HIV (14-12) |
| Virgin Islands | IHD (1-41) | Diabetes (3-6) | Stroke (1-07) | Back+Neck (0-66) | Violence (7-04) | Sense (1-35) | CKD (3-17) | Prostate C (5-5) | Skin (1-08) | HIV (29-92) |
| Tropical Latin America | IHD (0-62) | Violence (4-1) | Back+Neck (1-03) | Road Inj (1-03) | Stroke (0-68) | Sense (1-13) | Skin (1-03) | Diabetes (1-14) | LRI (1-09) | Migraine (1-05) |
| Brazil | IHD (0-61) | Violence (4-27) | Back+Neck (1-02) | Road Inj (1-04) | Stroke (0-68) | Sense (1-13) | Skin (1-03) | Diabetes (1-13) | LRI (1-13) | Migraine (1-04) |
| Paraguay | IHD (0-6) | Road Inj (1-04) | Diabetes (1-48) | Back+Neck (0-91) | Stroke (0-66) | Skin (1-06) | Congenital (1-0) | NN Preterm (0-93) | Sense (0-98) | Violence (1-72) |
| Southeast Asia, east Asia, and Oceania | Stroke (1-61) | IHD (0-73) | Back+Neck (0-89) | Road Inj (0-95) | COPD (1-51) | Sense (1-18) | Diabetes (1-09) | Lung C (1-42) | Skin (0-87) | Liver C (4-25) |
| East Asia | Stroke (1-83) | IHD (0-73) | Back+Neck (0-89) | Road Inj (1-03) | COPD (1-72) | Lung C (1-68) | Sense (1-2) | Liver C (5-42) | Skin (0-78) | Depression (1-01) |
| China | Stroke (1-83) | IHD (0-74) | Back+Neck (0-9) | Road Inj (1-01) | COPD (1-69) | Sense (1-2) | Congenital (1-71) | NN Preterm (5-45) | Skin (0-78) | Depression (1-02) |
| North Korea | Stroke (2-25) | IHD (1-05) | COPD (2-6) | Lung C (7-46) | Road Inj (0-96) | Congenital (0-93) | NN Preterm (0-56) | Back+Neck (1-36) | Sense (1-3) | LRI (0-29) |
| Taiwan (Province of China) | Diabetes (2-5) | Back+Neck (0-82) | IHD (0-41) | Stroke (0-81) | Sense (1-19) | Liver C (5-73) | Lung C (0-84) | Road Inj (1-05) | Skin (0-84) | CKD (2-23) |
| Southeast Asia | IHD (0-75) | Stroke (1-12) | Back+Neck (0-95) | Diabetes (1-4) | LRI (1-19) | Road Inj (0-75) | Sense (1-13) | Skin (1-07) | TB (5-73) | NN Preterm (0-87) |
| Cambodia | LRI (1-01) | Stroke (1-01) | IHD (0-58) | NN Preterm (0-67) | Road Inj (0-73) | Back+Neck (1-28) | Skin (1-13) | Sense (1-13) | NN Enceph (0-77) | Malaria (70-48) |
| Indonesia | IHD (0-91) | Stroke (1-26) | Diabetes (1-69) | TB (8-74) | Back+Neck (0-94) | NN Preterm (1-06) | Sense (1-07) | Road Inj (0-68) | Skin (1-0) | Diarrhoea (3-07) |
| Laos | LRI (1-7) | NN Preterm (1-69) | Congenital (1-88) | IHD (0-85) | NN Enceph (1-83) | Stroke (1-01) | Diarrhoea (0-82) | Road Inj (0-8) | Skin (1-16) | Back+Neck (1-02) |
| Malaysia | IHD (0-79) | Stroke (0-72) | Road Inj (1-08) | LRI (2-09) | Back+Neck (0-68) | Skin (1-14) | Diabetes (1-29) | Sense (0-72) | Depression (0-89) | COPD (0-9) |
| Maldives | IHD (0-56) | Back+Neck (0-79) | Skin (1-15) | Sense (0-97) | Iron (2-9) | COPD (0-88) | Stroke (0-34) | Migraine (0-81) | Diabetes (0-68) | CKD (1-21) |
| Mauritius | Diabetes (5-86) | IHD (0-97) | CKD (5-25) | Stroke (0-92) | Back+Neck (0-9) | Sense (1-33) | Skin (1-0) | COPD (1-08) | Iron (3-78) | Depression (0-99) |
| Myanmar | Stroke (1-05) | LRI (0-86) | Sense (1-45) | Road Inj (0-75) | IHD (0-41) | Back+Neck (1-04) | COPD (1-25) | Diabetes (1-11) | TB (3-05) | Skin (1-13) |
| Philippines | IHD (0-92) | LRI (1-66) | Stroke (1-03) | Diabetes (1-26) | Back+Neck (0-92) | Skin (1-22) | NN Preterm (0-93) | TB (4-99) | Congenital (0-99) | CKD (1-96) |
| Sri Lanka | IHD (0-83) | Diabetes (2-14) | Back+Neck (0-93) | Sense (1-39) | Stroke (0-68) | Self Harm (1-63) | Skin (0-98) | COPD (1-05) | Asthma (2-37) | Road Inj (0-52) |
| Seychelles | IHD (0-69) | LRI (2-18) | HTN HD (5-93) | Back+Neck (0-84) | Stroke (0-74) | CKD (2-92) | Sense (1-19) | Diabetes (1-23) | Skin (0-98) | Road Inj (0-63) |
| Thailand | Road Inj (1-32) | IHD (0-47) | Back+Neck (1-01) | Stroke (0-84) | Sense (1-35) | LRI (1-58) | Skin (1-13) | Diabetes (1-31) | COPD (1-39) | Liver C (5-35) |
| Timor-Leste | LRI (0-77) | Diarrhoea (1-46) | NN Preterm (0-87) | IHD (0-62) | Congenital (0-94) | Stroke (0-66) | NN Enceph (1-07) | Skin (1-13) | Sense (1-09) | Back+Neck (0-89) |
| Vietnam | Stroke (1-28) | IHD (0-54) | Road Inj (0-93) | Back+Neck (1-07) | Sense (1-19) | Lung C (2-31) | Diabetes (1-03) | Skin (1-0) | COPD (1-02) | Congenital (0-69) |
| Oceania | LRI (0-94) | IHD (1-39) | Stroke (1-41) | Diabetes (2-77) | COPD (2-16) | NN Preterm (0-68) | Congenital (1-01) | Asthma (3-88) | Road Inj (0-76) | Diarrhoea (0-52) |
| American Samoa | Diabetes (3-44) | IHD (0-57) | Skin (1-25) | Stroke (0-67) | Back+Neck (0-68) | CKD (2-45) | Sense (0-74) | LRI (0-99) | COPD (0-87) | Asthma (1-66) |
| Micronesia | IHD (1-35) | Diabetes (3-48) | Stroke (1-51) | CKD (2-71) | LRI (0-91) | COPD (1-4) | Self Harm (1-98) | Road Inj (0-69) | Skin (1-2) | Back+Neck (0-84) |
| Fiji | Diabetes (10-48) | IHD (1-54) | Stroke (1-17) | CKD (4-59) | LRI (2-54) | Back+Neck (0-78) | Asthma (3-59) | Skin (1-09) | NN Preterm (1-48) | Congenital (1-28) |
| Guam | IHD (1-38) | Diabetes (2-92) | Stroke (1-09) | Back+Neck (0-68) | Self Harm (1-4) | Lung C (1-07) | Skin (1-04) | CKD (2-84) | LRI (2-06) | COPD (1-44) |
| Kiribati | Diabetes (5-26) | IHD (1-39) | Stroke (1-77) | LRI (0-41) | NN Preterm (0-65) | Congenital (1-12) | TB (1-56) | NN Enceph (0-95) | Self Harm (2-91) | Diarrhoea (0-41) |
| Marshall Islands | Diabetes (8-59) | IHD (1-03) | Stroke (1-05) | LRI (0-77) | CKD (2-43) | NN Preterm (0-81) | Road Inj (0-6) | Skin (1-12) | Self Harm (1-75) | Back+Neck (0-88) |
| Northern Mariana Islands | Diabetes (1-77) | Back+Neck (0-62) | IHD (0-28) | Skin (0-97) | Stroke (0-47) | Road Inj (0-64) | Self Harm (0-8) | Migraine (0-7) | Sense (0-6) | CKD (1-53) |
| Papua New Guinea | LRI (0-77) | IHD (1-55) | Stroke (1-61) | COPD (2-7) | NN Preterm (0-64) | Diabetes (2-35) | Congenital (1-08) | Asthma (3-88) | Road Inj (0-91) | Diarrhoea (0-32) |
| Samoa | IHD (0-97) | Diabetes (2-31) | Stroke (0-93) | LRI (0-46) | Skin (1-18) | CKD (1-76) | Back+Neck (0-95) | Sense (0-99) | COPD (0-81) | Self Harm (1-17) |
| Solomon Islands | IHD (1-56) | Stroke (1-64) | Diabetes (3-27) | LRI (0-4) | CKD (2-07) | NN Preterm (0-46) | Congenital (0-79) | COPD (1-42) | Road Inj (0-62) | Asthma (2-52) |
| Tonga | Diabetes (3-25) | IHD (0-81) | Stroke (0-7) | LRI (0-65) | NN Preterm (0-73) | CKD (1-9) | Skin (1-18) | Back+Neck (0-93) | Sense (0-94) | Road Inj (0-49) |
| Vanuatu | IHD (1-92) | Stroke (1-73) | LRI (0-66) | Diabetes (2-53) | NN Preterm (0-69) | Congenital (1-05) | COPD (1-43) | CKD (1-85) | Road Inj (0-63) | Iron (1-41) |

(Figure 8 continues on next page)

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|-------------------------------------|--------------------------|-------------------------|-------------------------|-------------------|-------------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| North Africa and Middle East | IHD (0.92) | Conflict Terror (63.83) | Road Inj (1.16) | Congenital (1.69) | NN Preterm (1.52) | Back+Neck (1.06) | Stroke (0.62) | Diabetes (1.23) | LRI (1.06) | Sense (0.98) |
| North Africa and Middle East | IHD (0.92) | Conflict Terror (63.77) | Road Inj (1.16) | Congenital (1.69) | NN Preterm (1.52) | Back+Neck (1.06) | Stroke (0.62) | Diabetes (1.23) | LRI (1.06) | Sense (0.98) |
| Afghanistan | Conflict Terror (32.51) | LRI (0.48) | IHD (2.44) | Congenital (1.99) | Road Inj (2.05) | NN Preterm (0.66) | Stroke (1.37) | Meningitis (1.0) | TB (0.71) | Diarrhoea (0.11) |
| Algeria | IHD (0.74) | Congenital (1.66) | Back+Neck (1.09) | NN Preterm (1.53) | Road Inj (0.96) | Diabetes (1.25) | Sense (1.1) | Stroke (0.52) | Migraine (1.18) | Skin (0.88) |
| Bahrain | Diabetes (2.48) | IHD (0.43) | Back+Neck (0.93) | Migraine (1.19) | Skin (0.95) | Depression (1.22) | Road Inj (0.63) | Sense (0.83) | Oth MSK (1.52) | Anxiety (1.16) |
| Egypt | IHD (1.3) | Road Inj (1.09) | Stroke (0.81) | Back+Neck (1.08) | LRI (1.22) | Diabetes (1.36) | Congenital (1.22) | Diarrhoea (3.39) | Sense (1.08) | Cirr HepC (5.32) |
| Iran | IHD (0.71) | Road Inj (1.73) | Back+Neck (0.94) | NN Preterm (2.36) | Depression (1.59) | Congenital (1.8) | Diabetes (1.39) | Migraine (1.15) | Stroke (0.5) | Sense (0.92) |
| Iraq | Conflict Terror (166.54) | NN Preterm (1.3) | IHD (1.24) | Congenital (1.49) | Diabetes (1.59) | NN Sepsis (2.3) | Road Inj (0.78) | Stroke (0.68) | LRI (0.29) | Back+Neck (1.1) |
| Jordan | Congenital (1.93) | IHD (0.49) | NN Preterm (1.56) | Back+Neck (0.86) | Conflict Terror (36.87) | Diabetes (1.22) | Road Inj (0.71) | Skin (0.93) | Migraine (1.09) | Sense (0.86) |
| Kuwait | IHD (0.42) | Back+Neck (0.79) | Migraine (1.2) | Road Inj (0.94) | Congenital (1.7) | Skin (0.84) | Depression (1.06) | Diabetes (0.95) | Sense (0.72) | Anxiety (1.18) |
| Lebanon | IHD (0.68) | Back+Neck (0.82) | Conflict Terror (29.91) | Sense (1.1) | Migraine (1.15) | Skin (0.96) | Diabetes (1.21) | Depression (1.09) | Congenital (1.23) | Anxiety (1.31) |
| Libya | IHD (0.72) | Conflict Terror (51.15) | Back+Neck (0.85) | Road Inj (1.36) | Diabetes (1.37) | Migraine (1.13) | Stroke (0.52) | Sense (0.95) | Skin (0.91) | Depression (1.07) |
| Morocco | IHD (1.1) | Back+Neck (1.45) | Road Inj (0.94) | Diabetes (1.5) | Depression (1.88) | Stroke (0.59) | Sense (1.15) | TB (3.75) | NN Preterm (0.73) | Migraine (1.32) |
| Palestine | NN Preterm (0.97) | IHD (1.01) | Congenital (1.23) | Stroke (0.51) | Back+Neck (1.09) | Road Inj (0.52) | Diabetes (1.06) | NN Sepsis (1.26) | Skin (0.92) | Depression (1.32) |
| Oman | Road Inj (2.14) | IHD (0.58) | Back+Neck (0.9) | Diabetes (1.59) | Migraine (1.15) | Skin (0.83) | Depression (1.04) | Sense (0.68) | Stroke (0.32) | Congenital (0.72) |
| Qatar | Road Inj (1.75) | Back+Neck (0.84) | Diabetes (1.43) | Migraine (1.18) | IHD (0.24) | Depression (1.13) | Skin (0.82) | Congenital (1.22) | Sense (0.63) | Anxiety (1.14) |
| Saudi Arabia | IHD (0.47) | Road Inj (1.63) | Back+Neck (0.84) | Migraine (1.2) | Skin (0.95) | Sense (0.78) | Depression (0.96) | Stroke (0.39) | Diabetes (0.89) | Congenital (1.11) |
| Sudan | Congenital (2.65) | NN Preterm (1.54) | IHD (1.09) | Road Inj (1.45) | LRI (0.6) | Stroke (0.67) | Iron (2.05) | Back+Neck (1.26) | Diarrhoea (0.76) | Skin (0.97) |
| Syria | Conflict Terror (851.38) | IHD (1.17) | Back+Neck (1.0) | Stroke (0.5) | Sense (0.94) | Migraine (1.15) | Skin (0.89) | Congenital (0.69) | Depression (1.01) | Road Inj (0.4) |
| Tunisia | IHD (0.93) | Back+Neck (1.2) | Road Inj (0.97) | Diabetes (1.41) | Stroke (0.61) | Sense (1.18) | Migraine (1.22) | Skin (0.91) | Depression (1.23) | Congenital (0.93) |
| Turkey | IHD (0.56) | Back+Neck (1.09) | Diabetes (1.36) | Sense (1.07) | Congenital (1.4) | Migraine (1.12) | Skin (0.84) | Stroke (0.44) | Depression (1.03) | Lung C (0.98) |
| United Arab Emirates | Road Inj (2.17) | IHD (0.65) | Back+Neck (0.92) | Stroke (0.62) | Diabetes (1.37) | Migraine (1.14) | Skin (0.79) | Sense (0.76) | Depression (0.9) | CKD (1.64) |
| Yemen | Conflict Terror (171.01) | NN Preterm (1.4) | IHD (1.15) | Congenital (1.57) | Road Inj (1.39) | Diarrhoea (0.76) | Iron (2.73) | LRI (0.27) | Stroke (0.7) | STD (4.05) |
| South Asia | IHD (1.19) | Diarrhoea (1.95) | COPD (2.01) | LRI (0.79) | Stroke (0.77) | NN Preterm (0.8) | Iron (2.49) | TB (2.88) | Road Inj (0.67) | Sense (1.24) |
| South Asia | IHD (1.19) | Diarrhoea (1.95) | COPD (2.01) | LRI (0.79) | Stroke (0.77) | NN Preterm (0.8) | Iron (2.49) | TB (2.88) | Road Inj (0.67) | Sense (1.24) |
| Bangladesh | IHD (0.8) | Stroke (1.03) | LRI (0.38) | Back+Neck (1.19) | NN Enceph (0.95) | Sense (1.23) | COPD (1.1) | Migraine (1.41) | Oth NN (1.6) | Skin (0.95) |
| Bhutan | IHD (0.73) | LRI (0.73) | Iron (2.89) | NN Preterm (0.84) | Back+Neck (0.96) | COPD (1.24) | Congenital (0.76) | Stroke (0.48) | Migraine (1.27) | Diabetes (0.91) |
| India | IHD (1.22) | COPD (2.28) | Diarrhoea (2.47) | LRI (0.9) | Stroke (0.74) | Iron (3.0) | NN Preterm (0.87) | TB (3.61) | Sense (1.3) | Road Inj (0.69) |
| Nepal | IHD (1.06) | LRI (0.39) | NN Enceph (1.01) | COPD (1.67) | Stroke (0.64) | Oth NN (1.53) | Back+Neck (1.23) | Diarrhoea (0.34) | Road Inj (0.52) | Skin (1.0) |
| Pakistan | IHD (1.3) | NN Enceph (2.72) | Diarrhoea (1.36) | NN Preterm (0.86) | LRI (0.56) | Stroke (0.81) | Oth NN (2.71) | Road Inj (0.75) | TB (1.81) | Congenital (0.69) |
| Sub-Saharan Africa | Malaria (138.47) | HIV (54.6) | Diarrhoea (1.23) | LRI (0.78) | NN Enceph (1.55) | NN Preterm (0.77) | TB (1.69) | PEM (1.94) | NN Sepsis (2.07) | Congenital (0.91) |
| Southern sub-Saharan Africa | HIV (226.35) | LRI (3.43) | TB (17.53) | Diarrhoea (9.23) | Road Inj (1.56) | Violence (4.08) | Diabetes (2.03) | NN Preterm (1.6) | IHD (0.42) | Stroke (0.64) |

(Figure 8 continues on next page)

| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|-----------------------------------|------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| Botswana | HIV (166-06) | Diabetes (1-95) | TB (13-78) | LRI (1-95) | Diarrhoea (7-09) | IHD (0-43) | Road Inj (0-9) | Skin (1-14) | Stroke (0-57) | Back+Neck (0-67) |
| Lesotho | HIV (316-34) | TB (14-68) | Diarrhoea (5-4) | LRI (1-71) | Road Inj (1-51) | Violence (3-5) | NN Preterm (1-18) | Diabetes (2-16) | Oth NN (4-64) | Stroke (1-0) |
| Namibia | HIV (117-45) | Diarrhoea (6-46) | LRI (1-76) | Road Inj (1-05) | TB (6-58) | NN Preterm (1-15) | Oth NN (4-82) | Violence (1-92) | IHD (0-38) | Skin (1-18) |
| South Africa | HIV (287-36) | LRI (3-97) | Road Inj (1-99) | Violence (6-29) | TB (23-49) | Diabetes (2-47) | IHD (0-42) | Diarrhoea (8-27) | Stroke (0-63) | Back+Neck (0-7) |
| Swaziland | HIV (200-02) | Diarrhoea (7-1) | LRI (2-31) | TB (8-93) | Road Inj (1-29) | Diabetes (2-1) | NN Preterm (1-11) | Oth NN (4-58) | Violence (1-97) | IHD (0-43) |
| Zimbabwe | HIV (133-99) | Diarrhoea (2-09) | LRI (1-09) | TB (3-84) | NN Enceph (2-11) | NN Preterm (1-04) | Congenital (0-99) | PEM (3-05) | NN Sepsis (2-47) | Road Inj (0-73) |
| Western sub-Saharan Africa | Malaria (195-78) | Diarrhoea (1-4) | LRI (0-78) | HIV (39-08) | NN Enceph (1-95) | NN Preterm (0-92) | Congenital (1-44) | Meningitis (2-83) | NN Sepsis (2-32) | PEM (1-6) |
| Benin | Malaria (23-7) | Diarrhoea (0-71) | LRI (0-51) | NN Enceph (1-18) | NN Preterm (0-74) | Congenital (1-2) | Meningitis (1-31) | NN Sepsis (1-16) | HIV (9-78) | Iron (1-06) |
| Burkina Faso | Malaria (14-54) | LRI (0-62) | Diarrhoea (0-28) | Congenital (1-7) | NN Preterm (0-61) | Meningitis (1-25) | NN Enceph (0-7) | PEM (0-77) | NN Sepsis (1-18) | Iron (1-32) |
| Cameroon | Malaria (274-08) | HIV (76-33) | LRI (1-04) | Diarrhoea (0-96) | Congenital (1-78) | NN Preterm (0-99) | NN Enceph (1-53) | Meningitis (2-84) | NN Sepsis (2-14) | PEM (1-52) |
| Cape Verde | IHD (0-55) | HIV (14-08) | LRI (0-59) | Back+Neck (0-99) | NN Preterm (0-6) | Violence (1-45) | Skin (1-06) | Stroke (0-48) | Congenital (0-69) | Iron (1-71) |
| Chad | Diarrhoea (0-81) | LRI (0-89) | HIV (25-23) | Malaria (5-13) | NN Enceph (1-21) | NN Preterm (0-8) | Meningitis (1-66) | PEM (1-17) | STD (2-59) | Congenital (1-29) |
| Côte d'Ivoire | Malaria (89-87) | Diarrhoea (1-04) | LRI (50-5) | LRI (0-75) | NN Preterm (1-01) | Congenital (1-64) | NN Enceph (1-42) | IHD (0-75) | NN Sepsis (1-78) | Stroke (0-79) |
| The Gambia | LRI (0-39) | NN Preterm (0-74) | Diarrhoea (0-32) | HIV (20-9) | NN Enceph (0-9) | Congenital (1-0) | NN Sepsis (1-35) | Iron (1-42) | Meningitis (1-2) | Malaria (4-5) |
| Ghana | Malaria (431-66) | HIV (32-92) | NN Enceph (2-24) | LRI (0-72) | Congenital (1-11) | NN Sepsis (3-02) | NN Preterm (0-59) | Stroke (0-69) | IHD (0-49) | Meningitis (2-84) |
| Guinea | Malaria (19-15) | LRI (0-7) | NN Enceph (1-27) | Diarrhoea (0-27) | NN Preterm (0-68) | Congenital (1-47) | HIV (16-26) | Meningitis (1-37) | TB (0-78) | NN Sepsis (1-2) |
| Guinea-Bissau | HIV (45-01) | LRI (0-49) | Diarrhoea (0-4) | NN Enceph (1-13) | NN Preterm (0-75) | TB (1-34) | STD (3-3) | Meningitis (1-72) | IHD (1-02) | Congenital (1-16) |
| Liberia | Diarrhoea (0-57) | Malaria (10-62) | LRI (0-32) | HIV (17-67) | NN Enceph (0-86) | NN Preterm (0-48) | Congenital (0-97) | IHD (0-61) | Meningitis (0-92) | TB (0-6) |
| Mali | Malaria (11-12) | Diarrhoea (0-55) | NN Enceph (1-38) | NN Preterm (0-91) | LRI (0-3) | PEM (1-08) | Congenital (1-32) | STD (2-03) | Meningitis (1-1) | NN Sepsis (1-17) |
| Mauritania | LRI (0-62) | Diarrhoea (1-04) | NN Preterm (0-77) | NN Enceph (1-22) | Congenital (1-0) | NN Sepsis (2-06) | IHD (0-45) | Back+Neck (1-13) | Iron (1-43) | Skin (1-11) |
| Niger | Malaria (3-36) | Diarrhoea (0-52) | LRI (0-51) | Meningitis (1-0) | NN Enceph (0-74) | NN Preterm (0-48) | Congenital (1-0) | PEM (0-5) | NN Sepsis (0-93) | TB (0-32) |
| Nigeria | Malaria (594-41) | Diarrhoea (2-59) | HIV (50-48) | NN Enceph (2-98) | LRI (0-97) | NN Preterm (1-15) | Congenital (1-52) | NN Sepsis (3-49) | Meningitis (3-93) | PEM (2-71) |
| São Tomé and Príncipe | LRI (0-45) | Congenital (0-83) | Diarrhoea (0-34) | NN Enceph (0-72) | NN Preterm (0-43) | Stroke (0-57) | NN Sepsis (1-35) | Skin (1-08) | Back+Neck (0-95) | PEM (0-94) |
| Senegal | Diarrhoea (0-41) | LRI (0-39) | NN Preterm (0-58) | NN Enceph (0-87) | Congenital (1-11) | Malaria (7-08) | Meningitis (1-4) | NN Sepsis (1-27) | IHD (0-62) | TB (0-75) |
| Sierra Leone | Malaria (46-59) | Diarrhoea (0-87) | LRI (0-82) | NN Enceph (1-63) | Congenital (2-11) | NN Preterm (0-76) | Meningitis (2-1) | HIV (17-7) | TB (0-85) | NN Sepsis (1-36) |
| Togo | Malaria (83-04) | HIV (38-48) | LRI (0-49) | Diarrhoea (0-47) | NN Enceph (1-17) | NN Preterm (0-7) | Congenital (1-13) | IHD (0-64) | TB (0-87) | NN Sepsis (1-34) |
| Eastern sub-Saharan Africa | HIV (45-37) | LRI (0-57) | Diarrhoea (0-61) | Malaria (17-5) | NN Enceph (1-04) | TB (1-35) | NN Preterm (0-53) | PEM (1-28) | Meningitis (1-5) | NN Sepsis (1-37) |
| Burundi | Diarrhoea (0-63) | LRI (0-52) | Malaria (4-13) | TB (1-73) | NN Enceph (1-04) | PEM (1-11) | NN Preterm (0-65) | NN Sepsis (1-24) | Meningitis (0-85) | HIV (8-1) |
| Comoros | LRI (0-61) | Diarrhoea (0-9) | NN Enceph (1-04) | NN Preterm (0-59) | TB (1-53) | IHD (0-47) | PEM (1-5) | Skin (1-24) | NN Sepsis (1-57) | Meningitis (1-78) |
| Djibouti | HIV (44-64) | LRI (0-67) | PEM (2-36) | NN Preterm (0-7) | TB (1-61) | Diarrhoea (0-41) | NN Enceph (0-87) | NN Sepsis (1-73) | IHD (0-56) | Meningitis (1-73) |
| Eritrea | Diarrhoea (0-71) | LRI (0-52) | TB (1-99) | PEM (1-5) | NN Enceph (0-7) | NN Preterm (0-42) | Meningitis (1-51) | HIV (11-79) | NN Sepsis (1-2) | Skin (1-2) |
| Ethiopia | Diarrhoea (0-29) | LRI (0-34) | TB (0-98) | NN Enceph (0-7) | NN Sepsis (1-02) | IHD (0-62) | HIV (8-87) | NN Preterm (0-31) | PEM (0-61) | Meningitis (0-85) |
| Kenya | HIV (64-38) | Diarrhoea (3-17) | LRI (1-0) | NN Enceph (1-3) | NN Preterm (0-63) | Skin (1-25) | TB (1-6) | Malaria (92-97) | NN Sepsis (1-89) | Meningitis (2-45) |
| Madagascar | Diarrhoea (0-88) | LRI (0-68) | PEM (2-55) | NN Preterm (0-85) | Stroke (1-1) | Malaria (9-15) | NN Sepsis (1-48) | NN Enceph (0-68) | STD (2-54) | Meningitis (1-27) |
| Malawi | HIV (93-83) | Malaria (18-86) | LRI (0-6) | Diarrhoea (0-45) | NN Enceph (1-32) | NN Preterm (0-65) | PEM (1-13) | Meningitis (1-47) | TB (0-89) | NN Sepsis (1-35) |
| Mozambique | HIV (92-49) | Malaria (8-96) | LRI (0-46) | NN Enceph (0-98) | TB (1-18) | Diarrhoea (0-19) | NN Preterm (0-5) | NN Sepsis (1-2) | Oth NN (0-87) | STD (1-41) |
| Rwanda | LRI (0-55) | Diarrhoea (0-41) | TB (1-25) | NN Enceph (0-86) | Malaria (15-27) | HIV (16-46) | PEM (1-21) | NN Preterm (0-42) | Meningitis (1-41) | NN Sepsis (1-22) |

(Figure 8 continues on next page)




Figure 8: Leading ten causes of all-age DALYs with the ratio of observed to expected DALYs on the basis of Socio-demographic Index in 2016, by location

The ratio of observed to expected DALYs on the basis of Socio-demographic Index is provided in brackets for each cause and cells are colour-coded by ratio ranges (calculated to place a roughly equal number of cells into each bin). Shades of blue represent much lower observed DALY levels than expected on the basis of Socio-demographic Index, whereas red shows observed DALYs that exceed expected levels. Alcohol=alcohol use disorders. Alzheimer's=Alzheimer's disease and other dementias. Anxiety=anxiety disorders. Back+Neck=low back and neck pain. Cirr alc=cirrhosis due to alcohol use. Cirr HepC=cirrhosis due to hepatitis C. CKD=chronic kidney disease. CMP=cardiomyopathy and myocarditis. Colorect C=colon and rectum cancer. Conflict Terror=conflict and terrorism. Congenital=congenital anomalies. COPD=chronic obstructive pulmonary disease. Depression=depressive disorders. Diabetes=diabetes mellitus. Diarrhoea=diarrhoeal diseases. DR Congo=Democratic Republic of the Congo. Drugs=drug use disorders. HIV=HIV/AIDS. HTN HD=hypertensive heart disease. IHD=ischaemic heart disease. Iron=iron-deficiency anaemia. Liver C=liver cancer. LRI=lower respiratory infections. Lung C=lung, bronchus, and trachea cancers. NN Enceph=neonatal encephalopathy due to birth asphyxia and trauma. NN Preterm=neonatal preterm birth complications. NN Sepsis=neonatal sepsis and other neonatal infections. Oth Cardio=other cardiovascular and circulatory diseases. Oth MSK=other musculoskeletal disorders. Oth NN=Other neonatal disorders. PEM=protein-energy malnutrition. Prostate C=prostate cancer. Road Inj=road injuries. Sense=sense organ diseases. Skin=skin and subcutaneous diseases. STD=sexually transmitted diseases excluding HIV. Stomach C=stomach cancer. Stroke=Cerebrovascular disease. TB=tuberculosis. Violence=interpersonal violence.

GBD 2015 results.⁴ The conclusion that DALY convergence is likely to be accelerated by escalating increases in SDI remains valid. We have looked a step further in GBD 2016 and highlighted countries that have most improved and showed the largest declines in their DALY performance relative to that expected on the basis of their SDI.

The pace and efficiency of a country's transition through the SDI quintiles and the concurrent improvements in population summary health metrics is by no means inevitable, however, and analysis derived from the ratio of observed to expected DALYs on the basis of SDI has been informative. The exemplars were Nicaragua, Costa Rica, the Maldives, Peru, and Israel, which had the lowest levels of age-standardised DALYs relative to the levels expected on the basis of SDI in 2016. The appendix (pp 63–66) contains a full list of exemplars. These countries considerably outperformed health expectations relative to their development status. Conversely, Lesotho, Swaziland, South Africa, Fiji, and Botswana had the highest levels of age-standardised DALYs relative to those expected, reflecting a need for additional attention and examination of the reasons for these discrepancies. Lesotho had the highest ratio of age-standardised DALYs relative to that expected, which

indicates poor performance in terms of health outcomes for mortality and morbidity relative to their SDI status.

The reasons for the success in health outcomes of the five exemplars merit in-depth analyses. Conversely, with the exception of Fiji (which is heavily impacted by the high levels of diabetes seen throughout Oceania), the poor performers in 2016 were clearly those that were most devastated by the continuing HIV/AIDS epidemic and, although progress has been made in South Africa, Swaziland, Zimbabwe, and Lesotho are still disproportionately affected. Many different locations within the top ten exemplars and ten poor performers would make excellent case studies to assess the reasons for their relative rankings, particularly given the different health challenges and successes for each nation.

By highlighting exemplars and poor performers across the SDI quintiles, we have seen that the underlying influences that have contributed to the relative successes and failures of nations are substantially heterogeneous. A clear corollary is that the policy implications for reform will not be the same between quintiles. Nation-by-nation complexity in how countries progress through the SDI continuum and how population health summaries track is clearly too voluminous to document in this study by

nation, but we hope that the results of this study will spawn a plethora of derivative and policy-relevant research.

We have highlighted key exemplar nations and those whose progress could be improved, but there are clearly more general lessons that can be learned from the epidemiological transition across all causes and locations. The collection of London Declaration NTDs, along with malaria²⁸ and HIV, all show how effective domestic and international collaborations in fighting of specific infectious diseases can be. Whether or not these collaborations provide evidence of health-care improvements in one area, whether or not the benefits cascade, and whether or not non-health-specific interventions, such as poverty alleviation, education,²⁹ and family planning, are having synergistic effects are hard to differentiate,^{28,30} but a host of online tools exist that enable national experts to examine performance of and changes in the leading causes of disease burden, compare them with peers, and objectively assess what can be improved.

Since the concept of the DALY was introduced two decades ago,³¹ it has become a key metric for monitoring of population health and prioritisation within health sectors.^{32–39} The following are examples of policy makers, funders, or foundations that use DALYs in decision making: WHO, the World Bank, the National Institutes of Health, the US Centers for Disease Control and Prevention, the Bill & Melinda Gates Foundation, Gavi, the US President's Emergency Plan for AIDS Relief, the Global Fund, and the Wellcome Trust, as well as the Chinese Politburo, the National Institute for Health and Care Excellence, Indonesia Bappenas, and Public Health England at the national level.

Comparison of GBD 2016 with other global estimates and GBD 2015

The GBD study is the only source of comprehensive quantification of population health summary measures, including YLLs, YLDs, DALYs, and HALE. Specific efforts that are relevant to policy makers are being made to estimate burden within other organisations. Since GBD 2015, most organisations that we assessed have not produced updated estimates for DALYs or HALE. The exception to this is WHO, which has released updated Global Health Estimates for DALYs and HALE for 183 countries from 2000 to 2015.⁴⁰ These estimates draw heavily on the GBD 2015 results, with revisions to the all-cause mortality envelope^{40,41} as detailed in the GBD 2016 cause-specific mortality publication¹⁰ and to selected cause-specific disability weights and severity distributions for YLDs.⁴² WHO-adjusted disability weights for various causes are further detailed in the WHO Global Health Estimates Technical Paper.⁴¹

Disease-specific considerations

The SDI transition was prominent for many CMNN conditions, especially diarrhoea, lower respiratory

infection, HIV after 2005 (the year of the global peak), maternal disorders, and vaccine-preventable diseases like measles, tetanus, diphtheria, and pertussis. Focused programmes with community-wide targeting of health interventions—such as access to antenatal care,^{43,44} vaccination, malaria vector control (insecticide-treated bednets and indoor residual spraying),⁴⁵ and artemisinin combination therapies,⁴⁶ water and sanitation efforts,⁴⁷ and ART and prevention of mother-to-child transmission of HIV^{30,48,49}—are potentially responsible for DALY reductions in these conditions.⁵⁰ Further scale-up and maintenance of these interventions should continue, including a transition from non-governmental organisations to local ownership and financing. Corresponding improvements in health system functions related to survival of mothers and their children have not occurred,⁵¹ shown by the comparatively slow improvement in DALYs due to maternal and neonatal disorders compared with other CMNN causes and the growing burden of congenital birth defects, haemoglobinopathies and haemolytic anaemias, and sudden infant death syndrome, especially in the under-5 age group. Early detection of complications of pregnancy like micronutrient deficiencies, hypertensive disorders of pregnancy, pregnancy-transmitted and sexually transmitted infections, and congenital birth defects can help identify the pregnant women and newborns at highest risk, help ensure that appropriate curative treatments are administered, and put the females and newborns in a position to receive timely perinatal care when it is needed. Newborn screening for congenital birth defects and haemoglobinopathies can, at the very least, identify children with these conditions and facilitate maternal education about the importance of presentation for care early when a mother's child gets sick. Delays in seeking care can have deadly consequences, particularly for children with acute conditions. Education of mothers and fathers, especially first-time parents, about how to best care for their babies can help reduce avoidable injuries and deaths. At the same time, stark increases in neonatal sepsis DALYs, in which nosocomial infections of the newborn are included, highlight the crucial importance of being vigilant in infection control within hospitals and clinics.

This iteration of GBD estimated diseases and injuries for 5 year age groups older than 80 years for the first time: 80–84 years, 85–89 years, 90–94 years, and 95 years and older. This addition has provided a clearer picture than from previous GBD publications of the burden of disease in ageing populations, who are disproportionately affected by dementias and other NCDs compared with younger age groups. Ageing populations have increased substantially in size since 1990. As a result of this population shift and the epidemiological transition (which increases the proportion of burden of disease due to NCDs), age groups older than 80 years had increases in all-age DALYs between

1990 and 2016, with increases across all SDI quintiles. The proportion due to NCDs also increased. Therefore, research into these age groups and the diseases that continue to affect them over time should be prioritised.

Tuberculosis

We made important changes to the modelling strategy for tuberculosis in GBD 2016. For fatal tuberculosis, we first modelled the prevalence of active disease and latent infection, which we then used as covariates for the Cause of Death Ensemble model. For non-fatal tuberculosis, we strengthened our statistical triangulation approach, which enforces consistency between data for different parameters by modelling tuberculosis incidence, prevalence, and mortality among those with latent infection. Application of MIRs estimated on the basis of SDI to better reflect incidence in low-income and middle-income countries has also enhanced consistency between fatal and non-fatal estimates of tuberculosis. All of these changes have resulted in global tuberculosis DALYs that are 15% higher than in GBD 2015 for the year 2010, with prominent increases occurring in several African countries, including Uganda, the Central African Republic, and Zambia.

HIV/AIDS

A major HIV methods change for GBD 2016 was the distribution of ART coverage by age, sex, and CD4-positive cell count. We used two AIDS Indicator Surveys^{52,53} to predict the age-sex-CD4 cell distribution of ART coverage and applied the distributions to the input counts of people receiving ART in our HIV estimation model. This method shifted the coverage distribution to groups with higher CD4 cell counts, as was seen in the data. All of these changes have resulted in higher global HIV/AIDS DALYs than in GBD 2015 for the year 2010, with the most prominent increases occurring in several African countries, including South Africa, Botswana, Lesotho, and Swaziland. We have also systematically updated other key input parameters to the HIV/AIDS estimation process, such as the on-ART mortality rate and other demographic inputs, including HIV-free mortality.

Lower respiratory infection and diarrhoea

Total lower respiratory infection (increased by 0·701%) and diarrhoeal disease (increased by 15·5%) DALYs increased compared with GBD 2015 estimates for the year 2010. Like most CMNN causes, the lower respiratory infection and diarrhoeal disease DALY burden is due primarily to YLLs; more than 90% of global DALYs are from these causes. Several changes have been made to the under-5 mortality models that have large impacts on the DALY totals for these causes. We added several new model covariates on the basis of risk factors associated with lower respiratory infections and diarrhoeal diseases, including childhood stunting and suboptimal breastfeeding. Additionally, because of

India's large population size and disease burden, its Sample Registration System data changed the magnitude of YLLs. Inclusion of diarrhoeal diseases as a fatal discontinuity in YLL estimation also affected the overall estimation for GBD 2016. Overall, the YLLs due to lower respiratory infections decreased slightly (by 0·626%) for the year 2010 from GBD 2015 to GBD 2016 and those due to diarrhoeal diseases increased (by 14·3%). We believe that these modelling changes improved under-5 mortality estimates for lower respiratory infections and diarrhoea. Although responsible for a smaller contribution to DALYs than cause of death models, the non-fatal models of lower respiratory infections and diarrhoea included new data sources and processing. An example is that we now account for seasonality from the population-representative surveys that provide most prevalence data from sub-Saharan Africa and south Asia in the lower respiratory infection and diarrhoeal disease models. Another major update is a change in the estimated mean duration of illness on the basis of a set of updated systematic reviews. The duration of diarrhoea remained nearly the same as in GBD 2015, but that of lower respiratory infection decreased by about 20% compared with GBD 2015.

Malaria

Refinements to the methodological approach and addition of substantially more data than in GBD 2015 have led to changes in estimates of both malaria mortality and morbidity estimates in the GBD 2016 iteration, with resulting changes in DALYs. Globally, predicted trends in malaria were similar to GBD 2015, rising to a peak in 2005 before steadily declining. Overall malaria DALYs are lower in GBD 2016 than in GBD 2015, reflecting mainly lower estimates outside of Africa and particularly for India. Outside of Africa, and for lower-burden countries within Africa, estimates were informed for the first time by extensive subnational case-reporting data from routine surveillance systems. These were subsequently adjusted to account for under-reporting, misdiagnosis, and incompleteness, and then entered into a spatiotemporal geostatistical model to infer continuous surfaces of incidence rate before re-aggregating them to national and subnational totals. This approach led to notable reductions in estimated cases, YLDs, and DALYs in India, Myanmar, Indonesia, and Pakistan. In high-burden countries in sub-Saharan Africa, where the methods remained similar to GBD 2015, changes were relatively modest and reflected the inclusion of newly available cross-sectional parasite rate surveys or updates to data for malaria intervention coverage in recent years.

London Declaration diseases

The London Declaration was established in 2012 as a partnership of pharmaceutical companies, private

foundations, and global health organisations, committed to providing resources and expertise for controlling, eliminating, or eradicating ten NTDs: human African trypanosomiasis, Chagas disease, Guinea worm disease, leprosy, lymphatic filariasis, onchocerciasis, schistosomiasis, soil-transmitted helminths, blinding trachoma, and visceral leishmaniasis.⁵⁴ With this year's addition of Guinea worm disease to the GBD cause list, we now estimate DALYs for all ten of the London Declaration diseases and GBD estimates can, therefore, offer insight into progress. We estimate that total DALYs from the London Declaration NTDs have declined by 21·1% from 11·4 million (95% UI 6·8 million to 18·5 million) DALYs in 2010 before the London Declaration in 2012 to 9·0 million (5·3 million to 14·5 million) in 2016. Moreover, we find that DALY rates have declined for all ten of the London Declaration NTDs between 1990 and 2016, with the largest declines occurring for Guinea worm disease, reflecting the particular effectiveness of the Guinea worm eradication initiative,⁵⁵ and for human African trypanosomiasis.⁵⁶

Zika virus disease

The appearance of Zika virus cases in the Americas in 2015, along with the reported associations between Zika virus infection and microcephaly and Guillain-Barré syndrome, led WHO to declare Zika virus a Public Health Emergency of International Concern in February 2016.^{57–59} We estimated Zika virus disease for GBD 2016 in response to this heightened global concern and interest. We estimated four non-fatal outcomes of Zika virus infection: asymptomatic infection, symptomatic infection, Guillain-Barré syndrome caused by Zika virus infection, and congenital Zika syndrome. Despite the high incidence observed in the Americas in 2016 and understandable public concern, especially surrounding congenital Zika syndrome, we estimate that less than 0·01% of Zika virus infections result in Guillain-Barré syndrome, congenital Zika syndrome, or death. We estimate that 7·60 million (95% UI 5·70 million to 10·7 million) infections occurred in 2016, resulting in 2·70 million symptomatic Zika virus cases, 1880 cases of Zika-attributable Guillain-Barré syndrome, and 2400 congenital Zika syndrome births.⁸ Our estimate for congenital Zika syndrome is in line with official reports from the Pan American Health Organization of circa 2500 congenital Zika syndrome births.⁶⁰ However, those born with congenital Zika syndrome will have future disability.

Injuries

Global DALYs due to injuries generally contributed to a lower proportion of total burden than did those due to CMNN diseases and NCDs, and remained relatively stable as a proportion of total DALYs over time. Examination of these trends by SDI quintile, region, and cause of injury reveals much more variable patterns

than the global pattern, which are potentially related to a confluence of demographic changes, policy adoption and enforcement, access to high-quality trauma care, and political stability. Among high-SDI and high-middle-SDI countries, all-age DALYs due to road injuries have substantially declined since 1990, while DALYs due to road injuries in low-middle-SDI and low-SDI countries have risen, a trend that is particularly evident in recent years. By contrast, DALYs from drowning markedly declined among middle-SDI and low-middle-SDI countries. Disease burden from falls rose across the development spectrum, a trend primarily driven by population growth and ageing. Finally, the consequences of ongoing conflict, particularly in north Africa and the Middle East, and interpersonal violence, particularly in Latin America, on population health cannot be overlooked.⁶¹ Fatalities due to such conflict and violence have resulted in stagnated or decreasing life expectancy in many of these countries,^{7,62} and for those who survive, the long-term effects of such injuries could easily result in impaired movement and functioning, heightened risk of other disorders (eg, musculoskeletal conditions), and mental health challenges for an extended period after the end of the conflict.

Cerebrovascular disease

For GBD 2016, we modelled each stroke subtype independently to produce more reliable ratios of ischaemic to haemorrhagic stroke than for GBD 2015 and better match our independent estimation of subtype-specific stroke mortality. We avoided undercounting of stroke by reclassifying hospital admissions and deaths ascribed to unspecified stroke. Despite this method, haemorrhagic stroke remains a heterogeneous category that includes neonatal intraventricular haemorrhage and all other non-traumatic intracranial bleeding. These deaths in children younger than 5 years result in many more YLLs, and therefore many more DALYs, than from ischaemic stroke. Consistent with population-based and multinational studies of stroke subtype, GBD estimates higher incidence but much lower case fatality and YLLs due to ischaemic stroke among adults than due to haemorrhagic stroke.⁶³ This pattern appears true even for locations where this pattern was not previously thought to be the case, such as in China.⁶⁴ Future estimates can be improved by production of separate estimates of non-traumatic subarachnoid haemorrhage and paediatric stroke.

Mental and substance use disorders

Throughout multiple iterations of GBD, mental and substance use disorders have consistently been shown as the leading causes of YLDs worldwide, with burden present in both sexes across the lifespan. They are also strongly associated with premature mortality, although this association is not reflected in GBD YLL estimates for mental disorders as they are rarely coded as the direct

cause of death. Nevertheless, they contribute a substantial number of DALYs and this large contribution to burden has remained constant across time in all countries, including those with high or substantially improving SDI. Treatment rates remain very low^{65–67} and, even in high-income countries where treatment coverage has increased, the prevalence of the most common disorders has not changed.⁶⁸ To reduce the burden of these disorders, improved treatment coverage needs to include a focus on the quality of the intervention delivered. Additionally, identification and quantification of modifiable risk factors for mental and substance use disorders are vital for development of effective prevention strategies and are an area noted for expansion in future iterations of GBD.

Diabetes

We made several important improvements to the process of estimation of diabetes prevalence, including use of more data sources than GBD 2015 and development of a novel approach to standardise the definition of diabetes across different sources. In our assessment of diagnostic criteria for diabetes across different surveys, we identified more than 50 different definitions for diabetes based on various biomarkers (eg, fasting plasma glucose concentration, oral glucose tolerance test result, and glycated haemoglobin A1c concentration) and different levels of each biomarker. To standardise the definition of diabetes, we mostly focused on the surveys that had included fasting plasma glucose concentration as a diagnostic criterion and developed an ensemble model to characterise the distribution of fasting plasma glucose concentration at the population level in each age and sex group. Then, we used the fasting plasma glucose concentration distribution to convert various definitions of diabetes into the standard case definition. Using the ensemble model, we also estimated the prevalence of diabetes on the basis of the mean fasting plasma glucose concentration in places for which we only had data for mean fasting plasma glucose concentration. These changes allowed us to be more consistent than in GBD 2015 in our estimation of the prevalence of diabetes across countries and over time. As a result, our estimates of the prevalence of diabetes globally and in most regions are slightly lower than those reported in GBD 2015. The strong relationship between SDI and diabetes remained, mirroring global rises in overweight and obesity.⁶⁹

Cancer

DALYs for cancer have changed compared with GBD 2015; this change is predominantly due to lower YLD estimates for GBD 2016 than for GBD 2015. These improvements stem from adjustments made in the modelling of MIRs to better reflect differences in MIR based on SDI in data-sparse locations than in GBD 2015. In addition to stricter inclusion criteria for data used in the MIR modelling than in GBD 2015, we changed the

modelling approach and used the most parsimonious model with just SDI as a predictor of MIR.⁸ MIRs are used to estimate cancer incidence and prevalence from GBD cancer mortality estimates and therefore directly determine YLDs. Until cancer registry incidence data and accurate mortality statistics are widely available, validation of MIR is difficult in countries that lack these data sources. The Global Initiative for Cancer Registry Development and expansion of civil registration systems are therefore crucial to further improve estimation of cancer burden.

Future directions

Challenging data gaps exist in the severity distributions across sequelae for most diseases in the YLD literature. Most data sources for severity are from high-income countries, which probably leads to an underestimation of YLDs in low-income and middle-income countries where the severity of presentation of non-fatal illnesses might be worse than in high-income countries, frequently as a result of late diagnosis and underdiagnosis. Improvements can be made if disease-specific research focuses on routine use of a single established measure of severity in surveys and patient populations and if countries are able to link survey data to a general health assessment instrument, along with improvements in early diagnosis and treatment procedures before illnesses progress to high-severity presentations. Although uncertainty and sample variance will persist at various levels, gaining of greater geographical information on severity of diseases than at present will increase the accuracy of GBD models.

Potential is clearly huge for exploration of this work in relation to development assistance for health,^{70,71} as the connection between health financing and outcomes needs better understanding than at present. Improved understanding will probably help identify the reasons why certain countries have such an impressive record, whereas others are so ineffective with the resources that they have available, at all levels of SDI. More precise measurement of health burden than at present, across countries and at the subnational level, could be tied to health services financing and delivery to identify systematic associations and causal relationships. Additionally, risk factor data, health outcome data, health financing data, and other socioeconomic indicators can be combined to measure and assess health system performance. Furthermore, a core focus moving forward with GBD is to progressively increase the spatial resolution at which we implement estimations to help realise aspirations in precision public health.⁷² A key goal of a geographically refined DALY, at 5×5 km spatial resolution—starting with some key CMNN causes and under-5 mortality—is part of a long-term aspiration.⁷³

Finally, upcoming GBD work will focus on exploration of future health scenarios, examining the likely burden of disease under different possible trajectories of independent drivers of health. This framework will capture the complex past trends and interdependent

relationships in socioeconomic development, risk factors, interventions, morbidity, mortality, and population to both estimate the likely future burden of disease and enable comparison between scenarios based on different sets of assumptions. For instance, it can be used to analyse the likely effect of the introduction and scale-up of a new type of vaccination or different trends in funding for ART for HIV/AIDS. Extension of this work through to DALY and HALE analyses is also part of these future plans.

To look at cause-deleted HALE would also be valuable so that we could understand the remaining DALY burden in the absence of diseases and injuries that might be highly amenable to preventive or curative measures. Compelling examples would include HIV, vaccine-preventable and malnutrition-related diseases, a subset of NCDs that are particularly amenable to health-care or preventive measures (eg, tobacco-related or alcohol-related or some congenital birth defects), and injuries (eg, firearms or transport injuries). This approach would not be fully counterfactual, but would be a first and more rigorous way of showing what conditions contributed to the changes in HALE from 1990 (or 2006) to 2016 than was possible in this study.

Limitations

Despite our continued methodological advancements and data enrichments, this study has limitations. First, all limitations documented in the elements of the GBD estimation process that allow for DALY and HALE estimation^{2-4,12} will contribute to uncertainty in these summary measures. Second, these summary measures will also be influenced by data availability. Time lags in the reporting of health information by national authorities and thus their subsequent incorporation into the GBD estimation mean that some of the most recent changes in health states will not be captured. Relatedly, data deficiencies from populations in conflict zones (eg, Syria, Iraq, Yemen, South Sudan, and Afghanistan), autonomous subnational regions, and certain non-geographically based subpopulations (ie, migrants, refugees, and some indigenous people) limit the precision of some of our estimated levels and trends of disease burden.⁷⁴ Third, the relationship between DALYs, HALE, and SDI, although explanatory, cannot be viewed as causal. Fourth, a non-trivial assumption of the analyses is the independence of the uncertainty calculated for YLLs and YLDs. Because of the link between death and prevalence, a positive correlation probably exists between these uncertainties that we do not capture. As such, we probably underestimate the aggregated uncertainty for DALYs, although the primary source of uncertainty for DALYs comes from uncertainty in disability weights, which are unaffected by this limitation. In future iterations of GBD, this potential correlation will be explored using copula, a statistical method that models the dependence structure among multiple independent

marginal probability distributions to estimate correlation between them.⁷⁵

Conclusion

Many improvements have been made to GBD 2016 to allow for a clearer and more nuanced picture of the changing picture of global health than in GBD 2015. Among these improvements are inclusion of new studies and subnational data (notably in India) and many substantial improvements to modelling and analyses. Prominent results of these changes include higher global DALY estimates for tuberculosis, HIV/AIDS, lower respiratory infection, and diarrhoeal disease than in GBD 2015. We have for the first time discussed our DALY exemplars, countries with the lowest ratio of observed to expected DALYs, as well as our poor performers, those with the largest ratio of observed to expected DALYs. These exemplars and poor performers suggest a need to learn lessons from those with clear health gains and implement system-based strategies for those struggling. This analysis and increasingly more detailed results will allow for more informed public policy and health financing decisions as the world faces an absolute expansion of morbidity.

Globally, individuals could expect to live substantially longer lives in 2016 than they could in 1990. This improvement was due to a rapid decline in YLLs and more modest age-standardised declines in YLDs, leading to lower age-standardised DALY rates across the entire socioeconomic development spectrum in 2016 than in 1990, with a decrease of approximately a third for all causes. At the same time, populations can expect to spend more time with functional health loss due to absolute morbidity expansion than previously. Such improvements have been accompanied by rapid population growth and ageing and, against the backdrop of the epidemiological transition, have resulted in the paradox of a total expansion in DALY burden and thus an ever-increasing demand on health systems, domestic health financing, development assistance for health, and associated global health organisations. This increasing demand on health systems is ubiquitous across time, location, GBD region, and SDI, posing a huge opportunity for health-care innovators in prevention and treatment for morbidity reduction. Our analysis of exemplars and poor performers is also indicative of various practices that can hasten or slow the epidemiological transition and warrant deep investigation.

GBD 2016 DALYs and HALE Collaborators

Simon I Hay, Amanuel Alemu Abajobir, Kalkidan Hassen Abate, Cristiana Abbafati, Kaja M Abbas, Foad Abd-Allah, Abdishakur M Abdulle, Teshome Abuka Abebo, Semaw Ferede Abera, Victor Aboyans, Laith J Abu-Raddad, Ilana N Ackerman, Isaac A Adedeji, Olatunji Adetokunboh, Ashkan Afshin, Rakesh Aggarwal, Sutapa Agrawal, Anurag Agrawal, Aliasghar Ahmad Kiadaliri, Muktar Beshir Ahmed, Amani Nidhal Aichour, Ibtihel Aichour, Miloud Taki Eddine Aichour, Sneha Aiyar, Tomi F Akinyemiju, Nadia Akseer, Faris Hasan Al Lami,

Fares Alahdab, Ziyad Al-Aly, Khurshid Alam, Noore Alam, Tahiya Alam, Deena Alasfoor, Kefyalew Addis Alene, Raghib Ali, Reza Alizadeh-Navaei, Juma M Alkaabi, Ala'a Alkerwi, François Alla, Peter Allebeck, Christine Allen, Fatma Al-Maskari, Mohammad AbdulAziz AlMazroa, Rajaa Al-Raddadi, Ubai Alsharif, Shirina Alsowaidi, Benjamin M Althouse, Khalid A Altirkawi, Nelson Alvis-Guzman, Azmeraw T Amare, Erfan Amini, Walid Ammar, Yaw Ampem Amoako, Mustafa Geleto Ansha, Carl Abelardo T Antonio, Palwasha Anwari, Johan Årnlöv, Megha Arora, Al Artaman, Krishna Kumar Aryal, Solomon W Asgedom, Tesfay Mehari Atey, Niguse Tadele Atnafu, Leticia Avila-Burgos, Euripide Frinel G Arthur Avokpaho, Ashish Awasthi, Shally Awasthi, Beatriz Paulina Ayala Quintanilla, Mahmoud Reza Azarpazhooh, Peter Azzopardi, Tesleem Kayode Babalola, Umar Bacha, Alaa Badawi, Kalpana Balakrishnan, Marlana S Bannick, Aleksandra Barac, Suzanne L Barker-Collo, Till Bärnighausen, Simon Barquera, Lope H Barrero, Sanjay Basu, Robert Battista, Katherine E Battle, Bernhard T Baune, Shahrzad Bazargan-Hejazi, Justin Beardsley, Neeraj Bedi, Yannick Béjot, Bayu Begashaw Bekele, Michelle L Bell, Derrick A Bennett, James R Bennett, Isabela M Bensenor, Jennifer Benson, Adugnaw Berhane, Derbew Fikadu Berhe, Eduardo Bernabé, Balem Demtsu Betsu, Mircea Beuran, Addisu Shunu Beyene, Anil Bhansali, Samir Bhatt, Zulfiqar A Bhutta, Sibhatu Biadgilign, Kelly Bienhoff, Boris Bikbov, Charles Birungi, Stan Biryukov, Donal Bisanzio, Habtamu Mellie Bizuayehu, Fiona M Blyth, Dube Jara Boneya, Dipan Bose, Ibrahim R Bou-Orm, Rupert R A Bourne, Michael Brainin, Carol E G Brayne, Alexandra Brazinova, Nicholas J K Breitborde, Paul S Briant, Gabrielle Britton, Traolach S Brugha, Rachelle Buchbinder, Lemma Negesa Bulto Bulto, Blair Bumgarner, Zahid A Butt, Lucero Cahuana-Hurtado, Ewan Cameron, Ismael Ricardo Campos-Nonato, Hélène Carabin, Rosario Cárdenas, David O Carpenter, Juan Jesus Carrero, Austin Carter, Felix Carvalho, Daniel Casey, Carlos A Castañeda-Orjuela, Jacqueline Castillo Rivas, Chris D Castle, Ferrán Catalá-López, Jung-Chen Chang, Fiona J Charlson, Pankaj Chaturvedi, Honglei Chen, Mirriam Chibalabala, Chioma Ezinne Chibueze, Vesper Hichilombwe Chisumpa, Abdulaal A Chitheer, Rajiv Chowdhury, Devasahayam Jesudas Christopher, Liliana G Ciobanu, Massimo Cirillo, Danny Colombara, Leslie Trumbull Cooper, Cyrus Cooper, Paolo Angelo Cortesi, Monica Cortinovic, Michael H Criqui, Elizabeth A Cromwell, Marita Cross, John A Crump, Abel Fekadu Dadi, Koustuv Dalal, Albertino Damasceno, Lalit Dandona, Rakhi Dandona, José das Neves, Dragos V Davitoiu, Kairat Davletov, Barbora de Courten, Diego De Leo, Hans De Steur, Louisa Degenhardt, Selina Deiparine, Robert P Dellavalle, Kebede Deribe, Amare Deribew, Don C Des Jarlais, Subhojit Dey, Samath D Dharmaratne, Preet K Dhillon, Daniel Dicker, Shirin Djalalinia, Huyen Phuc Do, Klara Dokova, David Teye Doku, E Ray Dorsey, Kadine Priscila Bender dos Santos, Tim R Driscoll, Manisha Dubey, Bruce Bartholow Duncan, Beth E Ebel, Michelle Echko, Ziad Ziad El-Khatib, Ahmadali Enayati, Aman Yesuf Endries, Sergey Petrovich Ermakov, Holly E Erskine, Setegn Eshetie, Babak Eshtrati, Alireza Esteghamati, Kara Estep, Fanuel Belayneh Bekele Fanuel, Tamer Farag, Carla Sofia e Sa Farinha, André Faro, Farshad Farzadfar, Mir Sohail Fazeli, Valery L Feigin, Andrea B Feigl, Seyed-Mohammad Fereshtehjad, João C Fernandes, Alize J Ferrari, Tesfaye Regassa Feyissa, Irina Filip, Florian Fischer, Christina Fitzmaurice, Abraham D Flaxman, Nataliya Foigt, Kyle J Foreman, Richard C Franklin, Joseph J Frostad, Nancy Fullman, Thomas Fürst, Joao M Furtado, Neal D Futran, Emmanuela Gakidou, Alberto L Garcia-Basteiro, Teshome Gebre, Gebremedhin Berhe Gebregergs, Tsegaye Tewelde Gebrehiwot, Johanna M Geleijnse, Ayele Geleto, Bikila Lencha Gemechu, Hailay Abrrha Gesesew, Peter W Gething, Alireza Ghajar, Katherine B Gibney, Richard F Gillum, Ibrahim Abdelmageem Mohamed Ginawi, Melkamu Dedefo Gishu, Giorgia Giussani, William W Godwin, Kashish Goel, Shifalika Goenka, Ellen M Goldberg, Philimon N Gona, Amador Goodridge, Sameer Vali Gopalani, Richard A Gosselin, Carolyn C Gotay, Atsushi Goto, Alessandra Carvalho Goulart, Nicholas Graetz, Harish Chander Gugnani, Rajeev Gupta, Prakash C Gupta, Tanush Gupta, Vipin Gupta, Rahul Gupta, Reyna A Gutiérrez, Vladimir Hachinski, Nima Hafezi-Nejad, Alemayehu Desalegne Hailu, Gessesew Bugssa Hailu, Randah Ribhi Hamadeh, Samer Hamidi, Mouhanad Hammami, Alexis J Handal, Graeme J Hankey, Yuantao Hao, Hilda L Harb, Habtamu Abera Hareri, Josep Maria Haro, Kimani M Harun, James Harvey, Mohammad Sadegh Hassanvand, Rasmus Havmoeller, Roderick J Hay, Mohammad T Hedayati, Delia Hendrie, Nathaniel J Henry, Ileana Beatriz Heredia-Pi, Pouria Heydarpour, Hans W Hoek, Howard J Hoffman, Masako Horino, Nobuyuki Horita, H Dean Hosgood, Sorin Hostiuc, Peter J Hotez, Damian G Hoy, Aung Soe Htet, Guoqing Hu, John J Huang, Chantal Huynh, Kim Moesgaard Iburg, Ehimario Uche Igumbor, Chad Ikeda, Caleb Mackay Salpeter Irvine, Kathryn H Jacobsen, Nader Jahanmehri, Mihajlo B Jakovljevic, Peter James, Simerjot K Jassal, Mehdi Javanbakht, Sudha P Jayaraman, Panniyammakal Jeemon, Paul N Jensen, Vivekanand Jha, Guohong Jiang, Denny John, Catherine O Johnson, Sarah Charlotte Johnson, Jost B Jonas, Mikk Jürisson, Zubair Kabir, Rajendra Kadel, Amaha Kahsay, Rital Kamal, Chittaranjan Kar, Nadim E Karam, André Karch, Corine Kakizi Karema, Seyed M Karimi, Chante Karimkhani, Amir Kasaiean, Getachew Mullu Kassa, Nicholas J Kassebaum, Nigussie Assefa Kassaw, Anshul Kastor, Srinivasa Vittal Katikireddi, Anil Kaul, Norito Kawakami, Peter Njenga Keiyoro, Laura Kemmer, Andre Pascal Kengne, Andre Keren, Chandrasekharan Nair Kesavachandran, Yousef Saleh Khader, Ibrahim A Khalil, Ejaz Ahmad Khan, Young-Ho Khang, Abdullah T Khoja, Ardeshir Khosravi, Jagdish Khubchandani, Christian Kielsing, Yun Jin Kim, Daniel Kim, Ruth W Kimokoti, Yohannes Kinfu, Adnan Kisa, Katarzyna A Kissimova-Skarbek, Niranjana Kissoon, Mika Kivimaki, Ann Kristin Knudsen, Yoshihiro Kokubo, Dhaval Kolte, Jacek A Kopec, Soewarta Kosen, Georgios A Kotsakis, Parvaiz A Koul, Ai Koyanagi, Michael Kravchenko, Kristopher J Krohn, Barthelemy Kuate Defo, Burcu Kucuk Bicer, G Anil Kumar, Pushpendra Kumar, Hmwe H Kyu, Anton Carl Jonas Lager, Dharmesh Kumar Lal, Ratilal Laloo, Tea Lallukka, Nkurunziza Lambert, Qing Lan, Van C Lansingh, Anders Larsson, Janet L Leasher, Paul H Lee, James Leigh, Cheru Tesema Leshargie, Janni Leung, Ricky Leung, Miriam Levi, Yichong Li, Yongmei Li, Xiaofeng Liang, Patrick Legesse Liben, Stephen S Lim, Shai Linn, Angela Liu, Mick Y Liu, Shiwei Liu, Yang Liu, Rakesh Lodha, Giancarlo Logroscino, Katharine J Looker, Alan D Lopez, Stefan Lorkowski, Paulo A Lotufo, Rafael Lozano, Timothy C D Lucas, Raimundas Lunevicius, Ronan A Lyons, Erlyn Rachelle King Macarayan, Emilie R Maddison, Hassan Magdy Abd El Razek, Mohammed Magdy Abd El Razek, Carlos Magis-Rodriguez, Mahdi Mahdavi, Marek Majdan, Reza Majdzadeh, Azeem Majeed, Reza Malekzadeh, Rajesh Malhotra, Deborah Carvalho Malta, Abdullah A Mamun, Helena Manguerra, Treh Manhertz, Lorenzo G Mantovani, Chabila C Mapoma, Lyn M March, Laurie B Marczak, Jose Martinez-Raga, Paulo Henrique Viegas Martins, Francisco Rogerlândio Martins-Melo, Ira Martopullo, Winfried März, Manu Raj Mathur, Mohsen Mazidi, Colm McAlinden, Madeline McCaughey, John J McGrath, Martin McKee, Suresh Mehata, Toni Meier, Kidanu Gebremariam Meles, Peter Memiah, Ziad A Memish, Walter Mendoza, Melkamu Merid Mengesha, Mubarek Abera Mengistie, Desalegn Tadesse Mengistu, George A Mensah, Atte Meretoja, Tuomo J Meretoja, Haftay Berhane Mezgebe, Renata Micha, Anousha Millear, Ted R Miller, Shawn Minnig, Mojde Mirarefin, Erkin M Mirrakhimov, Awoke Misganaw, Shiva Raj Mishra, Philip B Mitchell, Karzan Abdulmuhsin Mohammad, Alireza Mohammadi, Shafiu Mohammed, Kedir Endris Mohammed, Muktar Sano Kedir Mohammed, Murali B V Mohan, Ali H Mokdad, Sarah K Mollenkopf, Lorenzo Monasta, Julio Cesar Montañez Hernandez, Marcella Montico,

Maziar Moradi-Lakeh, Paula Moraga, Lidia Morawska, Rintaro Mori, Shane D Morrison, Mark Moses, Cliff Mountjoy-Venning, Kalayu Birhane Mruts, Ulrich O Mueller, Kate Muller, Michele E Mudoch, Srinivas Murthy, Gudlavalleti Venkata Satyanarayana Murthy, Kamarul Imran Musa, Jean B Nachega, Gabriele Nagel, Mohsen Naghavi, Aliya Naheed, Kovin S Naidoo, Vinay Nangia, Jamal T Nasher, Gopalakrishnan Natarajan, Dumessa Edessa Negasa, Ionut Negoii, Ruxandra Irina Negoii, Charles R Newton, Josephine Wanjiku Ngunjiri, Cuong Tat Nguyen, Quyen Le Nguyen, Grant Nguyen, Trang Huyen Nguyen, Minh Nguyen, Emma Nichols, Dina Nur Anggraini Ningrum, Vuong Minh Nong, Ole F Norheim, Bo Norrving, Jean Jacques N Noubiap, Alypio Nyandwi, Carla Makhlof Obermeyer, Martin J O'Donnell, Felix Akpojone Ogbo, In-Hwan Oh, Anselm Okoro, Olanrewaju Oladimeji, Andrew Toyin Olagunju, Tinuke Oluwasefunmi Olagunju, Helen E Olsen, Bekajoko Olubukunola Oluasanya, Jacob Olusegun Oluasanya, Kanyin Ong, John Nelson Opio, Eyal Oren, Alberto Ortiz, Richard H Osborne, Aaron Osgood-Zimmerman, Majdi Osman, Erika Ota, Mayowa O Owolabi, Mahesh PA, Rosana E Pacella, Basant Kumar Panda, Jeyaraj D Pandian, Christina Papachristou, Eun-Kee Park, Charles D Parry, Mahboubeh Parsaeian, Snehal T Patil, Scott B Patten, George C Patton, Deepak Paudel, Katherine Paulson, Neil Pearce, David M Pereira, Krystle Marie Perez, Norberto Perico, Konrad Pesudovs, Carrie Beth Peterson, William Arthur Petri, Max Petzold, Michael Robert Phillips, Geoffrey Phipps, David M Pigott, Julian David Pillay, Christine Pinho, Michael A Piradov, Dietrich Plass, Martin A Pletcher, Svetlana Popova, Richie G Poulton, Farshad Pourmalek, Dorairaj Prabhakaran, Narayan Prasad, Carrie Purcell, Manorama Purwar, Mostafa Qorbani, Rynaz H S Rabiee, Amir Radfar, Anwar Rafay, Kazem Rahimi, Afarin Rahimi-Movaghar, Vafa Rahimi-Movaghar, Mahfuzar Rahman, Muhammad Aziz Rahman, Mohammad Hifz Ur Rahman, Rajesh Kumar Rai, Sasa Rajsic, Usha Ram, Chhabi Lal Ranabhat, Thara Rangaswamy, Zane Rankin, Paturi Vishnupriya Rao, Puja C Rao, Salman Rawaf, Sarah E Ray, Robert C Reiner, Nikolas Reinig, Marissa Reitsma, Giuseppe Remuzzi, Andre M N Renzaho, Serge Resnikoff, Satar Rezaei, Antonio L Ribeiro, Hirbo Shore Roba, Stephen R Robinson, David Rojas-Rueda, Mohammad Bagher Rokni, Luca Ronfani, Gholamreza Roshandel, Gregory A Roth, Dietrich Rothenbacher, Ambuj Roy, Enrico Rubagotti, George Mugambage Ruhago, Soheil Saadat, Mahdi Safdarian, Saeid Safiri, Rajesh Sagar, Ramesh Sahathevan, Mohammad Ali Sahrain, Joseph Salama, Muhammad Muhammad Saleh, Joshua A Salomon, Sundeep Santosh Salvi, Abdallah M Samy, Juan Ramon Sanabria, Maria Dolores Sanchez-Niño, Damian Santomauro, João Vasco Santos, Itamar S Santos, Milena M Santric Milicevic, Benn Sartorius, Maheswar Satpathy, Monika Sawhney, Sonia Saxena, Kathryn Schelonka, Maria Inés Schmidt, Ione J C Schneider, Ben Schöttker, Aletta E Schutte, David C Schwebel, Falk Schwendicke, Soraya Seedat, Sadaf G Sepanlou, Edson E Servan-Mori, Amira Shaheen, Masood Ali Shaikh, Mansour Shamsipour, Sheikh Mohammed Shariful Islam, Rajesh Sharma, Jayendra Sharma, Jun She, Peilin Shi, Kenji Shibuya, Chloe Shields, Mekonnen Sisay Shiferaw, Mika Shigematsu, Rahman Shiri, Reza Shirkoobi, Shreya Shirude, Kawkab Shishani, Haitham Shoman, Soraya Siabani, Abla Mehio Sibai, Inga Dora Sigfusdottir, Donald H Silberberg, João Pedro Silva, Diego Augusto Santos Silva, Dayane Gabriele Alves Silveira, Jasvinder A Singh, Virendra Singh, Om Prakash Singh, Narinder Pal Singh, Dharendra Narain Sinha, Eirimi Skiadaresi, Vegard Skirbekk, Erica Leigh Slepak, David L Smith, Mari Smith, Badr H A Sobaih, Eugene Sobngwi, Michael Soljak, Reed J D Sorensen, Tatiane Cristina Moraes Sousa, Luciano A Sposato, Chandrashekhara T Sreeramareddy, Vinay Srinivasan, Jeffrey D Stanaway, Vasiliki Stathopoulou, Nicholas Steel, Dan J Stein, Caitlyn Steiner, Sabine Steink, Mark Andrew Stokes, Lars Jacob Stovner, Bryan Strub, Michelle Subart,

Muawiyah Babale Sufiyan, Rizwan Suliankatchi Abdulkader, Bruno F Sunguya, Patrick J Sur, Soumya Swaminathan, Bryan L Sykes, Dillon Sylte, Cassandra E I Szoeki, Rafael Tabarés-Seisdedos, Santosh Kumar Tadakamadla, Getachew Redae Taffere, Jukka S Takala, Nikhil Tandon, David Tanne, Yihunie L Tarekegn, Mohammad Tavakkoli, Nuno Taveira, Hugh R Taylor, Teketo Kassaw Tegegne, Arash Tehrani-Banihashemi, Tesfalidet Tekelab, Girma Temam Shifa, Abdullah Sulieyman Terkawi, Dawit Jember Tesfaye, Belay Tsessema, JS Thakur, Ornwipa Thamsuwan, Alice M Theadom, Andrew M Theis, Katie E Thomas, Nihal Thomas, Robert Thompson, Amanda G Thrift, Ruoyan Tobe-Gai, Myriam Tobollik, Marcello Tonelli, Roman Topor-Madry, Miguel Tortajada, Mathilde Touvier, Jefferson Traebert, Bach Xuan Tran, Christopher Troeger, Thomas Truelsen, Derrick Tsoi, Emin Murat Tuzcu, Hayley Tymeson, Stefanos Tyrovolas, Kingsley Nnanna Ukwaja, Eduardo A Undurraga, Chigozie Jesse Uneke, Rachel Updike, Olalekan A Uthman, Benjamin S Chudi Uzochukwu, Job F M van Boven, Santosh Varughese, Tommi Vasankari, Lennert J Veerman, S Venkatesh, Narayanaswamy Venketasubramanian, Ramesh Vidavalur, Lakshmi Vijayakumar, Francesco S Violante, Abhishek Vishnu, Sergey K Vladimirov, Vasilii Victorovich Vlassov, Stein Emil Vollset, Theo Vos, Fiseha Wadilo, Tolassa Wakayo, Mitchell T Wallin, Yuan-Pang Wang, Scott Weichenthal, Elisabete Weiderpass, Robert G Weintraub, Daniel J Weiss, Andrea Werdecker, Ronny Westerman, Harvey A Whiteford, Tissa Wijeratne, Hywel C Williams, Charles Shey Wiysonge, Belete Getahun Woldeyes, Charles D A Wolfe, Rachel Woodbrook, Anthony D Woolf, Abdulhalik Workicho, Denis Xavier, Gelin Xu, Simon Yadgir, Mohsen Yaghoubi, Bereket Yakob, Lijing L Yan, Yuichiro Yano, Pengpeng Ye, Mahari Gidey Yihdego, Hassen Hamid Yimam, Paul Yip, Naohiro Yonemoto, Seok-Jun Yoon, Marcel Yotebieng, Mustafa Z Younis, Chuanhua Yu, Zoubida Zaidi, Maysaa El Sayed Zaki, Elias Asfaw Zegeye, Zerihun Menkalew Zenebe, Xueying Zhang, Yingfeng Zheng, Maigeng Zhou, Ben Zipkin, Sanjay Zodpey, Leo Zoeckler, Liesl Joanna Zuhlke, Christopher J L Murray.

Affiliations

Institute for Health Metrics and Evaluation (Prof S I Hay DSc, A Afshin MD, S Aiyar, T Alam MPH, C Allen BA, M Arora BSA, M S Bannick BS, J R Bennett BA, K Bienhoff MA, S Biryukov BS, P S Briant BS, B Bumgarner MBA, A Carter BS, D Casey MPH, C D Castle BA, F J Charlson PhD, D Colombara MPH, E A Cromwell PhD, Prof L Dandona MD, Prof R Dandona PhD, Prof L Degenhardt PhD, S Deiparine, D Dicker BS, M Echno BSc, H E Erskine PhD, K Estep MPA, T Farag PhD, A J Ferrari PhD, C Fitzmaurice MD, A D Flaxman PhD, K J Foreman PhD, J J Frostad MPH, N Fullman MPH, E Gakidou PhD, W W Godwin BS, E M Goldberg BS, N Graetz MPH, J Harvey BS, N J Henry BS/BA, C Huynh BA, C Ikeda BS, C M S Irvine BA, C O Johnson PhD, S C Johnson MSc, N J Kassebaum MD, L Kemmer PhD, I A Khalil MD, K J Krohn BA, H H Kyu PhD, Prof S S Lim PhD, P Y Liu MPH, A Liu BS, E R Maddison BS, H Manguerra BS, T Manhertz BA, L B Marczak PhD, I Martopullo MPH, M McGaughey BA, A Millar BA, S Minnig MS, M Mirarefin MPH, A Misganaw PhD, A H Mokdad PhD, S K Mollenkopf, M Moses MHS, K Muller MPH, Prof M Naghavi PhD, C Mountjoy-Venning BA, G Nguyen MPH, M Nguyen BS, E Nichols BA, H E Olson MA, K Ong PhD, A Osgood-Zimmerman MS, K Paulson BS, G Phipps PhD, D M Pigott DPhil, C Pinho BA, M Pletcher BS, C Purcell BS, Z Rankin MPH, P C Rao MPH, S E Ray BS, R C Reiner PhD, N Reinig BS, M B Reitsma BS, G A Roth MD, J Salama MSc, D Santomauro PhD, K Schelonka BA, C Shields BS, S Shirude MPH, E L Slepak MLIS, D L Smith PhD, M Smith MPA, R J D Sorensen MPH, V Srinivasan BA, J D Stanaway PhD, C Steiner MPH, B Strub BS, M Subart BA, P J Sur BA, D Sylte BA, O Thamsuwan PhD, A M Theis BA, K E Thomas PhD, R Thompson PhD, C Troeger MPH, D Tsoi BS, H Tymeson BA,

- R Updike AB, Prof S E Vollset DrPH, T Vos PhD, Prof H A Whiteford PhD, R Woodbrook MLIS, S Yadgir BS, Prof M Zhou PhD, B Zipkin BA, L Zoeckler BA, Prof C J L Murray DPhil, Harborview Injury Prevention and Research Center (B E Ebel MD), Division of Hematology, Department of Medicine (C Fitzmaurice MD), Center for Health Trends and Forecasts, Institute for Health Metrics and Evaluation (Prof M B Jakovljevic PhD), School of Dentistry (G A Kotsakis DDS), University of Washington, Seattle, WA, USA (B M Althouse PhD, N D Futran MD, K M Harun MPH, P N Jensen PhD, J Leung PhD, S D Morrison MD, K M Perez MD); Oxford Big Data Institute, Li Ka Shing Centre for Health Information and Discovery (K E Battle DPhil, D Bisanzio PhD, E Cameron PhD, P W Gething PhD, Prof S I Hay DSc, T C D Lucas PhD, D J Weiss PhD), Nuffield Department of Population Health (D A Bennett PhD), Nuffield Department of Medicine (D Bisanzio PhD, A Deribew PhD), NIHR Musculoskeletal Biomedical Research Centre (Prof C Cooper MD), University of Oxford, Oxford, UK (R Ali MSc, Prof V Jha DM, K Rahimi DM); School of Public Health (A A Abajobir MPH, F J Charlson PhD, H E Erskine PhD, A J Ferrari PhD, J Leung PhD, D Santomauro PhD, L J Veerman PhD, Prof H A Whiteford PhD), School of Dentistry (Prof R Lalloo PhD), University of Queensland, Brisbane, QLD, Australia (S R Mishra MPH); Department of Epidemiology, College of Health Sciences (M B Ahmed MPH), Jimma University, Jimma, Ethiopia (K H Abate MS, T T Gebrehiwot MPH, H A Gesesew MPH, M A Mengistie MS, T Wakayo MS, A Workicho MPH); La Sapienza University, Rome, Italy (C Abbafati PhD); Virginia Tech, Blacksburg, VA, USA (Prof K M Abbas PhD); Department of Neurology, Cairo University, Cairo, Egypt (Prof F Abd-Allah MD); New York University Abu Dhabi, Abu Dhabi, United Arab Emirates (A M Abdulle PhD); Hawassa University College of Medicine and Health Sciences, Hawassa, Ethiopia (T A Abebo MPH, D J Tesfaye MPH); School of Public Health (S F Abera MSc, G B Gebregergs MPH, K G Meles MPH, G R Taffere PhD), School of Pharmacy (D F Berhe MS), College of Health Sciences (D T Mengistu MS, K E Mohammed MPH), Mekelle University, Mekelle, Ethiopia (S W Asgedom PhD, T M Atey MS, B D Betsu MS, G B Hailu MSc, A Kahsay MPH, H B Mezgebe MS, Z M Zenebe MS); Food Security and Institute for Biological Chemistry and Nutrition, University of Hohenheim, Stuttgart, Germany (S F Abera MSc); Dupuytren University Hospital, Limoges, France (Prof V Aboyans PhD); Infectious Disease Epidemiology Group, Weill Cornell Medical College in Qatar, Doha, Qatar (L J Abu-Raddad PhD); Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive Medicine (I N Ackerman PhD, Prof R Buchbinder PhD), Department of Medicine, School of Clinical Sciences at Monash Health (Prof A G Thrift PhD), Monash University, Melbourne, VIC, Australia (Prof B de Courten PhD); Department of Psychiatry (Prof C D Parry PhD), Stellenbosch University, Cape Town, South Africa (O Adetokunboh MD, Prof J B Nachege PhD, Prof S Seedat PhD, Prof C S Wiysonge PhD); Olabisi Onabanjo University, Ago-Iwoye, Nigeria (I A Adedji MS); Sanjay Gandhi Postgraduate Institute of Medical Sciences, Lucknow, India (Prof R Aggarwal MD); Centre for Control of Chronic Conditions (P Jeemon PhD), Public Health Foundation of India, Gurugram, India (S Agrawal PhD, Prof L Dandona MD, Prof R Dandona PhD, P K Dhillon PhD, S Goenka PhD, G A Kumar PhD, D K Lal MD, M R Mathur PhD, Prof S Zodpey PhD); CSIR - Institute of Genomics and Integrative Biology, Delhi, India (A Agrawal PhD); Department of Internal Medicine, Baylor College of Medicine, Houston, TX, USA (A Agrawal PhD); Department of Clinical Sciences Lund, Orthopedics, Clinical Epidemiology Unit (A Ahmad Kiadaliri PhD), Skane University Hospital, Department of Clinical Sciences Lund, Neurology (Prof B Norrving PhD), Lund University, Lund, Sweden; University Ferhat Abbas of Setif, Setif, Algeria (A N Aichour BS); National Institute of Nursing Education, Setif, Algeria (I Aichour MS); High National School of Veterinary Medicine, Algiers, Algeria (M T Aichour MD); Department of Epidemiology (T F Akinyemiju PhD), University of Alabama at Birmingham, Birmingham, AL, USA (D C Schwebel PhD, J A Singh MD); Centre for Global Child Health, The Hospital for Sick Children, Toronto, ON, Canada (N Akseer MSc, Z A Bhutta PhD); Dalla Lana School of Public Health (N Akseer MSc), Department of Nutritional Sciences, Faculty of Medicine (A Badawi PhD), Centre for Addiction and Mental Health (S Popova PhD), University of Toronto, Toronto, ON, Canada; Baghdad College of Medicine, Baghdad, Iraq (F H Al Lami PhD); Mayo Clinic Foundation for Medical Education and Research, Rochester, MN, USA (F Alahdab MD); Syrian American Medical Society, Washington, DC, USA (F Alahdab MD); Washington University in St Louis, St Louis, MO, USA (Z Al-Aly MD); Murdoch Childrens Research Institute (K Alam PhD, P Azzopardi PhD, Prof G C Patton MD, R G Weintraub MBBS), Department of Paediatrics (P Azzopardi PhD), Melbourne School of Population and Global Health (Prof A D Lopez PhD), Department of Medicine (A Meretoja PhD), Institute of Health and Ageing (Prof C E I Szoekie PhD), Center for Youth Mental Health (L Vijayakumar PhD), The University of Melbourne, Melbourne, VIC, Australia (K Alam PhD, M A Rahman PhD, Prof H R Taylor AC, Prof T Wijeratne MD); Sydney School of Public Health (Prof T R Driscoll PhD), The University of Sydney, Sydney, NSW, Australia (K Alam PhD, F M Blyth PhD, M Cross PhD, J Leigh PhD, Prof L M March PhD); Department of Health, Queensland, Brisbane, QLD, Australia (N Alam MAppEpid); Ministry of Health, Al Khuwair, Oman (D Alasfoor MSc); Department of Epidemiology and Biostatistics, Institute of Public Health (K A Alene MPH), College of Medical and Health Sciences (B B Bekele PhD), University of Gondar, Gondar, Ethiopia (Prof A F Dadi MPH, S Eshetie MS, B Tessema PhD); Department of Global Health, Research School of Population Health, Australian National University, Canberra, ACT, Australia (K A Alene MPH); Gastrointestinal Cancer Research Center (R Alizadeh-Navaei PhD), Department of Medical Mycology and Parasitology, School of Medicine (Prof M T Hedayati PhD), Mazandaran University of Medical Sciences, Sari, Iran; College of Medicine and Health Sciences, United Arab Emirates University, Al Ain, United Arab Emirates (J M Alkaabi MD, Prof F Al-Maskari PhD, S Alsowaidi MD); Luxembourg Institute of Health, Strassen, Luxembourg (A Alkerwi PhD); School of Public Health, University of Lorraine, Nancy, France (Prof F Alla PhD); Department of Public Health Sciences (P Allebeck PhD, Z Z El-Khatib PhD, R H S Rabiee MPH), Department of Neurobiology, Care Sciences and Society, Division of Family Medicine and Primary Care (Prof J Ärnlöv PhD, S Fereshtehnejad PhD), Department of Medical Epidemiology and Biostatistics (Prof J J Carrero PhD, E Weiderpass PhD), Karolinska Institutet, Stockholm, Sweden (R Havmoeller PhD, A C J Lager PhD); Saudi Ministry of Health, Riyadh, Saudi Arabia (M A AlMazroa MD, Prof Z A Memish MD); Joint Program of Family and Community Medicine, Jeddah, Saudi Arabia (R Al-Raddadi PhD); Charité Universitätsmedizin, Berlin, Germany (U Alsharif MPH); Institute for Disease Modeling, Bellevue, WA, USA (B M Althouse PhD); King Saud University, Riyadh, Saudi Arabia (K A Altirkawi MD, B H A Sobaih MD); Universidad de Cartagena, Cartagena de Indias, Colombia (Prof N Alvis-Guzman PhD); School of Medicine (A T Amare MPH, Prof B T Baune PhD, L G Ciobanu MS), Discipline of Psychiatry, School of Medicine (A T Olagunju MS), University of Adelaide, Adelaide, SA, Australia; College of Medicine and Health Sciences, Bahir Dar University, Bahir Dar, Ethiopia (A T Amare MPH); Uro-Oncology Research Center (E Amini MD), Endocrinology and Metabolism Research Center (Prof A Esteghamati MD, N Hafezi-Nejad MD, A Kasaean PhD), Non-Communicable Diseases Research Center (E Amini MD, F Farzadfar MD, A Khosravi PhD, M Parsaeian PhD), Department of Internal Medicine (A Ghajar MD), Center for Air Pollution Research, Institute for Environmental Research (M S Hassanvand PhD), Multiple Sclerosis Research Center, Neuroscience Institute (P Heydarpour MD, M A Sahraian MD), Hematology-Oncology and Stem Cell Transplantation Research Center (A Kasaean PhD), Institute of Health Policy and Management (M Mahdavi PhD), Knowledge Utilization Research Center and Community Based Participatory Research Center (Prof R Majdzadeh PhD), Digestive Diseases Research Institute (Prof R Malekzadeh MD, G Roshandel PhD, S G Sepanlou PhD), Department of Epidemiology and Biostatistics, School of Public Health (M Parsaeian PhD), Iranian National Center for Addiction Studies (INCAS) (A Rahimi-Movaghar MD), Sina Trauma and Surgery

Research Center (Prof V Rahimi-Movaghar MD, S Saadat PhD, M Safdarian MD), Institute for Environmental Research (M Shamsipour PhD), Cancer Research Center (Prof R Shirkoobi PhD), Tehran University of Medical Sciences, Tehran, Iran (Prof M B Rokni PhD); Ministry of Public Health, Beirut, Lebanon (W Ammar PhD, I R Bou-Orm MD, H L Harb MPH); Department of Medicine, Komfo Anokye Teaching Hospital, Kumasi, Ghana (Y A Amoako MD); West Hararghe Zonal Health Department, Chiro, Ethiopia (M G Ansha MPH); Department of Health Policy and Administration, College of Public Health, University of the Philippines Manila, Manila, Philippines (C A T Antonio MD); Self-employed, Kabul, Afghanistan (P Anwari MS); School of Health and Social Studies, Dalarna University, Falun, Sweden (Prof J Årnlöv PhD); University of Manitoba, Winnipeg, MB, Canada (A Artaman PhD); Nepal Health Research Council, Kathmandu, Nepal (K K Aryal MPH); University of Oslo, Oslo, Norway (K K Aryal MPH, A S Htet MPH); Department of Public Health, College of Health Sciences (M G Yihdego MPH), Mizan Tepi University, Mizan Teferi, Ethiopia (N T Atnafu MS, M S K Mohammed MS, H H Yimam MPH); PMA 2020, Addis Ababa, Ethiopia (N T Atnafu MS); National Institute of Public Health, Cuernavaca, Mexico (L Avila-Burgos PhD, S Barquera PhD, L Cahuana-Hurtado PhD, I R Campos-Nonato PhD, I B Heredia-Pi PhD, R Lozano PhD, J C Montañez Hernandez MSc, Prof E E Servan-Mori MSc); Institut de Recherche Clinique du Bénin (IRCB), Cotonou, Benin (E F G A Avokpaho MPH); Laboratoire d'Etudes et de Recherche-Action en Santé (LERAS Afrique), Parakou, Benin (E F G A Avokpaho MPH); Indian Institute of Public Health, Gandhinagar, India (A Awasthi PhD); King George's Medical University, Lucknow, India (S Awasthi MD); The Judith Lumley Centre for Mother, Infant and Family Health Research (B P Ayala Quintanilla PhD), La Trobe University, Melbourne, VIC, Australia (M A Rahman PhD); Peruvian National Institute of Health, Lima, Peru (B P Ayala Quintanilla PhD); Ghaem Hospital, Mashhad University of Medical Sciences, Mashhad, Iran (M R Azarpazhooh MD); Department of Clinical Neurological Sciences, London, ON, Canada (M R Azarpazhooh MD); Burnet Institute, Melbourne, VIC, Australia (P Azzopardi PhD); Wardliparingga Aboriginal Research Unit, South Australian Health and Medical Research Institute, Adelaide, South Australia, Australia (P Azzopardi PhD); Department of Community Health and Primary Care (T K Babalola MS), Department of Psychiatry, College of Medicine (A T Olagunju MS), University of Lagos, Lagos, Nigeria; School of Health Sciences, University of Management and Technology, Lahore, Pakistan (U Bacha PhD); Public Health Agency of Canada, Toronto, ON, Canada (A Badawi PhD); Department of Environmental Health Engineering, Sri Ramachandra University, Chennai, India (K Balakrishnan PhD); Institute of Social Medicine (M M Santric Milicevic PhD), Centre School of Public Health and Health Management (M M Santric Milicevic PhD), Faculty of Medicine (A Barac PhD), University of Belgrade, Belgrade, Serbia; School of Psychology, University of Auckland, Auckland, New Zealand (S L Barker-Collo PhD); Department of Global Health and Population (Prof T Barnighausen MD, A B Feigl ScD, J A Salomon PhD), Harvard T H Chan School of Public Health (I R Campos-Nonato PhD), Ariadne Labs (E R K Macarayan PhD), Harvard Medical School (M Osman MD), Harvard University, Boston, MA, USA; Africa Health Research Institute, Mtubatuba, South Africa (Prof T Barnighausen MD); Institute of Public Health, Heidelberg University, Heidelberg, Germany (Prof T Barnighausen MD, S Mohammed PhD); Department of Industrial Engineering, School of Engineering, Pontificia Universidad Javeriana, Bogota, Colombia (L H Barrero ScD); Stanford University, Stanford, CA, USA (S Basu PhD); Doctor Evidence, Santa Monica, CA, USA (R Battista MBA, M S Fazeli MD); College of Medicine, Charles R Drew University of Medicine and Science, Los Angeles, CA, USA (Prof S Bazargan-Hejazi PhD); David Geffen School of Medicine, University of California at Los Angeles, Los Angeles, CA, USA (Prof S Bazargan-Hejazi PhD); Oxford University, Ho Chi Minh City, Vietnam (J Beardsley MBChB); College of Public Health and Tropical Medicine, Jazan, Saudi Arabia (N Bedi MD); University Hospital and Medical School of Dijon, University of Burgundy, Dijon, France (Prof Y Béjot PhD); Department of Public Health, College of Health Sciences (M G Yihdego MPH), Mizan Tepi University, Mizan Aman, Ethiopia (B B Bekele PhD); Yale University, New Haven, CT, USA (Prof M L Bell PhD, J J Huang MD); Center for Clinical and Epidemiological Research Center, Hospital Universitario (A C Goulart PhD), Internal Medicine Department (Prof I S Santos PhD), University of São Paulo, São Paulo, Brazil (I M Bensenor PhD, Prof P A Lotufo DrPH); College of Health Sciences (A Berhane PhD), Public Health Department, College of Health (K B Mruts MPH), Debre Berhan University, Debre Berhan, Ethiopia; Department of Psychiatry (Prof H W Hoek MD), University Medical Center Groningen (D F Berhe MS), University of Groningen, Groningen, Netherlands (J F M van Boven PhD); Division of Health and Social Care Research (Prof C D Wolfe MD), King's College London, London, UK (E Bernabé PhD, Prof R J Hay DM); Carol Davila University of Medicine and Pharmacy, Bucharest, Romania (Prof M Beuran PhD, D V Davitoiu PhD, S Hostiu PhD, I Negoi PhD, R I Negoi PhD); Emergency Hospital of Bucharest, Bucharest, Romania (Prof M Beuran PhD, I Negoi PhD); College of Health and Medical Sciences (A S Beyene MPH, M M Mengesha MPH, H S Roba MPH), Haramaya University, Harar, Ethiopia (L N B Bulto MS, A Geleto MPH, M D Gishu MS, D E Negasa MS, M S Shiferaw MS); Postgraduate Institute of Medical Education and Research, Chandigarh, India (A Bhansali DM); Department of Infectious Disease Epidemiology (S Bhatt DPhil, T Fürst PhD), Department of Primary Care & Public Health (Prof A Majeed MD, M Soljak PhD), Imperial College London, London, UK (K J Foreman PhD, Prof S Rawaf MD, S Saxena MD, H Shoman MPH); Centre of Excellence in Women and Child Health, Aga Khan University, Karachi, Pakistan (Z A Bhutta PhD); Independent Public Health Consultants, Addis Ababa, Ethiopia (S Biadgilign MPH); IRCCS - Istituto di Ricerche Farmacologiche Mario Negri, Bergamo, Italy (B Bikbov MD, M Cortinovis Biotech D, N Perico MD, Prof G Remuzzi MD); Department of Epidemiology and Public Health (Prof M Kivimaki PhD), University College London, London, UK (C Birungi MS, M R Mathur PhD); Department of Public Health (D J Boneya MPH), Debre Markos University, Debre Markos, Ethiopia (H M Bizuayehu MPH, G M Kassa MS, C T Leshargie MPH, T K Tegegne MPH); World Bank, Washington, DC, USA (D Bose PhD); Vision & Eye Research Unit, Anglia Ruskin University, Cambridge, UK (Prof R R A Bourne MD); Danube-University Krems, Krems, Austria (Prof M Brainin PhD); Cambridge Institute of Public Health, Cambridge, UK (Prof C E G Bryane MD); Faculty of Health Sciences and Social Work, Department of Public Health, Faculty of Medicine, Comenius University, Bratislava, Slovakia (A Brazinova MD); The Ohio State University, Columbus, OH, USA (Prof N J K Breitborde PhD, M Yotebieng PhD); INDICASAT AIP, Panama, Panama (G Britton PhD); University of Leicester, Leicester, UK (Prof T S Brugha PhD); Monash Department of Clinical Epidemiology, Cabrini Institute, Melbourne, VIC, Australia (Prof R Buchbinder PhD); Al Shifa Trust Eye Hospital, Rawalpindi, Pakistan (Z A Butt PhD); Department of Biostatistics and Epidemiology, University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA (H Carabin PhD); Metropolitan Autonomous University, Mexico City, Mexico (R Cárdenas ScD); University at Albany, Rensselaer, NY, USA (Prof D O Carpenter MD); Faculty of Pharmacy (Prof F Carvalho PhD), i3S - Instituto de Investigação e Inovação em Saúde (J das Neves PhD), INEB - Instituto de Engenharia Biomédica (J das Neves PhD), Faculty of Medicine (J V Santos MD), UCIBIO@REQUIMTE, Toxicology Group, Faculty of Pharmacy (J P Silva PhD), University of Porto, Porto, Portugal; Colombian National Health Observatory, Instituto Nacional de Salud, Bogota, Colombia (C A Castañeda-Orjuela MSc); Epidemiology and Public Health Evaluation Group, Public Health Department, Universidad Nacional de Colombia, Bogota, Colombia (C A Castañeda-Orjuela MSc); Caja Costarricense de Seguro Social, San Jose, Costa Rica (Prof J Castillo Rivas MPH); Universidad de Costa Rica, San Pedro, Montes de Oca, Costa Rica (Prof J Castillo Rivas MPH); Department of Medicine, University of Valencia, INCLIVA Health Research Institute and CIBERSAM, Valencia, Spain (F Catalá-López PhD, Prof R Tabarés-Seisdedos PhD); Clinical Epidemiology Program, Ottawa Hospital Research Institute,

Ottawa, ON, Canada (F Catalá-López PhD); School of Nursing, College of Medicine, National Taiwan University, Taipei, Taiwan (Prof J Chang PhD); Queensland Centre for Mental Health Research, Brisbane, QLD, Australia (F J Charlson PhD, H E Erskine PhD, A J Ferrari PhD, D Santomauro PhD, Prof H A Whiteford PhD); Tata Memorial Hospital, Mumbai, India (Prof P Chaturvedi MD); Michigan State University, East Lansing, MI, USA (H Chen PhD); Crowd Watch Africa, Lusaka, Zambia (M Chibabalala BS); National Center for Child Health and Development, Setagaya ku, Japan (C E Chibueze PhD); University of Zambia, Lusaka, Zambia (V H Chisumpa MPhil, C C Mapoma PhD); University of Witwatersrand, Johannesburg, South Africa (V H Chisumpa MPhil); Ministry of Health, Baghdad, Iraq (A A Chitheer MD); Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK (R Chowdhury PhD); Christian Medical College, Vellore, India (Prof D J Christopher MD, Prof S Varughese DM); University of Salerno, Baronissi, Italy (Prof M Cirillo MD); Mayo Clinic, Jacksonville, FL, USA (L T Cooper MD); MRC Lifecourse Epidemiology Unit and NIHR Biomedical Research Centre, University of Southampton, Southampton, UK (Prof C Cooper MD); Research Centre on Public Health (CESP), University of Milan-Bicocca, Monza, Italy (P A Cortesi PhD); VA San Diego (S K Jassal MD), University of California, San Diego, La Jolla, CA, USA (M H Criqui MD); Centre for International Health, Dunedin School of Medicine (Prof J A Crump MD), University of Otago, Dunedin, New Zealand (Prof R G Poulton PhD); Flinders University, Adelaide, South Australia, Australia (Prof A F Dadi MPH, H A Gesesew MPH, Prof K Pesudovs PhD); Centre for Injury Prevention and Safety Promotion, School of Health and Medical Sciences, Orebro University, Orebro, Sweden (Prof K Dalal PhD); Faculty of Medicine, Eduardo Mondlane University, Maputo, Mozambique (Prof A Damasceno PhD); Republican Institute of Cardiology and Internal Diseases, Almaty, Kazakhstan (K Davletov PhD); School of Public Health, Kazakh National Medical University, Almaty, Kazakhstan (K Davletov PhD); Griffith University, Brisbane, QLD, Australia (Prof D De Leo DSc); Ghent University, Ghent, Belgium (H De Steur PhD, A Workicho MPH); National Drug and Alcohol Research Centre (Prof L Degenhardt PhD), Brien Holden Vision Institute (Prof S Resnikoff MD), School of Optometry and Vision Science (Prof S Resnikoff MD), University of New South Wales, Kensington, NSW, Australia (Prof P B Mitchell MD); University of Colorado School of Medicine and the Colorado School of Public Health, Aurora, CO, USA (R P Dellavalle MD); Brighton and Sussex Medical School, Brighton, UK (K Deribe MPH); School of Public Health (K Deribe MPH, A D Hailu MPH), Addis Ababa University, Addis Ababa, Ethiopia (H A Hareri MS, N A Kassaw MPH, G Temam Shifa MPH, B G Woldeyes MPH, M G Yihdego MPH); KEMRI-Wellcome Trust Research Programme, Kilifi, Kenya (A Deribew PhD); Mount Sinai Beth Israel, New York, NY, USA (Prof D C Des Jarlais PhD); Icahn School of Medicine at Mount Sinai, New York, NY, USA (Prof D C Des Jarlais PhD, A Vishnu PhD); Department of Community Medicine, Faculty of Medicine, University of Peradeniya, Peradeniya, Sri Lanka (S D Dharmaratne MD); Undersecretary for Research & Technology, Ministry of Health & Medical Education, Tehran, Iran (S Djalalinia PhD); Institute for Global Health Innovations, Duy Tan University, Da Nang, Vietnam (H P Do MSc, C T Nguyen MSc, Q L Nguyen MD, T H Nguyen MSc, V M Nong MSc); Department of Social Medicine, Faculty of Public Health, Medical University - Varna, Varna, Bulgaria (K Dokova PhD); University of Cape Coast, Cape Coast, Ghana (D T Doku PhD); University of Tampere, Tampere, Finland (D T Doku PhD); University of Rochester Medical Center, Rochester, NY, USA (E R Dorsey MD); Universidade do Estado de Santa Catarina, Florianópolis, Brazil (Prof K P B dos Santos MA); International Institute for Population Sciences, Mumbai, India (M Dubey MPhil, A Kastor MPhil, P Kumar MPhil, B K Panda MPhil, M H U Rahman MPhil, Prof U Ram PhD); Federal University of Rio Grande do Sul, Porto Alegre, Brazil (B B Duncan PhD, C Kieling MD, Prof M I Schmidt MD); University of North Carolina, Chapel Hill, NC, USA (B B Duncan PhD); Department of Global Health and Social Medicine, Harvard Medical School, Kigali, Rwanda (Z Z El-Khatib PhD); School of Public Health and Health Sciences Research Center, Sari, Iran (Prof A Enayati PhD); Arba Minch University, Arba Minch, Ethiopia (A Y Endries MPH, G Temam Shifa MPH); The Institute of Social and Economic Studies of Population, Russian Academy of Sciences, Moscow, Russia (Prof S P Ermakov DSc); Federal Research Institute for Health Organization and Informatics, Ministry of Health of the Russian Federation, Moscow, Russia (Prof S P Ermakov DSc); Ministry of Health and Medical Education, Tehran, Iran (B Eshrati PhD); Arak University of Medical Sciences, Arak, Iran (B Eshrati PhD); Hawassa University, Hawassa, Ethiopia (F B B Fanuel MPH); Wolaita Sodo University, Wolaita Sodo, Ethiopia (F B B Fanuel MPH, F Wadilo MS); DGS Directorate General of Health, Lisboa, Portugal (C S E S Farinha MSc); Universidade Aberta, Lisboa, Portugal (C S E S Farinha MSc); Federal University of Sergipe, Aracaju, Brazil (Prof A Faro PhD); National Institute for Stroke and Applied Neurosciences (V L Feigin PhD), Auckland University of Technology, Auckland, New Zealand (A M Theadom PhD); CBQF - Center for Biotechnology and Fine Chemistry - Associate Laboratory, Faculty of Biotechnology, Catholic University of Portugal, Porto, Portugal (J C Fernandes PhD); Wollega University, Nekemte, Ethiopia (T R Feyissa MPH, T Tekelab MS); Kaiser Permanente, Fontana, CA, USA (I Filip MD); School of Public Health, Bielefeld University, Bielefeld, Germany (F Fischer PhD); Fred Hutchinson Cancer Research Center, Seattle, WA, USA (C Fitzmaurice MD); Institute of Gerontology, Academy of Medical Science, Kyiv, Ukraine (N Foigt PhD); James Cook University, Townsville, QLD, Australia (R C Franklin PhD); Department of Epidemiology and Public Health (T Fürst PhD), Swiss Tropical and Public Health Institute, Basel, Switzerland (C K Karema MSc); University of Basel, Basel, Switzerland (T Fürst PhD); Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, Brazil (J M Furtado MD); Manhiça Health Research Center, Manhiça, Mozambique (A L Garcia-Basteiro MSc); Barcelona Institute for Global Health, Barcelona, Spain (A L Garcia-Basteiro MSc); The Task Force for Global Health, Decatur, GA, USA (T Gebre PhD); Division of Human Nutrition, Wageningen University, Wageningen, Netherlands (J M Geleijnse PhD); University of Newcastle, Newcastle, NSW, Australia (A Geleto MPH, T Tekelab MS); Mada Walabu University, Bale Goba, Ethiopia (B L Gemechu MPH); The Peter Doherty Institute for Infection and Immunity, The University of Melbourne & The Royal Melbourne Hospital, Melbourne, VIC, Australia (K B Gibney MBBS); Howard University, Washington, DC, USA (R F Gillum MD); College of Medicine, University of Hail, Hail, Saudi Arabia (I A Ginawi MD); Kersa Health and Demographic Surveillance System, Harar, Ethiopia (M D Gishu MS); IRCCS - Istituto di Ricerche Farmacologiche Mario Negri, Milan, Italy (G Giussani BiolD); Mayo Clinic, Rochester, MN, USA (K Goel MD); University of Massachusetts Boston, Boston, MA, USA (Prof P N Gona PhD); Instituto de Investigaciones Científicas y Servicios de Alta Tecnología - INDICASAT-AIP, Ciudad del Saber, Panama (A Goodridge PhD); Department of Health and Social Affairs, Government of the Federated States of Micronesia, Palikir, Federated States of Micronesia (S V Gopalani MPH); University of California, San Francisco, San Francisco, CA, USA (R A Gosselin MD); University of British Columbia, Vancouver, BC, Canada (C C Gotay PhD, Prof N Kisson MD, J A Kopec PhD, S Murthy MD, F Pourmalek PhD); Division of Epidemiology, Center for Public Health Sciences, National Cancer Center, Tokyo, Japan (A Goto PhD); Center of Check of Hospital Sirio Libanes, São Paulo, Brazil (A C Goulart PhD); Departments of Microbiology and Epidemiology & Biostatistics, Saint James School of Medicine, The Quarter, Anguilla (Prof H C Gugnani PhD); Eternal Heart Care Centre and Research Institute, Jaipur, India (R Gupta PhD); Healis - Sekhsaria Institute for Public Health, Navi Mumbai, India (P C Gupta DSc); Montefiore Medical Center, Bronx, NY, USA (T Gupta MD); Albert Einstein College of Medicine, Bronx, NY, USA (T Gupta MD, Prof H D Hosgood PhD); Department of Anthropology, University of Delhi, Delhi, India (V Gupta PhD); West Virginia Bureau for Public Health, Charleston, WV, USA (R Gupta MD); National Institute of Psychiatry Ramon de la Fuente, Mexico City, Mexico (R A Gutiérrez PhD); Department of Clinical Neurological Sciences (L A Sposato MD), Western University,

London, ON, Canada (Prof V Hachinski DSc, T O Olagunju MD); Department of Psychosocial Science (A K Knudsen PhD), Department of Global Public Health and Primary Care (Prof S E Vollset DrPH), University of Bergen, Bergen, Norway (A D Hailu MPH, Prof O F Norheim PhD); Kilde Awlaelo Health and Demographic Surveillance System, Mekelle, Ethiopia (G B Hailu MSc); Arabian Gulf University, Manama, Bahrain (Prof R R Hamadeh DPhil); Hamdan Bin Mohammed Smart University, Dubai, United Arab Emirates (S Hamidi DrPH); Wayne County Department of Health and Human Services, Detroit, MI, USA (M Hammami MD); University of New Mexico, Albuquerque, NM, USA (A J Handal PhD); School of Medicine and Pharmacology, University of Western Australia, Perth, WA, Australia (Prof G J Hankey MD); Harry Perkins Institute of Medical Research, Nedlands, WA, Australia (Prof G J Hankey MD); Western Australian Neuroscience Research Institute, Nedlands, WA, Australia (Prof G J Hankey MD); School of Public Health (Prof Y Hao PhD), Zhongshan Ophthalmic Center (Y Zheng MD), Sun Yat-sen University, Guangzhou, China; Parc Sanitari Sant Joan de Déu - CIBERSAM, Sant Boi de Llobregat (Barcelona), Spain (J M Haro MD); Kenyatta University, Nairobi, Kenya (K M Harun MPH); International Foundation for Dermatology, London, UK (Prof R J Hay DM); School of Public Health (D Hendrie PhD), Centre for Population Health (T R Miller PhD), Curtin University, Perth, WA, Australia; Department of Epidemiology, Mailman School of Public Health (Prof H W Hoek MD), Columbia University, New York, NY, USA (Prof V Skirbekk PhD); Epidemiology and Statistics Program, National Institute on Deafness and Other Communication Disorders (H J Hoffman MA), Center for Translation Research and Implementation Science, National Heart, Lung, and Blood Institute (G A Mensah MD), National Institutes of Health, Bethesda, MD, USA; Bureau of Child, Family & Community Wellness, Nevada Division of Public and Behavioral Health, Carson City, NV, USA (M Horino MPH); Department of Pulmonology, Yokohama City University Graduate School of Medicine, Yokohama, Japan (N Horita MD); College of Medicine, Baylor University, Houston, Texas, USA (P J Hotez PhD); Public Health Division, The Pacific Community, Noumea, New Caledonia (D G Hoy PhD); International Relations Division, Ministry of Health, Nay Pyi Taw, Myanmar (A S Htet MPhil); Department of Epidemiology and Health Statistics, School of Public Health, Central South University, Changsha, China (G Hu PhD); National Centre for Register-Based Research, Aarhus School of Business and Social Sciences (Prof J J McGrath PhD), Aarhus University, Aarhus, Denmark (K M Iburg PhD); US Centers for Disease Control and Prevention, Pretoria, South Africa (Prof E U Igumbor PhD); School of Public Health, University of the Western Cape, Cape Town, South Africa (Prof E U Igumbor PhD); Department of Global and Community Health, George Mason University, Fairfax, VA, USA (K H Jacobsen PhD); School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran (N Jahanmehr PhD); Faculty of Medical Sciences, University of Kragujevac, Kragujevac, Serbia (Prof M B Jakovljevic PhD); Department of Epidemiology, Harvard T H Chan School of Public Health, Channing Division of Network Medicine, Brigham & Women's Hospital, Harvard Medical School, Boston, MA, USA (P James ScD); University of Aberdeen, Aberdeen, UK (M Javanbakht PhD); Department of Surgery, Virginia Commonwealth University, Richmond, VA, USA (S P Jayaraman MD); Centre for Chronic Disease Control, New Delhi, India (P Jeemon PhD, D Prabhakaran DM); The George Institute for Global Health, New Delhi, India (Prof V Jha DM); Tianjin Centers for Disease Control and Prevention, Tianjin, China (G Jiang MD); International Center for Research on Women, New Delhi, India (D John MPH); Department of Ophthalmology, Medical Faculty Mannheim, Ruprecht-Karls-University Heidelberg, Mannheim, Germany (Prof J B Jonas MD); Institute of Family Medicine and Public Health, University of Tartu, Tartu, Estonia (M Jürisson MD); University College Cork, Cork, Ireland (Z Kabir PhD); London School of Economics and Political Science, London, UK (R Kadel MPH); CSIR - Indian Institute of Toxicology Research, Lucknow, India (R Kamal MSc, C N Kesavachandran PhD); SCB Medical College, Cuttack, India (Prof C Kar DM); University of Balamand, Beirut, Lebanon (N E Karam MD); Epidemiological and Statistical Methods Research Group, Helmholtz Centre for Infection Research, Braunschweig, Germany (A Karch MD); Hannover-Braunschweig Site, German Center for Infection Research, Braunschweig, Germany (A Karch MD); Quality and Equity Health Care, Kigali, Rwanda (C K Karema MSc); University of Washington Tacoma, Tacoma, WA, USA (S M Karimi PhD); Case Western University Hospitals, Cleveland, OH, USA (C Karimkhani MD); MRC/CSO Social & Public Health Sciences Unit, University of Glasgow, Glasgow, UK (S V Katikireddi PhD); Oklahoma State University, Tulsa, OK, USA (A Kaul MD); School of Public Health (Prof N Kawakami MD), University of Tokyo, Tokyo, Japan (K Shibuya MD); Institute of Tropical and Infectious Diseases, Nairobi, Kenya (P N Keiyoro PhD); School of Continuing and Distance Education, Nairobi, Kenya (P N Keiyoro PhD); UKZN Gastrointestinal Cancer Research Centre (Prof B Sartorius PhD), South African Medical Research Council, Cape Town, South Africa (A P Kengne PhD); Department of Psychiatry (Prof D J Stein PhD), University of Cape Town, Cape Town, South Africa (A P Kengne PhD, J J N Noubiap MD); Assuta Hospitals, Assuta Hashalom, Tel Aviv, Israel (Prof A Keren MD); Department of Community Medicine, Public Health and Family Medicine, Jordan University of Science and Technology, Irbid, Jordan (Prof Y S Khader ScD); Health Services Academy, Islamabad, Pakistan (E A Khan MD); Department of Health Policy and Management, Seoul National University College of Medicine, Seoul, South Korea (Prof Y Khang MD); Institute of Health Policy and Management, Seoul National University Medical Center, Seoul, South Korea (Prof Y Khang MD); Iranian Ministry of Health and Medical Education, Tehran, Iran (A Khosravi PhD); Mohammed Ibn Saudi University, Riyadh, Saudi Arabia (A T Khoja MD); Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA (A T Kjoja MD, Prof J B Nachega PhD); Department of Nutrition and Health Science, Ball State University, Muncie, IN, USA (J Khubchandani PhD); Hospital de Clinicas de Porto Alegre, Porto Alegre, Brazil (C Kielsing MD); School of Medicine, Xiamen University Malaysia Campus, Sepang, Malaysia (Y J Kim PhD); Department of Health Sciences, Northeastern University, Boston, MA, USA (Prof D Kim DrPH); Simmons College, Boston, MA, USA (R W Kimokoti MD); Centre for Research and Action in Public Health, Canberra, ACT, Australia (Y Kinfu); Oslo University, Oslo, Norway (Prof A Kisa PhD); Institute of Public Health, Faculty of Health Sciences (R Topor-Madry PhD), Jagiellonian University Medical College, Krakow, Poland (K A Kissimova-Skarbek PhD); Clinicum, Faculty of Medicine (Prof M Kivimaki PhD), Finnish Institute of Occupational Health, Work Organizations, Work Disability Program, Department of Public Health, Faculty of Medicine (T Lallukka PhD, R Shiri PhD), University of Helsinki, Helsinki, Finland (T J Meretoja PhD); Center for Disease Burden (A K Knudsen PhD, Prof S E Vollset DrPH), Norwegian Institute of Public Health, Oslo, Norway (Prof V Skirbekk PhD); Department of Preventive Cardiology, National Cerebral and Cardiovascular Center, Suita, Japan (Y Kokubo PhD); Division of Cardiology, Brown University, Providence, RI, USA (D Kolte MD); Center for Community Empowerment, Health Policy and Humanities, National Institute of Health Research & Development, Jakarta, Indonesia (S Kosen MD); Sher-i-Kashmir Institute of Medical Sciences, Srinagar, India (Prof P A Koul MD); Research and Development Unit, Parc Sanitari Sant Joan de Deu (CIBERSAM), Barcelona, Spain (A Koyanagi MD); Research Center of Neurology, Moscow, Russia (M Kravchenko PhD, Prof M A Piradov DSc); Department of Social and Preventive Medicine, School of Public Health (Prof B Kuate Defo PhD), Department of Demography and Public Health Research Institute (Prof B Kuate Defo PhD), University of Montreal, Montreal, QC, Canada; Institute of Public Health, Hacettepe University, Ankara, Turkey (B Kucuk Bicer PhD); Centre for Epidemiology and Community Medicine, Stockholm County Council, Solna, Sweden (A C J Lager PhD); Ministry of Public Health and Fight Against AIDS, Mukaza, Burundi (N Lambert MD); National Cancer Institute, Rockville, MD, USA (Q Lan PhD); Help Me See, Inc, New York, NY, USA (V C Lansingh PhD); Instituto Mexicano de Oftalmologia, Queretaro, Mexico (V C Lansingh PhD); Department of Medical Sciences, Uppsala University, Uppsala, Sweden (Prof A Larsson PhD); College of Optometry, Nova Southeastern University, Fort Lauderdale,

FL, USA (J L Leasher OD); Hong Kong Polytechnic University, Hong Kong, China (P H Lee PhD); State University of New York, Albany, Rensselaer, NY, USA (R Leung PhD); Tuscany Regional Centre for Occupational Injuries and Diseases, Florence, Italy (M Levi PhD); Department of Data Management, Peking University Clinical Research Institute, Beijing, China (Y Li PhD); National Center for Chronic and Noncommunicable Disease Control and Prevention (Y Li PhD, S Liu PhD, P Ye MPH, Prof M Zhou PhD), Chinese Center for Disease Control and Prevention, Beijing, China (Prof X Liang MD); San Francisco VA Medical Center, San Francisco, CA, USA (Y Li PhD); Samara University, Samara, Ethiopia (M L Liben MPH); University of Haifa, Haifa, Israel (Prof S Linn MD); Emory University, Atlanta, GA, USA (Prof Y Liu PhD, Prof M R Phillips MD); All India Institute of Medical Sciences, New Delhi, India (R Lodha MD, Prof R Malhotra MS, A Roy DM, R Sagar MD, Prof N Tandon PhD); University of Bari, Bari, Italy (Prof G Logroscino PhD); University of Bristol, Bristol, UK (K J Looker PhD); Institute of Nutrition, Friedrich Schiller University Jena, Jena, Germany (Prof S Lorkowski PhD); Competence Cluster for Nutrition and Cardiovascular Health (nutriCARD) Halle-Jena-Leipzig, Jena, Germany (Prof S Lorkowski PhD); Aintree University Hospital National Health Service Foundation Trust, Liverpool, UK (Prof R Lunevicius PhD); School of Medicine, University of Liverpool, Liverpool, UK (Prof R Lunevicius PhD); Farr Institute, Swansea University, Swansea, UK (Prof R A Lyons MD); Ateneo de Manila University, Manila, Philippines (E R K Macarayan PhD); Mansoura Faculty of Medicine, Mansoura, Egypt (H Magdy Abd El Razek MBBCh); Aswan University Hospital, Aswan Faculty of Medicine, Aswan, Egypt (M Magdy Abd El Razek MBBCh); National Center for the Prevention and Control of HIV/AIDS, Secretariat of Health, Mexico City, Mexico (C Magis-Rodriguez PhD); Erasmus University Rotterdam, Rotterdam, Netherlands (M Mahdavi PhD); Faculty of Health Sciences and Social Work, Department of Public Health, Trnava University, Trnava, Slovakia (M Majdan PhD); National Institute of Health Research, Tehran, Iran (Prof R Majdzadeh PhD); Universidade Federal de Minas Gerais, Belo Horizonte, Brazil (Prof D C Malta PhD, P H V Martins PhD); The University of Queensland, Brisbane, QLD, Australia (Prof A A Mamun PhD); University of Milano Bicocca, Monza, Italy (Prof L G Mantovani DSc); Hospital Universitario Doctor Peset, Valencia, Spain (J Martinez-Raga PhD, M Tortajada PhD); CEU Cardinal Herrera University, Moncada, Spain (J Martinez-Raga PhD); Federal Institute of Education, Science and Technology of Ceará, Caucaia, Brazil (F R Martins-Melo PhD); Synlab Academy, Mannheim, Germany (Prof W März MD); Clinical Institute of Medical and Chemical Laboratory Diagnostics, Medical University of Graz, Graz, Austria (Prof W März MD); Key State Laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China (M Mazidi PhD); University Hospitals Bristol NHS Foundation Trust, Bristol, UK (C McAlinden PhD); Public Health Wales, Swansea, UK (C McAlinden PhD); Queensland Centre for Mental Health Research, The Park Centre for Mental Health, Wacol, QLD, Australia (Prof J J McGrath PhD); Queensland Brain Institute (Prof J J McGrath PhD), University of Queensland, Brisbane, QLD, Australia (S R Mishra MPH); London School of Hygiene and Tropical Medicine, London, UK (Prof M McKee DSc, Prof G V S Murthy MD, Prof N Pearce PhD); Ipas Nepal, Kathmandu, Nepal (S Mehata PhD); Martin Luther University Halle-Wittenberg, Halle (Saale), Germany (T Meier PhD); University of West Florida, Pensacola, FL, USA (P Memiah PhD); College of Medicine, Alfaisal University, Riyadh, Saudi Arabia (Prof Z A Memish MD); United Nations Population Fund, Lima, Peru (W Mendoza MD); Department of Neurology (A Meretoja PhD), Comprehensive Cancer Center, Breast Surgery Unit (T J Meretoja PhD), Helsinki University Hospital, Helsinki, Finland; Friedman School of Nutrition Science and Policy (R Micha PhD), Tufts University, Boston, MA, USA (P Shi PhD); Pacific Institute for Research & Evaluation, Calverton, MD, USA (T R Miller PhD); Hunger Action Los Angeles, Los Angeles, CA, USA (M Mirarefin MPH); Kyrgyz State Medical Academy, Bishkek, Kyrgyzstan (Prof E M Mirrakhimov PhD); National Center of Cardiology and Internal Disease, Bishkek, Kyrgyzstan (Prof E M Mirrakhimov PhD); Nepal Development Society, Chitwan, Nepal (S R Mishra MPH); University of Salahaddin, Erbil, Iraq (K A Mohammad PhD); ISHIK University, Erbil, Iraq (K A Mohammad PhD); Neuroscience Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran (A Mohammadi PhD); Health Systems and Policy Research Unit (S Mohammed PhD), Ahmadu Bello University, Zaria, Nigeria (M B Sufiyan MBA); Narayana Health, Bangalore, India (Prof M B V Mohan MD); Institute for Maternal and Child Health, IRCCS "Burlo Garofolo", Trieste, Italy (L Monasta DSc, M Montico MSc, L Ronfani PhD); Department of Community Medicine, Preventive Medicine and Public Health Research Center (M Moradi-Lakeh MD, A Tehrani-Banihashemi PhD), Gastrointestinal and Liver Disease Research Center (GILDR; M Moradi-Lakeh MD), Iran University of Medical Sciences, Tehran, Iran (M Yaghoubi MSc); Lancaster Medical School, Lancaster University, Lancaster, UK (P Moraga PhD); International Laboratory for Air Quality and Health (L Morawska PhD), Institute of Health and Biomedical Innovation (R E Pacella PhD), Queensland University of Technology, Brisbane, QLD, Australia; National Center for Child Health and Development, Setagaya, Japan (R Mori PhD, R Tobe-Gai PhD); Competence Center Mortality-Follow-Up of the German National Cohort (A Werdecker PhD), Federal Institute for Population Research, Wiesbaden, Germany (Prof U O Mueller PhD, R Westerman PhD); Indian Institute of Public Health, Public Health Foundation of India, Hyderabad, India (Prof G V S Murthy MD); School of Medical Sciences, University of Science Malaysia, Kubang Kerian, Malaysia (K I Musa MD); Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA, USA (Prof J B Nachega PhD); International Centre for Diarrhoeal Disease Research, Bangladesh (icddr), Dhaka, Bangladesh (A Naheed PhD, S M Shariful Islam PhD); Public Health Medicine, School of Nursing and Public Health (Prof B Sartorius PhD), Discipline of Public Health Medicine, School of Nursing and Public Health (B Yakob PhD), University of KwaZulu-Natal, Durban, South Africa (Prof K S Naidoo PhD, E A Zegeye MS); Suraj Eye Institute, Nagpur, India (V Nangia MD); Ministry of Public Health and Population, Sana'a, Yemen (J T Nasher MSc); Madras Wellcome College, Chennai, India, India (Prof G Natarajan DM); KEMRI Wellcome Trust, Kilifi, Kenya (Prof C R Newton MD); University of Nairobi, Nairobi, Kenya (J W Ngunjiri PhD); Department of Public Health, Semarang State University, Semarang City, Indonesia (D N A Ningrum MPH); Graduate Institute of Biomedical Informatics, College of Medical Science and Technology, Taipei Medical University, Taipei City, Taiwan (D N A Ningrum MPH); Medical Diagnostic Centre, Yaounde, Cameroon (J J N Noubiap MD); Rwanda Ministry of Health, Kigali, Rwanda (A Nyandwi MPH); American University of Beirut, Beirut, Lebanon (Prof C M Obermeyer DSc, Prof A M Sibai PhD); National University of Ireland Galway, Galway, Ireland (M J O'Donnell PhD); Centre for Health Research (F A Ogo MPH), Western Sydney University, Penrith, NSW, Australia (Prof A M N Ranzhoo PhD); Department of Preventive Medicine, School of Medicine, Kyung Hee University, Seoul, South Korea (Prof I Oh PhD); Society for Family Health, Abuja, Nigeria (A Okoro MPH); Human Sciences Research Council (HSRC), South Africa and University of KwaZulu-Natal, Durban, South Africa (O Oladimeji MS); Department of Psychiatry, Lagos University Teaching Hospital, Lagos, Nigeria (A T Olagunju MS); McMaster University, Hamilton, ON, Canada (A T Olagunju MD); Center for Healthy Start Initiative, Lagos, Nigeria (B O Olusanya PhD, J O Olusanya MBA); Lira District Local Government, Lira Municipal Council, Uganda (J N Opio MPH); University of Arizona, Tucson, AZ, USA (Prof E Oren PhD); IIS-Fundacion Jimenez Diaz-UAM, Madrid, Spain (Prof A Ortiz PhD); Deakin University, Geelong, VIC, Australia (Prof R H Osborne PhD); YBank, Cambridge, MA, USA (M Osman MD); St Luke's International University, Tokyo, Japan (E Ota PhD); Department of Medicine, Ibadan, Nigeria (M O Owolabi Dr Med); Blossom Specialist Medical Center, Ibadan, Nigeria (M O Owolabi Dr Med); JSS Medical College, JSS University, Mysore, India (Prof M PA DNB) Christian Medical College Ludhiana, Ludhiana, India (J D Pandian DM); Charité University Medicine Berlin, Berlin, Germany (C Papachristou PhD); Department of Medical Humanities and Social Medicine, College of Medicine, Kosin University, Busan, South Korea (E Park PhD); Alcohol,

Tobacco & Other Drug Research Unit (Prof C D Parry PhD), South African Medical Research Council, Cape Town, South Africa (A P Kengne PhD); Krishan Institute of Medical Sciences, Deemed University, School of Dental Sciences, Karad, India (S T Patil MDS); Department of Community Health Sciences (Prof S B Patten PhD), University of Calgary, Calgary, AB, Canada (Prof M Tonelli MD); UK Department for International Development, Lalitpur, Nepal (D Paudel PhD); REQUIMTE/LAQV, Laboratório de Farmacognosia, Departamento de Química, Faculdade de Farmácia, Universidade do Porto, Porto, Portugal (Prof D M Pereira PhD); Aalborg University, Aalborg Esst, Denmark (C B Peterson PhD); Department of Anesthesiology (A S Terkawi MD), University of Virginia, Charlottesville, VA, USA (W A Petri MD); Health Metrics Unit, University of Gothenburg, Gothenburg, Sweden (Prof M Petzold PhD); University of the Witwatersrand, Johannesburg, South Africa (Prof M Petzold PhD); Shanghai Jiao Tong University School of Medicine, Shanghai, China (Prof M R Phillips MD); Durban University of Technology, Durban, South Africa (J D Pillay PhD); Exposure Assessment and Environmental Health Indicators, German Environment Agency, Berlin, Germany (D Plass DrPH, M Tobollik MPH); Sanjay Gandhi Post Graduate Institute of Medical Sciences, Lucknow, India (Prof N Prasad DM); Intergrowth 21st Study Research Centre, Nagpur, India (Prof M Purwar MD); Non-Communicable Diseases Research Center, Alborz University of Medical Sciences, Karaj, Iran (M Qorbani PhD); A T Still University, Kirksville, MO, USA (A Radfar MD); Contech International Health Consultants, Lahore, Pakistan (A Rafay MS); Contech School of Public Health, Lahore, Pakistan (A Rafay MS); Research and Evaluation Division, BRAC, Dhaka, Bangladesh (M Rahman PhD); Society for Health and Demographic Surveillance, Suri, India (R K Rai MPH); ERAWEB Program, University for Health Sciences, Medical Informatics and Technology, Hall in Tirol, Austria (S Rajsic MD); Department of Preventive Medicine, Wonju College of Medicine, Yonsei University, Wonju, South Korea (C L Ranabhat PhD); Health Science Foundation and Study Center, Kathmandu, Nepal (C L Ranabhat PhD); Schizophrenia Research Foundation, Chennai, India (T Rangaswamy PhD); Diabetes Research Society, Hyderabad, India (Prof P V Rao MD); Diabetes Research Center, Hyderabad, India (Prof P V Rao MD); Azienda Socio-Sanitaria Territoriale, Papa Giovanni XXIII, Bergamo, Italy (Prof G Remuzzi MD); Department of Biomedical and Clinical Sciences "L Sacco", University of Milan, Milan, Italy (Prof G Remuzzi MD); Research Center for Environmental Determinants of Health, School of Public Health (S Rezaei PhD), Kermanshah University of Medical Sciences, Kermanshah, Iran (S Siabani PhD); Hospital das Clínicas da Universidade Federal de Minas Gerais, Belo Horizonte, Brazil (Prof A L Ribeiro MD); RMIT University, Bundoora, VIC, Australia (Prof S R Robinson PhD); Campus MAR, Barcelona Biomedical Research Park (PRBB), ISGlobal Instituto de Salud Global de Barcelona, Barcelona, Spain (D Rojas-Rueda PhD); Golestan Research Center of Gastroenterology and Hepatology, Golestan University of Medical Sciences, Gorgan, Iran (G Roshandel PhD); Institute of Epidemiology and Medical Biometry, Ulm University, Ulm, Germany (Prof G Nagel PhD, Prof D Rothenbacher MD); Universidad Tecnica del Norte, Ibarra, Ecuador (E Rubagotti PhD); Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania (G M Ruhago PhD, B F Sunguya PhD); Managerial Epidemiology Research Center, Department of Public Health, School of Nursing and Midwifery, Maragheh University of Medical Sciences, Maragheh, Iran (S Safiri PhD); Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur, Malaysia (R Sahathevan PhD); Ballarat Health Service, Ballarat, VIC, Australia (R Sahathevan PhD); Development Research and Projects Center, Abuja, Nigeria (M M Saleh MPH); Chest Research Foundation, Pune, India (S S Salvi MD); Faculty of Science, Ain Shams University, Cairo, Egypt (A M Samy PhD); J Edwards School of Medicine (J R Sanabria MD), Department of Public Health (M Sawhney PhD), Marshall University, Huntington, WV, USA; Case Western Reserve University, Cleveland, OH, USA (J R Sanabria MD); IIS-Fundacion Jimenez Diaz, Madrid, Spain (M D Sanchez-Niño PhD); Department of Community Medicine, Information and Health Decision Sciences, Center for Health Technology and Services

Research - CINTESIS, Porto, Portugal (J V Santos MD); Centre of Advanced Study in Psychology, Utkal University, Bhubaneswar, India (M Satpathy PhD); Federal University of Santa Catarina, Florianópolis, Brazil (I J C Schneider PhD); Division of Clinical Epidemiology and Ageing Research, German Cancer Research Center, Heidelberg, Germany (B Schöttker MPH); Institute of Health Care and Social Sciences, FOM University, Essen, Germany (B Schöttker MPH); Hypertension in Africa Research Team (HART), North-West University, Potchefstroom, South Africa (Prof A E Schutte PhD); UKZN Gastrointestinal Cancer Research Centre (Prof B Sartorius PhD), South African Medical Research Council, Potchefstroom, South Africa (Prof A E Schutte PhD); Charité Berlin, Berlin, Germany (F Schwendicke PhD); Department of Public Health, An-Najah University, Nablus, Palestine (A Shaheen PhD); Independent Consultant, Karachi, Pakistan (M A Shaikh MD); The George Institute for Global Health, Sydney, NSW, Australia (S M Shariful Islam PhD); Indian Institute of Technology Ropar, Rupnagar, India (R Sharma MA); Ministry of Health, Thimphu, Bhutan (J Sharma MPH); Department of Pulmonary Medicine, Zhongshan Hospital, Fudan University, Shanghai, China (J She MD); National Institute of Infectious Diseases, Tokyo, Japan (M Shigematsu PhD); Sandia National Laboratories, Albuquerque, NM, USA (M Shigematsu PhD); Washington State University, Spokane, WA, USA (K Shishani PhD); University of Technology Sydney, Sydney, NSW, Australia (S Siabani PhD); Reykjavik University, Reykjavik, Iceland (I D Sigfusdottir PhD); Federal University of Santa Catarina, Florianópolis, Brazil (D A S Silva PhD); Brasília University, Brasília, Brazil (D G A Silveira MD); University of Pennsylvania, Philadelphia, PA, USA (D H Silberberg MD); Asthma Bhawan, Jaipur, India (V Singh MD); Department of Medicine, Institute of Medical Sciences, Banaras Hindu University, Varanasi, India (O P Singh PhD); Max Hospital, Gaziabad, India (Prof N P Singh MD); School of Preventive Oncology, Patna, India (D N Sinha PhD); WHO FCTC Global Knowledge Hub on Smokeless Tobacco, National Institute of Cancer Prevention, Noida, India (D N Sinha PhD); Hywel Dda University Health Board, Carmarthen, UK (E Skiadaresi MD); Bristol Eye Hospital, Bristol, UK (E Skiadaresi MD); King Khalid University Hospital, Riyadh, Saudi Arabia (B H A Sobaih MD); University of Yaoundé, Yaoundé, Cameroon (Prof E Sobngwi PhD); Yaoundé Central Hospital, Yaoundé, Cameroon (Prof E Sobngwi PhD); National School of Public Health/Oswaldo Cruz Foundation, Rio de Janeiro, Brazil (Prof T C M Sousa MPH); Department of Community Medicine, International Medical University, Kuala Lumpur, Malaysia (C T Sreeramareddy MD); Attikon University Hospital, Athens, Greece (V Stathopoulou PhD); University of East Anglia, Norwich, UK (Prof N Steel PhD); Public Health England, London, UK (Prof N Steel PhD); South African Medical Research Council Unit on Anxiety & Stress Disorders, Cape Town, South Africa (Prof D J Stein PhD); Department of Dermatology, University Hospital Muenster, Muenster, Germany (S Steinke DrMed); Deakin University, Burwood, VIC, Australia (Prof M A Stokes PhD); Department of Neuroscience, Norwegian University of Science and Technology, Trondheim, Norway (Prof L J Stovner PhD); Norwegian Advisory Unit on Headache, St Olavs Hospital, Trondheim, Norway (Prof L J Stovner PhD); Ministry of Health, KSA, Riyadh, Saudi Arabia (R Suliankatchi Abdulkader MD); Indian Council of Medical Research, New Delhi, India (S Swaminathan MD); Departments of Criminology, Law & Society, Sociology, and Public Health, University of California, Irvine, Irvine, CA, USA (Prof B L Sykes PhD); Griffith University, Gold Coast, QLD, Australia (S K Tadakamadla PhD); WSH Institute, Ministry of Manpower, Singapore, Singapore (J S Takala DSc); Tampere University of Technology, Tampere, Finland (J S Takala DSc); Chaim Sheba Medical Center, Tel Hashomer, Israel (Prof D Tanne MD); Tel Aviv University, Tel Aviv, Israel (Prof D Tanne MD); Ethiopian Public Health Association, Addis Ababa, Ethiopia (Y L Tarekn MS); New York Medical Center, Valhalla, NY, USA (M Tavakkoli MD); Instituto Superior de Ciências da Saúde Egas Moniz, Almada, Portugal (Prof N Taveira PhD); Faculty of Pharmacy, Universidade de Lisboa, Lisboa, Portugal (Prof N Taveira PhD); Department of Anesthesiology, King Fahad Medical City, Riyadh, Saudi Arabia (A S Terkawi MD); Outcomes Research Consortium

(A S Terkawi MD), Cleveland Clinic, Cleveland, OH, USA (Prof E M Tuzcu MD); School of Public Health, Post Graduate Institute of Medical Education and Research, Chandigarh, India (Prof J Thakur MD); Christian Medical College Vellore, Vellore, India (Prof N Thomas PhD); Faculty of Health Sciences, Wroclaw Medical University, Wroclaw, Poland (R Topor-Madry PhD); School of Medicine, University of Valencia, Valencia, Spain (M Tortajada PhD); INSERM (French National Institute for Health and Medical Research), Paris, France (M Touvier PhD); University of Southern Santa Catarina, Palhoça, Brazil (Prof J Traebert PhD); Johns Hopkins University, Baltimore, MD, USA (B X Tran PhD); Department of Neurology, Rigshospitalet, University of Copenhagen, Copenhagen, Denmark (T Truelsén DMSc); Hanoi Medical University, Hanoi, Vietnam (B X Tran PhD); Parc Sanitari Sant Joan de Déu, Fundació Sant Joan de Déu, Universitat de Barcelona, CIBERSAM, Barcelona, Spain (S Tyrovolas PhD); Department of Internal Medicine, Federal Teaching Hospital, Abakaliki, Nigeria (K N Ukwaja MD); School of Government, Pontificia Universidad Católica de Chile, Santiago, Chile (E A Undurraga PhD); Ebonyi State University, Abakaliki, Nigeria (C J Uneke PhD); Warwick Medical School, University of Warwick, Coventry, UK (O A Uthman PhD); University of Nigeria, Nsukka, Enugu Campus, Enugu, Nigeria (Prof B S C Uzochukwu MD); UKK Institute for Health Promotion Research, Tampere, Finland (Prof T Vasankari PhD); National Centre for Disease Control, Delhi, India (S Venkatesh MD); Raffles Neuroscience Centre, Raffles Hospital, Singapore, Singapore (N Venketasubramanian MBBS); Weill Cornell Medical College, New York, NY, USA (R Vidavalur MD); VHS SNEHA, Chennai, India (L Vijayakumar PhD); University of Bologna, Bologna, Italy (Prof F S Violante MD); Federal Research Institute for Health Organization and Informatics, Moscow, Russia (S K Vladimirov PhD); National Research University Higher School of Economics, Moscow, Russia (Prof V V Vlassov MD); VA Medical Center, Washington, DC, USA (M T Wallin MD); Neurology Department, Georgetown University, Washington, DC, USA (M T Wallin MD); University of São Paulo Medical School, São Paulo, Brazil (Y Wang PhD); McGill University, Ottawa, ON, Canada (S Weichenthal PhD); Department of Research, Cancer Registry of Norway, Institute of Population-Based Cancer Research, Oslo, Norway (E Weiderpass PhD); Department of Community Medicine, Faculty of Health Sciences, University of Tromsø, The Arctic University of Norway, Tromsø, Norway (E Weiderpass PhD); Genetic Epidemiology Group, Folkhälsan Research Center, Helsinki, Finland (E Weiderpass PhD); Royal Children's Hospital, Melbourne, VIC, Australia (R G Weintraub MBBS); University of Melbourne, Melbourne, VIC, Australia (R G Weintraub MBBS); German National Cohort Consortium, Heidelberg, Germany (R Westerman PhD); West Herts Hospitals NHS Trust, Watford, UK (M E Murdoch MD); Western Health, Footscray, VIC, Australia (Prof T Wijeratne MD); Centre of Evidence Based Dermatology, University of Nottingham, Nottingham, UK (Prof H C Williams DSc); South African Medical Research Council, Cochrane South Africa, Cape Town, South Africa (Prof C S Wiysonge PhD); National Institute for Health Research Comprehensive Biomedical Research Centre, Guy's & St Thomas' NHS Foundation Trust and King's College London, London, UK (Prof C D Wolfe MD); Royal Cornwall Hospital, Truro, UK (Prof A D Woolf MBBS); St John's Medical College and Research Institute, Bangalore, India (Prof D Xavier MD); Department of Neurology, Jinling Hospital, Nanjing University School of Medicine, Nanjing, China (Prof G Xu PhD); School of Public Health, University of Saskatchewan, Saskatoon, SK, Canada (M Yaghoubi MSc); Global Health Research Center, Duke Kunshan University, Kunshan, China (Prof L L Yan PhD); Department of Preventive Medicine, Northwestern University, Chicago, IL, USA (Y Yano MD); Social Work and Social Administration Department (Prof P Yip PhD), The Hong Kong Jockey Club Centre for Suicide Research and Prevention (Prof P Yip PhD), University of Hong Kong, Hong Kong, China; Department of Biostatistics, School of Public Health, Kyoto University, Kyoto, Japan (N Yonemoto MPH); Department of Preventive Medicine, College of Medicine, Korea University, Seoul, South Korea (S Yoon PhD); School of Public Health, University of Kinshasa, Kinshasa, Democratic Republic of the Congo

(M Yotebieng PhD); Jackson State University, Jackson, MS, USA (Prof M Z Younis DrPH); Department of Epidemiology and Biostatistics, School of Public Health and Global Health Institute, Wuhan University, Wuhan, China (Prof C Yu PhD); University Hospital of Setif, Setif, Algeria (Prof Z Zaidi DSc); Faculty of Medicine, Mansoura University, Mansoura, Egypt (Prof M E Zaki PhD); Ethiopian Public Health Institute, Addis Ababa, Ethiopia (E A Zegeye MS); University of Texas School of Public Health, Houston, TX, USA (X Zhang MS); MD Anderson Cancer Center, Houston, TX, USA (X Zhang MS); and Red Cross War Memorial Children's Hospital, Cape Town, South Africa (L J Zuhlke PhD).

Contributors

Please see the appendix for more detailed information about individual authors' contributions to the research, divided into the following categories: managing the estimation process; writing the first draft of the manuscript; providing data or critical feedback on data sources; developing methods or computational machinery; applying analytical methods to produce estimates; providing critical feedback on methods or results; drafting the work or revising it critically for important intellectual content; extracting, cleaning, or cataloguing data; designing or coding figures and tables; and managing the overall research enterprise.

Declaration of interests

Laith J Abu-Raddad acknowledges the support of the Qatar National Research Fund (NPRP 9-040-3-008) who provided the main funding for generating the data provided to the GBD-IHME effort. Anurag Arawal acknowledges support from the Wellcome Trust DBT India Alliance Senior Fellowship; and reports personal fees from AstraZeneca. Ashish Awasthi received funding from DST, Government of India through INSPIRE Faculty scheme. The scientific work of Aleksandra Barac is part of the Project No III45005 granted by Ministry of Education, Science and Technological Development of the Republic of Serbia. Till Bärnighausen was supported by the Alexander von Humboldt Foundation through the Alexander von Humboldt Professor award, funded by the Federal Ministry of Education and Research; the Wellcome Trust; the European Commission; the Clinton Health Access Initiative; and from NICHD of NIH (R01-HD084233), NIA of NIH (P01-AG041710), NIAID of NIH (R01-AI124389 and R01-AI112339) as well as FIC of NIH (D43-TW009775); this research was supported by NIH National Center for Advancing Translational Science (NCATS) UCLA CTSI Grant Number UL1TR001881. Yannick Béjot reports grants and personal fees from AstraZeneca, personal fees from Daiichi-Sankyo, personal fees from Pfizer-BMS, personal fees from MSD, personal fees from Bayer, personal fees from Covidien, and grants and personal fees from Boehringer-Ingelheim. Boris Bikbov has received funding from the European Union's Horizon 2020 research and innovation programme under Marie Skłodowska-Curie grant agreement No 703226; and acknowledges that work related to this paper has been done on the behalf of the GBD Genitourinary Disease Expert Group supported by the International Society of Nephrology (ISN). Rupert Bourne acknowledges support from the Brien Holden Vision Institute. Rachelle Buchbinder is funded by an Australian National Health and Medical Research Council (NHMRC) Senior Principal Research Fellowship. Lucero Cahuana-Hurtado acknowledges support from the Instituto Nacional de Salud Pública (INSP) in Mexico. Juan-Jesus Carrero acknowledges support from Stockholm County Council Swedish Heart and Lung Foundation. Cyrus Cooper reports personal fees from Alliance for Better Bone Health, Amgen, Eli Lilly, GSK, Medtronic, Merck, Novartis, Pfizer, Servier, Takeda, and UCB. José das Neves was supported in his contribution to this work by a Fellowship from Fundação para a Ciência e a Tecnologia, Portugal (SFRH/BPD/92934/2013.); Barbara de Courten is supported by National Heart Foundation Future Leader Fellowship (100864). Louisa Degenhardt is supported by an Australian National Health and Medical Research Council (NHMRC) Principal Research Fellowship; the National Drug and Alcohol Research Centre at the University of NSW is supported by funding from the Australian Government under the Substance Misuse Prevention and Service Improvements Grants Fund. Kebede Deribe is funded by a Wellcome Trust Intermediate Fellowship in Public Health and Tropical Medicine (grant number 201900); Mir Sohail Fazeli reports personal fees from Doctor Evidence LLC.

João Fernandes gratefully acknowledges funding from FCT—Fundação para a Ciência e a Tecnologia (grant number UID/Multi/50016/2013). Katherine B Gibney is supported by an NHMRC Early Career Fellowship. Shifalika Goenka is supported by the Bernard Lown Scholars in Cardiovascular Health Program, Harvard School of Public Health (2015–17) and a Wellcome Trust (Grant No 096735/B/11/Z). Amador Goodridge acknowledges support from Sistema Nacional de Investigación (SNI) de Panamá and Secretaría Nacional de Ciencia, Tecnología e Innovación (SENACYT). Simon I Hay is funded by grants from the Bill & Melinda Gates Foundation (OPP1106023, OPP1119467, OPP1093011, and OPP1132415). Shariful Islam received postdoctoral research fellowship from The George Institute for Global Health and career transition grants from High Blood Pressure Research Foundation of Australia; the Ministry of Education Science and Technological Development of the Republic of Serbia has co-financed Serbian parts of this GBD related contribution throughout the Grant OI 175 014; publication of results was not contingent upon Minsitry's prior censorship or approval. Peter James is supported by R00 CA201542 from NCI. Panniyammakal Jeemon acknowledges support from the Clinical and public health intermediate fellowship from the Wellcome Trust and Department of Biotechnology, India Alliance (2015–2020). Srinivasa Vittal Katikireddi reports grants from Chief Scientist Office and grants from Medical Research Council, during the conduct of the study. Anil Kaul has received funding (HR14-065) from the Oklahoma Council for Advancement of Science & Technology (OCAST). Christian Kieling has received support from Brazilian governmental research funding agencies Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul (Fapergs), and Hospital de Clínicas de Porto Alegre (FIPE/HCPA). Ai Koyanagi's work was supported by the Miguel Servet contract financed by the CP13/00150 and PI15/00862 projects, integrated into the National R + D + I and funded by the ISCIII—General Branch Evaluation and Promotion of Health Research—and the European Regional Development Fund (ERDF-FEDER). Tea Lallukka is supported by the Academy of Finland (Grants #287488 and #294096). Cheru T Leshargie would like to thank Debreworkos University for the arrangement of internate service to communicate for tasks such as registering and communicating with GBD organiser and to download the manuscript and other supportive document, send on comments, and sending the co-author form for the GBD. Miriam Levi acknowledges institutional support received from CeRIMP, Regional Centre for Occupational Diseases and Injuries, Local Health Unit Tuscany Center, Florence, Italy. Katharine J Looker thanks the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Evaluation of Interventions at the University of Bristol, in partnership with Public Health England (PHE), for research support; and received separate funding from the World Health Organization and Sexual Health 24 during the course of this study; the views expressed are those of the authors and not necessarily those of the National Health Service, the NIHR, the Department of Health or Public Health England. Azeem Majeed and Imperial College London are grateful for support from the NW London NIHR Collaboration for Leadership in Applied Health Research & Care. Francisco Martins-Melo acknowledges support from the postdoctoral Fellowship, Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES; Brazilian public agency). Winifred März reports grants and personal fees from Siemens Diagnostics, grants and personal fees from Aegerion Pharmaceuticals, grants and personal fees from AMGEN, grants and personal fees from AstraZeneca, grants and personal fees from Danone Research, personal fees from Hoffmann LaRoche, personal fees from MSD, grants and personal fees from Pfizer, personal fees from Sanofi, personal fees from Synageva, grants and personal fees from BASF, grants from Abbott Diagnostics, grants and personal fees from Numares AG, grants and personal fees from Berlin-Chemie, and other support from Synlab Holding Deutschland GmbH. Mohsen Mazidi was supported by The World Academy of Sciences studentship of the Chinese Academy of Sciences. John McGrath received John Cade Fellowship APP1056929 from the National Health and Medical Research Council, and Niels Bohr Professorship from the Danish

National Research Foundation. Toni Meier would like to acknowledge institutional support from the “Competence Cluster of Nutrition and Cardiovascular Health (nutriCARD)—Jena-Halle-Leipzig. Philip Mitchell's research is supported by an Australian NHMRC Program Grant number 1037196. Ulrich Mueller has received financial support from the German National Cohort Study (grant 01ER1511D). Charles Newton is supported by the Wellcome Trust, UK. Olanrewaju Oladimeji is a Senior Research Specialist at the Human Sciences Research Council (HSRC), South Africa and Doctoral Candidate at the University of KwaZulu-Natal (UKZN), South Africa; we acknowledge the institutional support by leveraging on the existing organizational research facilities at HSRC and UKZN. AO was supported by intensificacion ISCIII FEDER funds and RETIC REDINREN. Richard Osborne was funded in part through an Australian National Health and Medical Research Council (NHMRC) Senior Research Fellowship #APP1059122. MO is supported by U54HG007479 from the NIH, USA. Charles Parry would like to acknowledge support from the South African Medical Research Council. Norberto Perico acknowledges that work related to this paper has been done on behalf of the GBD Genitourinary Disease Expert Group supported by the International Society of Nephrology (ISN). Konrad Pesudovs is support by Flinders University. William Petri is support by NIH grant AI043596; Kazem Rahimi is funded by an NIHR Career Development Fellowship and is supported by the National Institute for Health Research (NIHR) Biomedical Research Centre (BRC) and the Oxford Martin School. Giuseppe Remuzzi acknowledges that the work related to this paper has been done on behalf of GBD Genitourinary Disease Expert Group supported by the International Society of Nephrology (ISN). Maria Dolores Sanchez-Niño is supported by FIS PI15/00298 (ISCIII, Spanish Government). Aletta E Schutte received support from the South African Medical Research Council and National Research Foundation (DST/NRF SARCHI Programme). Jeffrey Stanaway reports grants from Merck. Cassandra E I Szeke reports grants from National Medical Health Research Council, during the conduct of the study; grants from Lundbeck, grants from Alzheimer's Association, outside the submitted work; and has a patent PCT/AU2008/001556 issued; Rafael Tabarés-Seisdedos was supported in part by grant PROMETEOI1/2015/021 from Generalitat Valenciana and the national grants PI14/00894 and PIE14/00031 from ISCIII-FEDER. Amanda G Thrift was provided fellowship support from the National Health & Medical Research Council (NHMRC; 1042600). Stefano Tyrovola's work was supported by the Foundation for Education and European Culture (IPEP), the Sara Borrell postdoctoral programme (reference no CD15/00019 from the Instituto de Salud Carlos III (ISCIII - Spain) and the Fondos Europeo de Desarrollo Regional (FEDER). Job F M van Boven's work was supported by the University Medical Center Groningen, University of Groningen, The Netherlands. Ronny Westerman would like to acknowledge that this work is on behalf of the German National Cohort funded by the German Ministry of Education and Research. Lijing I Yan is partly supported by the National Natural Sciences Foundation of China grants (71233001 and 71490732). Marcel Yotebieng is partially supported by the NIAID U01AI096299-01 and the NICHD R01HD087993. All other authors declare no competing interests.

Acknowledgments

Research reported in this publication was supported by the Bill & Melinda Gates Foundation, the National Institute on Aging of the National Institutes of Health (award P30AG047845), and the National Institute of Mental Health of the National Institutes of Health (award R01MH110163). The content is solely the responsibility of the authors and does not necessarily represent the official views of the Bill & Melinda Gates Foundation or the National Institutes of Health. The Palestinian Central Bureau of Statistics granted the researchers access to relevant data in accordance with licence number SLN2014-3-170, after subjecting data to processing aiming to preserve the confidentiality of individual data in accordance with the General Statistics Law - 2000. The researchers are solely responsible for the conclusions and inferences drawn upon available data. We thank the Russia Longitudinal Monitoring Survey, RLMS-HSE, conducted by the National Research University Higher School of Economics and ZAO “Demoscope” together with the Carolina Population Center, University of North

Carolina at Chapel Hill, and the Institute of Sociology RAS for making these data available. This study has been realised using the data collected by the Swiss Household Panel (SHP), which is based at the Swiss Centre of Expertise in the Social Sciences FORS. The project is financed by the Swiss National Science Foundation from the Framingham Heart Study of the National Heart Lung and Blood Institute of the National Institutes of Health and Boston University School of Medicine. This work was supported by the National Heart, Lung and Blood Institute's Framingham Heart Study (contract number N01-HC-25195). The Health and Retirement Study (HRS) is sponsored by the National Institute on Aging (grant number NIA U01AG009740) and is conducted by the University of Michigan. This research used data from the National Health Survey 2003. We are grateful to the Ministry of Health, Survey copyright owner, allowing him to have the database. All results of the study are those of the authors and in no way committed to the Ministry. This research uses data from Add Health, a programme project designed by J Richard Udry, Peter S Bearman, and Kathleen Mullan Harris, and funded by a grant P01-HD31921 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development, with cooperative funding from 17 other agencies. Special acknowledgment is due to Ronald R Rindfuss and Barbara Entwisle for assistance in the original design. People interested in obtaining data files from Add Health should contact Add Health, Carolina Population Center, 123 West Franklin Street, Chapel Hill, NC 27516-2524, USA (addhealth@unc.edu). No direct support was received from grant P01-HD31921 for this analysis. The data reported here have been supplied by the US Renal Data System (USRDS). The interpretation and reporting of these data are the responsibility of the author(s) and in no way should be seen as an official policy or interpretation of the US Government. HBS-C is an international study carried out in collaboration with WHO/EURO. The International Coordinator of the 1997–98, 2001–02, 2005–06 and 2009–10 surveys was Candace Currie and the Data Bank Manager for the 1997–98 survey was Bente Wold, whereas for the following survey, Oddrun Samdal was the Databank Manager. A list of principal investigators in each country can be found at <http://www.hbsc.org>. Data used in the preparation of this article were obtained from the Pooled Resource Open-Access ALS Clinical Trials (PRO-ACT) Database. In 2011, Prize4Life, in collaboration with the Northeast ALS Consortium, and with funding from the ALS Therapy Alliance, formed the Pooled Resource Open-Access ALS Clinical Trials (PRO-ACT) Consortium. The data available in the PRO-ACT Database has been volunteered by PRO-ACT Consortium members. This paper uses data from SHARE Waves 1, 2, 3 (SHARELIFE), 4, 5, and 6 (DOIs: 10.6103/SHARE.w1.600, 10.6103/SHARE.w2.600, 10.6103/SHARE.w3.600, 10.6103/SHARE.w4.600, 10.6103/SHARE.w5.600, 10.6103/SHARE.w6.600); see Börsch-Supan and colleagues (2013) for methodological details. The SHARE data collection has been primarily funded by the European Commission through FP5 (QLK6-CT-2001-00360), FP6 (SHARE-I3: RII-CT-2006-062193, COMPARE: CIT5-CT-2005-028857, SHARELIFE: CIT4-CT-2006-028812), and FP7 (SHARE-PREP: 211909, SHARE-LEAP: 227822, SHARE M4: 261982). Additional funding from the German Ministry of Education and Research, the Max Planck Society for the Advancement of Science, the US National Institute on Aging (U01-AG09740-13S2, P01-AG005842, P01-AG08291, P30-AG12815, R21-AG025169, Y1-AG-4553-01, IAG_BSR06-11, OGH_A-04-064, HHSN271201300071C), and various national funding sources is gratefully acknowledged (see www.share-project.org). This manuscript is based on data collected and shared by the International Vaccine Institute (IVI). This manuscript was not prepared in collaboration with investigators of IVI and does not necessarily reflect the opinions or views of IVI. Collection of these data was made possible by the US Agency for International Development (USAID) under the terms of cooperative agreement GPO-A-00-08-000_D3-00. The opinions expressed are those of the authors and do not necessarily reflect the views of USAID or the US Government. Data for this research was provided by MEASURE Evaluation, funded by the US Agency for International Development (USAID). Views expressed do not necessarily reflect those of USAID, the US Government, or MEASURE Evaluation.

References

- Mathers CD, Sadana R, Salomon JA, Murray CJ, Lopez AD. Healthy life expectancy in 191 countries, 1999. *Lancet* 2001; **357**: 1685–91.
- Murray CJ, Vos T, Lozano R, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; **380**: 2197–223.
- Murray CJ, Barber RM, Foreman KJ, et al. Global, regional, and national disability-adjusted life years (DALYs) for 306 diseases and injuries and healthy life expectancy (HALE) for 188 countries, 1990–2013: quantifying the epidemiological transition. *Lancet* 2015; **386**: 2145–91.
- Kassebaum NJ, Arora M, Barber RM, et al. Global, regional, and national disability-adjusted life-years (DALYs) for 315 diseases and injuries and healthy life expectancy (HALE), 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016; **388**: 1603–58.
- Murray CJ, Salomon JA, Mathers CD, Lopez AD. Summary measures of population health. Concepts, ethics, measurement and applications. Geneva: World Health Organization, 2002.
- Murray CJ. Quantifying the burden of disease: the technical basis for disability-adjusted life years. *Bull World Health Organ* 1994; **72**: 429–45.
- GBD 2016 Mortality Collaborators. Global, regional, and national under-5 mortality, adult mortality, age-specific mortality, and life expectancy, 1970–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017; **390**: 1084–50.
- GBD 2016 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017; **390**: 1211–59.
- Sullivan DF. A single index of mortality and morbidity. *HSMHA Health Rep* 1971; **86**: 347–54.
- GBD 2016 Causes of Death Collaborators. Global, regional, and national age-sex specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017; **390**: 1151–210.
- GBD 2016 Risk Factors Collaborators. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017; **390**: 1343–420.
- Murray CJ, Ezzati M, Flaxman AD, et al. GBD 2010: design, definitions, and metrics. *Lancet* 2012; **380**: 2063–66.
- Stevens GA, Alkema L, Black RE, et al. Guidelines for Accurate and Transparent Health Estimates Reporting: the GATHER statement. *Lancet* 2016; **388**: e19–23.
- Stevens GA, Alkema L, Black RE, et al. Guidelines for Accurate and Transparent Health Estimates Reporting: the GATHER statement. *PLoS Med* 2016; **13**: e1002056.
- Wang H, Naghavi M, Allen C, et al. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016; **388**: 1459–544.
- Flaxman AD, Vos T, Murray CJ, eds. An integrative meta-regression framework for descriptive epidemiology, first edn. Seattle: University of Washington Press, 2015.
- Stanaway JD, Shepard DS, Undurraga EA, et al. The global burden of dengue: an analysis from the Global Burden of Disease Study 2013. *Lancet Infect Dis* 2016; **16**: 712–23.
- Gething PW, Casey DC, Weiss DJ, et al. Mapping *Plasmodium falciparum* mortality in Africa between 1990 and 2015. *N Engl J Med* 2016; **375**: 2435–45.
- Salomon JA, Vos T, Hogan DR, et al. Common values in assessing health outcomes from disease and injury: disability weights measurement study for the Global Burden of Disease Study 2010. *Lancet* 2012; **380**: 2129–43.
- Salomon JA, Haagsma JA, Davis A, et al. Disability weights for the Global Burden of Disease 2013 study. *Lancet Glob Health* 2015; **3**: e712–23.
- Omran AR. The epidemiologic transition: a theory of the epidemiology of population change. *Milbank Q* 2005; **83**: 731–57.

- 22 WHO. The world health report 2000. Health systems: improving performance. Geneva: World Health Organization, 2000.
- 23 Bloom DE, Cafiero ET, Jané-Llopis E, Abrahams-Gessel S, Bloom LR. The global economic burden of noncommunicable diseases. Geneva: World Economic Forum, 2011.
- 24 Fries JF. Aging, natural death, and the compression of morbidity. *N Engl J Med* 1980; **303**: 130–35.
- 25 Mundel T. Honing the priorities and making the investment case for global health. *PLoS Biol* 2016; **14**: e1002376.
- 26 Institute for Health Metrics and Evaluation. A hand up: global progress towards universal education. Seattle, Institute for Health Metrics and Evaluation, 2015.
- 27 Jamison DT, Summers LH, Alleyne G, et al. Global health 2035: a world converging within a generation. *Lancet* 2013; **382**: 1898–955.
- 28 Bhatt S, Weiss DJ, Cameron E, et al. The effect of malaria control on *Plasmodium falciparum* in Africa between 2000 and 2015. *Nature* 2015; **526**: 207–11.
- 29 Gakidou E, Cowling K, Lozano R, Murray CJ. Increased educational attainment and its effect on child mortality in 175 countries between 1970 and 2009: a systematic analysis. *Lancet* 2010; **376**: 959–74.
- 30 Wang H, Wolock TM, Carter A, et al. Estimates of global, regional, and national incidence, prevalence, and mortality of HIV, 1980–2015: the Global Burden of Disease Study 2015. *Lancet HIV* 2016; **3**: e361–87.
- 31 Murray CJ, Lopez AD, eds. Global Burden of Disease: a comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projected to 2020, 1st edn. Cambridge: Harvard School of Public Health, 1996.
- 32 Stuckler D, King L, Robinson H, McKee M. WHO's budgetary allocations and burden of disease: a comparative analysis. *Lancet* 2008; **372**: 1563–69.
- 33 Catalá-López F, García-Altés A, Álvarez-Martín E, Génova-Maleras R, Morant-Ginestar C. Does the development of new medicinal products in the European Union address global and regional health concerns? *Popul Health Metr* 2010; **8**: 34.
- 34 Chalmers I, Bracken MB, Djulbegovic B, et al. How to increase value and reduce waste when research priorities are set. *Lancet* 2014; **383**: 156–65.
- 35 Emdin CA, Oduyayo A, Hsiao AJ, et al. Association between randomised trial evidence and global burden of disease: cross sectional study (Epidemiological Study of Randomized Trials—ESORT). *BMJ* 2015; **350**: h117.
- 36 Lam J, Lord SJ, Hunter KE, Simes RJ, Vu T, Askie LM. Australian clinical trial activity and burden of disease: an analysis of registered trials in National Health Priority Areas. *Med J Australia* 2015; **203**: 97–101.
- 37 Mitchell RJ, McClure RJ, Olivier J, Watson WL. Rational allocation of Australia's research dollars: does the distribution of NHMRC funding by National Health Priority Area reflect actual disease burden? *Med J Australia* 2009; **191**: 648–52.
- 38 Aoun S, Pennebaker D, Pascal R. To what extent is health and medical research funding associated with the burden of disease in Australia? *Aust N Z J Public Health* 2004; **28**: 80–86.
- 39 Lamarre-Cliche M, Castilloux AM, LeLorier J. Association between the burden of disease and research funding by the Medical Research Council of Canada and the National Institutes of Health. A cross-sectional study. *Clin Invest Med* 2001; **24**: 83–89.
- 40 WHO. Global Health Estimates 2015: deaths by cause, age, sex, by country and by region, 2000–2015. Geneva: World Health Organization, 2016. http://www.who.int/healthinfo/global_burden_disease/estimates/en/index1.html (accessed March 14, 2017).
- 41 WHO. WHO methods and data sources for global burden of disease estimates 2000–2015. Geneva: World Health Organization, 2017.
- 42 WHO. WHO Member State DALY estimates, 2000–2015. http://www.who.int/healthinfo/global_burden_disease/estimates/en/ (accessed April 24, 2017).
- 43 Kuznik A, Lamorde M, Nyabigambo A, Manabe YC. Antenatal syphilis screening using point-of-care testing in sub-Saharan African countries: a cost-effectiveness analysis. *PLoS Med* 2010; **10**: e1001545.
- 44 Vogel JP, Habib NA, Souza JP, et al. Antenatal care packages with reduced visits and perinatal mortality: a secondary analysis of the WHO Antenatal Care Trial. *Reprod Health* 2013; **10**: 19.
- 45 Bhatt S, Weiss DJ, Cameron E, et al. The effect of malaria control on *Plasmodium falciparum* in Africa between 2000 and 2015. *Nature* 2015; **526**: 207–11.
- 46 Baird JK. Effectiveness of antimalarial drugs. *N Engl J Med* 2005; **352**: 1565–77.
- 47 Fewtrell L, Kaufmann RB, Kay D, Enanoria W, Haller L, Colford JM. Water, sanitation, and hygiene interventions to reduce diarrhoea in less developed countries: a systematic review and meta-analysis. *Lancet Infect Dis* 2005; **5**: 42–52.
- 48 Quinn TC. HIV epidemiology and the effects of antiviral therapy on long-term consequences. *AIDS* 2008; **22**: S7–12.
- 49 Stover J, Johnson P, Hallett T, Marston M, Becquet R, Timaeus IM. The Spectrum projection package: improvements in estimating incidence by age and sex, mother-to-child transmission, HIV progression in children and double orphans. *Sex Transm Infect* 2010; **86**: ii16–21.
- 50 WHO. Towards a grand convergence for child survival and health. A strategic review of options for the future building on lessons learnt from IMNCI. Geneva: World Health Organization, 2016.
- 51 GBD 2015 Healthcare Access and Quality Collaborators. Healthcare Access and Quality Index based on mortality from causes amenable to personal health care in 195 countries and territories, 1990–2015: a novel analysis from the Global Burden of Disease Study 2015. *Lancet* 2017; **390**: 231–66.
- 52 National AIDS and STI Control Programme. Kenya AIDS Indicator Survey 2012: final report. Nairobi: National AIDS and STI Control Programme, 2014.
- 53 Ministry of Health. Uganda AIDS Indicator Survey 2011. Kampala: Ministry of Health, 2012.
- 54 Uniting to Combat Neglected Tropical Diseases. The London Declaration on neglected tropical diseases. <http://unitingtocombatntds.org/resource/london-declaration> (accessed May 8, 2017).
- 55 Hopkins DR. Disease eradication. *N Engl J Med* 2013; **368**: 54–63.
- 56 Brun R, Blum J, Chappuis F, Burri C. Human African trypanosomiasis. *Lancet* 2010; **375**: 148–59.
- 57 WHO. Fifth meeting of the Emergency Committee under the International Health Regulations (2005) regarding microcephaly, other neurological disorders and Zika virus. Nov 18, 2016. <http://www.who.int/mediacentre/news/statements/2016/zika-fifth-ec/en/> (accessed May 8, 2017).
- 58 Bogoch II, Brady OJ, Kraemer MU, et al. Potential for Zika virus introduction and transmission in resource-limited countries in Africa and the Asia-Pacific region: a modelling study. *Lancet Infect Dis* 2016; **16**: 1237–45.
- 59 Wikan N, Smith DR. Zika virus: history of a newly emerging arbovirus. *Lancet Infect Dis* 2016; **16**: e119–26.
- 60 Pan American Health Organization. Zika cumulative cases. http://www.paho.org/hq/index.php?option=com_content&view=article&id=12390:zika-cumulative-cases&catid=8424:contents&Itemid=42090&lang=en (accessed May 8, 2017).
- 61 GBD 2015 Eastern Mediterranean Region Collaborators, Mokdad AH. Danger ahead: the burden of diseases, injuries, and risk factors in the eastern Mediterranean region, 1990–2015. *Int J Public Health* 2017; published online Aug 3. DOI:10.1007/s00038-017-1017-y.
- 62 Gómez-Dantés H, Fullman N, Lamadrid-Figueroa H, et al. Dissonant health transition in the states of Mexico, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2016; **388**: 2386–402.
- 63 O'Donnell MJ, Xavier D, Liu L, et al. Risk factors for ischaemic and intracerebral haemorrhagic stroke in 22 countries (the INTERSTROKE study): a case-control study. *Lancet* 2010; **376**: 112–23.
- 64 Zhang LF, Yang J, Hong Z, et al. Proportion of different subtypes of stroke in China. *Stroke* 2003; **34**: 2091–96.
- 65 Andrade LH, Alonso J, Mneimneh Z, et al. Barriers to mental health treatment: results from the WHO World Mental Health surveys. *Psychol Med* 2014; **44**: 1303–17.
- 66 Degenhardt L, Glantz M, Evans-Lacko S, et al. Estimating treatment coverage for people with substance use disorders: an analysis of data from the World Mental Health surveys. *World Psychiatry* (in press).

- 67 Mathers BM, Degenhardt L, Ali H, et al. HIV prevention, treatment, and care services for people who inject drugs: a systematic review of global, regional, and national coverage. *Lancet* 2010; **375**: 1014–28.
- 68 Jorm AF, Patten SB, Brugha TS, Mojtabai R. Has increased provision of treatment reduced the prevalence of common mental disorders? Review of the evidence from four countries. *World Psychiatry* 2017; **16**: 90–99.
- 69 Ng M, Fleming T, Robinson M, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014; **384**: 766–81.
- 70 Global Burden of Disease Health Financing Collaborator Network. Evolution and patterns of global health financing 1995–2014: development assistance for health, and government, prepaid private, and out-of-pocket health spending in 184 countries. *Lancet* **389**: 1981–2004.s
- 71 Global Burden of Disease Health Financing Collaborator Network. Future and potential spending on health 2015–40: development assistance for health, and government, prepaid private, and out-of-pocket health spending in 184 countries. *Lancet* **389**: 2005–30.
- 72 Dowell SF, Blazes D, Desmond-Hellmann S. Four steps to precision public health. *Nature* 2016; **540**: 189–91.
- 73 Golding N, Burstein R, Longbottom J, et al. Mapping under-5 and neonatal mortality in Africa, 2000–15: a baseline analysis for the Sustainable Development Goals. *Lancet* 2017. [http://dx.doi.org/S0140-6736\(17\)31758-0](http://dx.doi.org/S0140-6736(17)31758-0) (in press).
- 74 Roberts L. Hunger amplifies infectious diseases for millions fleeing the violence of Boko Haram. April 6, 2017. <http://www.sciencemag.org/news/2017/04/hunger-amplifies-infectious-diseases-millions-fleeing-violence-boko-haram> (accessed March 24, 2017).
- 75 Hoff PD. A first course in Bayesian statistical methods. New York: Springer New York, 2009.