

Accepted Manuscript

The independent and joint associations of physical activity and body mass index with myocardial infarction: The Tromsø Study

Marius Renninger, Maja-Lisa Løchen, Ulf Ekelund, Laila A. Hopstock, Lone Jørgensen, Ellisiv B. Mathiesen, Inger Njølstad, Henrik Schirmer, Tom Wilsgaard, Bente Morseth



PII: S0091-7435(18)30283-4
DOI: [doi:10.1016/j.ypmed.2018.09.005](https://doi.org/10.1016/j.ypmed.2018.09.005)
Reference: YPMED 5491
To appear in: *Preventive Medicine*
Received date: 11 April 2018
Revised date: 7 August 2018
Accepted date: 11 September 2018

Please cite this article as: Marius Renninger, Maja-Lisa Løchen, Ulf Ekelund, Laila A. Hopstock, Lone Jørgensen, Ellisiv B. Mathiesen, Inger Njølstad, Henrik Schirmer, Tom Wilsgaard, Bente Morseth, The independent and joint associations of physical activity and body mass index with myocardial infarction: The Tromsø Study. *Ypmed* (2018), doi:[10.1016/j.ypmed.2018.09.005](https://doi.org/10.1016/j.ypmed.2018.09.005)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The independent and joint associations of physical activity and body mass index with myocardial infarction: The Tromsø Study

Authors:

Marius Renninger^{1,2}, Maja-Lisa Løchen^{3,4}, Ulf Ekelund^{5,6}, Laila A. Hopstock⁷, Lone Jørgensen^{7,8}, Ellisiv B. Mathiesen^{9,10}, Inger Njølstad³, Henrik Schirmer^{4,9}, Tom Wilsgaard³, Bente Morseth^{1,3,11}

Author affiliations:

¹School of Sport Sciences, UiT The Arctic University of Norway, Tromsø, Norway

²Department of Medical Informatics, Biometry and Epidemiology (IBE), Ludwig-Maximilians-Universität München, Munich, Germany

³Department of Community Medicine, UiT The Arctic University of Norway, Tromsø, Norway

⁴Department of Cardiology, University Hospital of North Norway, Tromsø, Norway

⁵Department of Sport Medicine, Norwegian School of Sport Sciences, Oslo, Norway

⁶Norwegian Institute for Public Health, Oslo, Norway

⁷Department of Health and Care Sciences, UiT The Arctic University of Norway, Tromsø, Norway

⁸Department of Clinical Therapeutic Services, University Hospital of Northern Norway, Tromsø, Norway

⁹Department of Clinical Medicine, UiT The Arctic University of Norway, Tromsø, Norway

¹⁰Department of Neurology, University Hospital of North Norway, Tromsø, Norway

¹¹Centre for Clinical Research and Education, University Hospital of North Norway Trust, Tromsø, Norway

Corresponding author:

Bente Morseth

School of Sport Sciences, UiT The Arctic University of Norway, 9037 Tromsø, Norway

E-mail: bente.morseth@uit.no

Telephone +47 77660769

Total word count: 3094

ACCEPTED MANUSCRIPT

Abstract

Physical activity and overweight are associated with myocardial infarction (MI). However, their joint association with MI remains unclear. Our objective was to examine the independent and joint association between leisure-time physical activity (LTPA), body mass index (BMI) and MI. This prospective cohort study included 16572 men and women (47.5% women) aged 20-54 years who took part in the second Tromsø Study. At baseline in 1979-80 LTPA was assessed by questionnaire. Data on MI was collected and adjudicated through hospital and causes of death registries between 1979 and 2013. Cox proportional hazards models were used to examine the independent and joint associations between LTPA, BMI and MI. The final sample included 16104 individuals. During a median follow up of 34 years, 1613 incident cases of MI were recorded. Physical inactivity and elevated BMI were both independently associated with MI (p for trend 0.02 and < 0.001). In joint analyses, normal weight, inactive individuals had a 20% higher risk of MI compared to their active counterparts (hazard ratio (HR) 1.20 (1.02-1.41)). The highest risk of MI was seen in obese, inactive individuals when compared to normal weight, active individuals (HR 3.20 (2.30-4.44)). The risk of MI increased with increasing BMI regardless of the activity level. HRs were lower for active compared to inactive individuals within the same BMI category. The findings suggest that LTPA and BMI are independently associated with risk of MI. LTPA seems to attenuate but not eliminate the risk of MI associated with excess bodyweight.

Keywords: Exercise, epidemiology, myocardial infarction, obesity, overweight

Introduction

The incidence and mortality rates of coronary heart disease (CHD) have declined in Western countries in recent years ^{1,2}. This trend is also seen in the Norwegian population, with a 24% decrease in the incidence rate for acute myocardial infarction (MI) among those 45 years or older between 2001-2009 ³. The decrease in CHD incidence rates are mainly explained by favorable changes in cardiovascular disease (CVD) risk factors such as decreasing smoking prevalence, systolic blood pressure and total cholesterol values ^{2,4,5}, and possibly a favorable increase in leisure time physical activity (LTPA) ^{2,5}.

LTPA is an important protective factor for the development of CHD with a risk reduction of about 20-30% in active compared to inactive individuals ^{6,7}. Excess bodyweight on the contrary has shown to increase the risk for CHD and in particular MI by 20-60% in overweight and obese individuals respectively, compared with those of normal weight ⁸⁻¹⁰. Most studies examining the joint effects of physical activity and body mass index (BMI) on CVD and CHD have reported that physical activity attenuated but did not fully eliminate the increased CVD risk associated with overweight and obesity ¹¹⁻¹⁶. However, one recent study showed that overweight and obese active individuals do not have an elevated CVD risk when compared to normal weight, active individuals ¹⁷. Most previous studies were conducted in single sex only ^{11-13,15}, and examined either the associations of physical activity and obesity with CVD ^{12,16,17} or CHD ^{11,13-15}. The exposure of interest in earlier research was LTPA ^{11-13,15,17} or a combination of occupational and leisure-time physical activity ^{14,16}.

The aim of this study was therefore to assess the relationship between LTPA, BMI and MI in both sexes. We examined whether the association between LTPA and MI risk differs across BMI categories and if LTPA can counteract the potential excess risk of MI associated with overweight and obesity.

Methods

Study population

The data were derived from the Tromsø Study, a population-based, prospective cohort study with repeated surveys (Tromsø 1-7, 1974-2016). The study design and data collection are described elsewhere¹⁸. For the present study, data from 16572 men and women aged 20-54 years who took part in the second Tromsø Study survey in 1979-1980 (participation rate 74%) were included.

After excluding individuals with incomplete data on BMI, LTPA, daily smoking, blood pressure treatment, self-reported diabetes (n = 412) and subjects with prevalent MI at baseline (n = 56), the final sample included 8235 men (51.1%) and 7869 women (48.9%). The Tromsø study is approved by the Norwegian Data Inspectorate and the Regional Committee of Research Ethics.

Measurements

Assessment of LTPA and BMI

At baseline in 1979-1980, LTPA was self-reported using a validated questionnaire¹⁹ that was first introduced by Saltin and Grimby in 1968 and participants were categorized into four physical activity categories. I (inactive): reading, watching TV, or other sedentary activity; II (moderately active): walking, cycling, or other forms of exercise at least 4 h/week; III (highly active): participation in recreational sports, heavy gardening, etc. at least 4 h/week; IV (vigorously active): participation in hard training or sports competitions regularly several times a week.

Height and weight were measured to the nearest centimeter and half-kilogram without shoes and with light clothing. BMI was calculated as weight in kilograms divided by the square of height in meters (kg/m^2). For the combined BMI/LTPA categories, normal-weight was defined as $\text{BMI} < 25\text{kg}/\text{m}^2$, overweight as $\text{BMI} 25\text{-}<30\text{kg}/\text{m}^2$ and obesity as $\text{BMI} \geq 30\text{kg}/\text{m}^2$. Individuals in activity categories II, III and IV were classified as physically active according to physical activity recommendations²⁰, whereas individuals in category I were classified as inactive.

Assessment of covariates

Information about covariates was collected at baseline by self-report and physical examinations at the study site by trained technicians. Self-reported data on current smoking (yes/no), diabetes (yes/no), years of education (No. of years including primary and secondary school) and blood pressure treatment (yes/no) were derived from the questionnaire. Blood pressure was measured with a mercury sphygmomanometer and stethoscope using standard procedures, and total cholesterol was derived from non-fasting blood samples analyzed by standard methods at the Department of Laboratory Medicine of the University Hospital of North Norway.

Assessment of MI

The outcome in this study was first-ever MI that occurred between the baseline examination in 1979-1980 and the end of follow-up 31 December 2013. The following diagnostic codes in the discharge diagnosis register at the University Hospital of North Norway, the only hospital serving the area of Tromsø, were included to identify all possible incident MI cases: International Classification of Diseases (ICD) 8 codes 410-414; ICD 9 codes 410-414 and ICD 10 codes I20-I25. When appropriate, discharge letters from other hospitals were collected as well. To identify fatal out-of-hospital cases of MI as well as deaths that occurred outside Tromsø, the Tromsø Study participant list was linked with the National Cause of Death Registry at the Norwegian Institute of Public Health. Death certificates for those with an underlying or contributing diagnosis of CVD or sudden death were retrieved and additional information from autopsy records and records from general practitioners, ambulance services and nursing homes was collected. Thereafter, cases were validated by reviewing the medical records and death certificates by trained physicians from an independent end-point committee ⁵.

Information on migration was obtained through the Population Registry of Norway.

Each participant contributed person-time from the baseline examination in 1979-80 until the date of the first MI, migration, death or end of follow-up 31 December 2013, whichever came first.

Statistical analyses

Cox proportional hazard models were used to estimate hazard ratios (HR) and 95% confidence intervals (CI) for the association between independent and joint BMI/LTPA categories and the risk of MI. To determine the associations between BMI and MI, and between LTPA and MI, normal-weight and inactive individuals, respectively, were used as the reference category. For the joint BMI/LTPA analyses, individuals were grouped into six categories and normal-weight active individuals were used as the reference category. Three different models were used in the Cox regression analyses. Model 1 was adjusted for age and sex; Model 2 was adjusted for age, sex, daily smoking and BMI in the LTPA model and LTPA in the BMI model. In an additional model, systolic and diastolic blood pressure, total cholesterol, and self-reported diabetes and hypertension treatment were added as covariates (Model 3). Since these variables might be considered as intermediate factors in the causal pathway of BMI/LTPA and MI, model 2 was used as the main model. Possible interactions between BMI and LTPA were evaluated by adding multiplicative interaction terms to the models. To examine the associations between LTPA and MI in different weight groups, individuals were stratified into three BMI groups (normal-weight, overweight, obese). Interactions between sex and BMI group, and sex and LTPA level were tested in all models and no significant modification by sex was observed. Similarly, there was no indication of a BMI by LTPA interaction in any of the models (data not shown).

In sensitivity analyses we additionally adjusted models 2 and 3 for years of education as a proxy for socio-economic status in individuals with valid data on education (n=14002), and the results were virtually unchanged (data not shown). To reduce the chance of reverse causation, we conducted a sensitivity analysis where individuals with MI within the first two years of follow-up were excluded (n=38), with virtually unchanged results.

Proportional hazard assumptions were examined for each model and were not violated. All analyses were performed using IBM SPSS Statistics for Windows, version 24 (IBM Corp., Armonk, N.Y., USA). The first author M.R. and senior author B.M. had full access to all the data in the study and take responsibility for its integrity and the data analysis.

Results

Baseline characteristics

In total, 7869 women (48.9%) and 8235 men (51.1%) were included in the analyses. During a median follow-up of 34.1 years (33.8 and 34.2 years for the 25th and the 75th percentile, respectively; 499196 person-years), 344 women (aged 40.6y, SD 6.5) and 1269 men (aged 41.6y, SD 8.8) had an incident MI. The overall incidence of MI was 3.18/1000 person-years, 1.35/1000 person-years in women and 5.18/1000 person-years in men (Table 1). Descriptive data are presented in Table 1.

LTPA and MI

In general, we observed an inverse relationship between LTPA and risk of MI (Table 2). In Model 1 (adjusted for age and sex), moderately and highly active individuals had 23% and 29% reduced risk of MI [HRs 0.77, 95% CI 0.68-0.87 and 0.71, 95% CI 0.62-0.83] compared with the inactive group. The MI risk among individuals who reported vigorous activity was halved compared with the inactive group [HR 0.44, 95% CI 0.29-0.65]. Adding smoking and BMI to the model (Model 2) attenuated these associations, although the associations were still statistically significant in the moderate and vigorous activity group. In the additional model (Model 3), the associations were further weakened but moderately active individuals still showed a significantly reduced risk of MI compared to inactive individuals [HR 0.87, 95% CI 0.77-0.99].

BMI and MI

The risk of MI increased monotonically with elevated BMI, by 19%, 21% and 12% for every two units of BMI increment in Model 1, 2, and 3 [HR 1.19, 95% CI 1.16-1.23; HR 1.21, 95% CI 1.18-1.25 and HR 1.12, 95% CI 1.09-1.16] (Table 2). When individuals were categorized into BMI groups, those in the overweight and obese groups had an elevated risk for MI with the highest risk in the obese group [HR 2.53 95% CI 2.10-3.05; HR 2.70 95% CI 2.24-3.26] in Model 1 and Model 2, respectively. In the additional model (Model 3), the HRs were attenuated although still statistically significant. The MI risk in overweight and obese individuals was 29% and 88% higher compared to normal-weight individuals [HRs 1.29, 95% CI 1.16-1.44 and 1.88, 95% CI 1.55-2.28].

Joint associations of LTPA and BMI with MI

In Model 1, the risk of MI was 37% higher in normal weight, inactive individuals than in their active counterparts [HR 1.37, 95% CI 1.16-1.61] (Table 3). When adjusting for covariates (Models 2 and 3), the risk increase was attenuated, although statistically significant [HRs 1.20, 95% CI 1.02-1.41 and 1.17, 95% CI 1.00 -1.38]. The highest risk of MI was seen in obese, inactive individuals, with a 3-fold risk compared to normal-weight, active individuals in Model 2 [HR 3.20, 95% CI 2.30-4.44]. Irrespective of the activity level, the risk of MI increased with increasing BMI (Table 3). However, the risk estimates were lower for active individuals compared to inactive individuals within the same weight category.

The risk of MI was higher in inactive than active individuals across BMI groups (Table 4). This was observed in crude and adjusted models in the normal-weight group, whereas the association in multivariable models was non-significant in overweight and obese individuals.

Discussion

In this cohort of young and middle-aged individuals followed for 34 years, both elevated BMI and leisure-time physical inactivity were independently associated with an increased risk of MI in a dose-response manner. Active individuals in the same weight category showed lower risk estimates than their inactive counterparts. However, the magnitude of association seems to be stronger for obesity compared to physical inactivity. Being obese and inactive was associated with the highest risk of MI and LTPA seems to attenuate but not eliminate the risk for MI associated with overweight and obesity.

Possible mechanisms

Previous studies have shown that the relationship between obesity and CHD is substantially mediated by traditional CVD risk factors, mainly blood pressure, cholesterol and blood glucose, explaining about half of the increased risk associated with high BMI²¹. In this study, the risk estimates for the association between BMI and MI were attenuated following adjustment for CVD risk factors, particularly in the obese group.

Similarly, adjusting for traditional CVD risk factors such as blood pressure, total cholesterol levels, and diabetes status attenuated the association between LTPA and MI. This may partially be explained by the mediating effects of these factors on the association between physical inactivity with increased risk of MI corroborating previous findings²². Others have suggested that the pronounced risk reduction in active compared with inactive individuals may include enhanced endothelial function and a positive impact on the autonomic nervous system, such as improved peripheral baroreflex function²³.

Both the association between LTPA and MI and the association between BMI and MI seem to be largely mediated by CVD risk factors (blood pressure, total cholesterol, diabetes status), and adjustment for these factors might reflect over-adjustment, as indicated by the attenuated risk estimates in Model 3.

Comparison with other studies

The association between increased physical activity and a decreased risk of MI observed in our study is consistent with findings from several previous studies^{15,22,24}, and the associations between overweight and obesity with risk of MI have been reported previously^{16,25}.

Our findings regarding the joint association between LTPA, BMI and MI are consistent with those of several observational studies. In a 20-year follow-up of 88 393 women in the Nurses' Health Study¹³, high levels of adiposity and physical inactivity independently predicted increased risk of CHD. The highest CHD risk was seen in obese and inactive women, with a more than 3-fold risk of CHD compared with the normal-weight active group. In agreement with the present study, LTPA attenuated but did not eliminate the adverse effects of overweight on CHD risk¹³. These results were recently supported by data from the Nurses' Health Study II¹¹, suggesting that LTPA was protective against CHD across BMI categories. Further, results from the Women's Health Study,¹⁵ indicate similar findings, including independent associations between self-reported LTPA and BMI with CHD risk. However, the reported risk estimates for obese inactive women were lower than in our study. This may be explained by the shorter follow-up time and a different assessment of LTPA as well as the inclusion of

percutaneous transluminal coronary angioplasty and coronary artery bypass graft in the outcome definition¹⁵. Carlsson et al.¹² examined the associations between repeated measurements of BMI and self-reported LTPA in the ULSAM-study with risk of CVD and reported similar results as those we report here. Furthermore, our findings are in contrast to studies that identified the obesity paradox. In our study, individuals with a higher BMI also had a higher risk of MI, while active individuals had a lower risk of MI than inactive individuals independent of their BMI. Taken together, it seems that LTPA attenuates but does not fully eliminate the increased risk of adiposity associated with MI. This suggests that public health initiatives should strive to reduce obesity and simultaneously increase physical activity in populations optimally reducing the risk of MI and other CVD. It should be emphasized that physical activity is important also in the obese and should be encouraged.

Strengths and limitations

The study has several strengths, including the prospective design with a long follow-up period of 34 years, rigorous outcome ascertainment, a large number of MI cases, a population-based sample and minimal loss to follow-up.

A limitation of the present study is that we did not use updated information on the exposure variables. This might lead to some misclassification due to changes in LTPA and BMI during follow-up.

Self-reported physical activity is prone to misclassification error. However, due to the prospective design of the present study, reported physical activity level is not biased by future disease status, and misclassification of physical activity will most likely be non-differential. Furthermore, Emaus et al.²⁶ observed overestimation of physical activity by self-report in the participants of the Tromsø study, which will most likely lead to underestimation of the true association between LTPA and MI.

The stronger magnitude of associations observed between BMI compared to LTPA and MI may be explained by the differences in measurement precision of the two exposure variables²⁷. Height and weight were measured objectively by trained technicians, whereas LTPA was self-reported and therefore more likely to be affected by random measurement error.

There is inconsistent evidence as to whether BMI, waist-to-hip ratio (WHR) or waist-circumference (WC) is the superior indicator of obesity. While a widely recognized case-control study showed a stronger relation between WHR and MI ²⁸, a more recent collaborative analysis of 58 cohort studies refuted these results and showed a similar association of BMI, WHR and WC with CVD risk ²⁹. Furthermore, the long-term reproducibility of BMI was higher than that of WHR or WC ²⁹. Thus, although BMI might slightly underestimate the association of obesity and MI, BMI seems to be a valid indicator of overweight and obesity in large population-based studies.

Finally, we cannot exclude the possibility that our observations are influenced by unmeasured confounders (e.g. diet and genotype).

Conclusion

This prospective cohort study suggests an independent association between BMI and LTPA with incident MI. LTPA reduced the risk of MI in both normal-weight and overweight individuals, but did not fully eliminate the increased risk of MI associated with overweight and obesity. These findings underline the importance of public health interventions targeted at weight loss and increasing LTPA to reduce the incidence of MI and other cardiovascular diseases.

Acknowledgments: None

Sources of Funding: This work is funded by the Northern Norway Regional Health Authority.

Conflict of Interest: none

References

1. Yeh RW, Sidney S, Chandra M, Sorel M, Selby JV, Go AS. Population Trends in the Incidence and Outcomes of Acute Myocardial Infarction. *New England Journal of Medicine*. 2010;362(23):2155-2165.
2. Hardoon SL, Whincup PH, Lennon LT, Wannamethee SG, Capewell S, Morris RW. How much of the recent decline in the incidence of myocardial infarction in British men can be explained by changes in cardiovascular risk factors? Evidence from a prospective population-based study. *Circulation*. 2008;117(5):598-604.
3. Sulo G, Iglund J, Nygard O, Vollset SE, Ebbing M, Tell GS. Favourable trends in incidence of AMI in Norway during 2001-2009 do not include younger adults: a CVDNOR project. *European journal of preventive cardiology*. 2014;21(11):1358-1364.
4. Eggen AE, Mathiesen EB, Wilsgaard T, Jacobsen BK, Njolstad I. Trends in cardiovascular risk factors across levels of education in a general population: is the educational gap increasing? The Tromso study 1994-2008. *Journal of epidemiology and community health*. 2014;68(8):712-719.
5. Mannsverk J, Wilsgaard T, Mathiesen EB, et al. Trends in Modifiable Risk Factors Are Associated With Declining Incidence of Hospitalized and Nonhospitalized Acute Coronary Heart Disease in a Population. *Circulation*. 2016;133(1):74-81.
6. Sattelmair J, Pertman J, Ding EL, Kohl HW, 3rd, Haskell W, Lee IM. Dose response between physical activity and risk of coronary heart disease: a meta-analysis. *Circulation*. 2011;124(7):789-795.
7. Sofi F, Capalbo A, Cesari F, Abbate R, Gensini GF. Physical activity during leisure time and primary prevention of coronary heart disease: an updated meta-analysis of cohort studies. *Eur J Cardiovasc Prev Rehabil*. 2008;15(3):247-257.
8. Mongraw-Chaffin ML, Peters SA, Huxley RR, Woodward M. The sex-specific association between BMI and coronary heart disease: a systematic review and meta-analysis of 95 cohorts with 1.2 million participants. *The lancet Diabetes & endocrinology*. 2015;3(6):437-449.
9. Zhu J, Su X, Li G, Chen J, Tang B, Yang Y. The incidence of acute myocardial infarction in relation to overweight and obesity: a meta-analysis. *Archives of medical science : AMS*. 2014;10(5):855-862.
10. Yusuf S, Hawken S, Ounpuu S, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet (London, England)*. 2004;364(9438):937-952.
11. Chomistek AK, Henschel B, Eliassen AH, Mukamal KJ, Rimm EB. Frequency, Type, and Volume of Leisure-Time Physical Activity and Risk of Coronary Heart Disease in Young Women. *Circulation*. 2016;134(4):290-299.
12. Carlsson AC, Arnlov J, Sundstrom J, Michaëlsson K, Byberg L, Lind L. Physical activity, obesity and risk of cardiovascular disease in middle-aged men during a median of 30 years of follow-up. *European journal of preventive cardiology*. 2016;23(4):359-365.
13. Li TY, Rana JS, Manson JE, et al. Obesity as compared with physical activity in predicting risk of coronary heart disease in women. *Circulation*. 2006;113(4):499-506.

14. Arsenault BJ, Rana JS, Lemieux I, et al. Physical inactivity, abdominal obesity and risk of coronary heart disease in apparently healthy men and women. *International journal of obesity (2005)*. 2010;34(2):340-347.
15. Weinstein AR, Sesso HD, Lee IM, et al. The joint effects of physical activity and body mass index on coronary heart disease risk in women. *Arch Intern Med*. 2008;168(8):884-890.
16. Hu G, Tuomilehto J, Silventoinen K, Barengo N, Jousilahti P. Joint effects of physical activity, body mass index, waist circumference and waist-to-hip ratio with the risk of cardiovascular disease among middle-aged Finnish men and women. *European heart journal*. 2004;25(24):2212-2219.
17. Koolhaas CM, Dhana K, Schoufour JD, Ikram MA, Kavousi M, Franco OH. Impact of physical activity on the association of overweight and obesity with cardiovascular disease: The Rotterdam Study. *European journal of preventive cardiology*. 2017;24(9):934-941.
18. Jacobsen BK, Eggen AE, Mathiesen EB, Wilsgaard T, Njølstad I. Cohort profile: the Tromsø Study. *International journal of epidemiology*. 2012;41(4):961-967.
19. Aires N, Selmer R, Thelle D. The validity of self-reported leisure time physical activity, and its relationship to serum cholesterol, blood pressure and body mass index. A population based study of 332,182 men and women aged 40-42 years. *Eur J Epidemiol*. 2003;18(6):479-485.
20. Haskell WL, Lee IM, Pate RR, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Circulation*. 2007;116(9):1081-1093.
21. Lu Y, Hajifathalian K, Ezzati M, Woodward M, Rimm EB, Danaei G. Metabolic mediators of the effects of body-mass index, overweight, and obesity on coronary heart disease and stroke: a pooled analysis of 97 prospective cohorts with 1.8 million participants. *Lancet (London, England)*. 2014;383(9921):970-983.
22. Mora S, Cook N, Buring JE, Ridker PM, Lee IM. Physical activity and reduced risk of cardiovascular events: potential mediating mechanisms. *Circulation*. 2007;116(19):2110-2118.
23. Joyner MJ, Green DJ. Exercise protects the cardiovascular system: effects beyond traditional risk factors. *The Journal of physiology*. 2009;587(Pt 23):5551-5558.
24. Sesso HD, Paffenbarger RS, Lee I-M. Physical Activity and Coronary Heart Disease in Men. *The Harvard Alumni Health Study*. 2000;102(9):975-980.
25. Marin A, Medrano MJ, Gonzalez J, et al. Risk of ischaemic heart disease and acute myocardial infarction in a Spanish population: observational prospective study in a primary-care setting. *BMC public health*. 2006;6:38.
26. Emaus A, Degerstrøm J, Wilsgaard T, et al. Does a variation in self-reported physical activity reflect variation in objectively measured physical activity, resting heart rate, and physical fitness? Results from the Tromsø study. *Scandinavian journal of public health*. 2010;38(5 Suppl):105-118.
27. Hutcheon JA, Chioloro A, Hanley JA. Random measurement error and regression dilution bias. *BMJ (Clinical research ed)*. 2010;340:c2289.
28. Yusuf S, Hawken S, Ounpuu S, et al. Obesity and the risk of myocardial infarction in 27,000 participants from 52 countries: a case-control study. *Lancet (London, England)*. 2005;366(9497):1640-1649.
29. Wormser D, Kaptoge S, Di Angelantonio E, et al. Separate and combined associations of body-mass index and abdominal adiposity with cardiovascular disease: collaborative analysis of 58 prospective studies. *Lancet (London, England)*. 2011;377(9771):1085-1095.

Tables:

TABLE 1: Baseline characteristics and incidence of MI by BMI-LTPA categories.

The Tromsø Study 1979-2013

n = 16104	normal weight / active	normal weight / inactive	overweight / active	overweight / inactive	obese / active	obese / inactive
n (% of all)	9390 (58.3)	2340 (14.5)	3003 (18.6)	841 (5.2)	395 (2.5)	135 (0.8)
Men (%)	45.6	42.1	69.3	71.7	52.9	57.8
Age (years)	33.2 (8.5)	32.8 (8.4)	37.4 (8.7)	37.5 (8.8)	39.0 (8.5)	38.5 (8.3)
Body height (cm)	170.4 (9.2)	169.5 (9.0)	172.2 (9.1)	172.3 (9.1)	169.1 (9.7)	169.7 (11.1)
Body weight (kg)	64.2 (9.4)	62.9 (9.5)	79.2 (9.0)	79.9 (9.2)	92.8 (11.7)	96.3 (14.7)
BMI (kg/m ²)	22.0 (1.8)	21.8 (1.9)	26.7 (1.3)	26.8 (1.4)	32.4 (2.5)	33.4 (3.6)
Systolic blood-pressure (mmHg)	123.8 (13.5)	123.4 (13.5)	130.9 (15.0)	130.8 (14.1)	135.9 (15.8)	136.5 (17.5)
Diastolic blood-pressure (mmHg)	78.5 (9.8)	78.3 (10.1)	84.1 (10.5)	84.6 (10.4)	89.6 (10.8)	89.9 (10.6)
Total cholesterol (mmol/l)	5.6 (1.2)	5.7 (1.2)	6.3 (1.2)	6.3 (1.3)	6.6 (1.4)	6.9 (1.2)
Incidence (cases/1000py)	2.2	2.6	5.5	6.6	8.3	10.6
Smoking in % (n)	47.4 (4448)	62.4 (1461)	43.1 (1294)	54.7 (460)	40.3 (159)	53.3 (72)
Treatment for hypertension, % (n)	0.9 (81)	1.0 (24)	3.0 (89)	3.2 (27)	7.8 (31)	10.4 (14)
Diabetes, % (n)	0.2 (22)	0.3 (8)	0.5 (14)	0.2 (2)	1.5 (6)	0.0 (0)

Data are presented as mean (SD) or % (n).

Table 2: Risk of MI in relation to LTPA and BMI. The Tromsø Study 1979-2013.**Data are Hazard Ratios (95% CI).**

LTPA level	n	MI events	Model 1	Model 2	Model 3
Low	3316	388	1.00 (ref.)	1.00 (ref.)	1.00 (ref.)
Moderate	8963	837	0.77 (0.68-0.87)	0.87 (0.77-0.98)	0.87 (0.77-0.99)
High	3278	362	0.71 (0.62-0.83)	0.88 (0.76-1.02)	0.91 (0.79-1.06)
Vigorous	547	26	0.44 (0.29-0.65)	0.63 (0.42-0.95)	0.70 (0.47-1.05)
P for trend			< 0.001	0.02	0.10
Per physical activity category			0.83 (0.77-0.89)	0.92 (0.86-0.99)	0.94 (0.88-1.01)
BMI category					
Underweight	402	10	0.83 (0.44-1.56)	0.73 (0.39-1.36)	0.80 (0.42-1.49)
Normal weight	11328	825	1.00 (ref.)	1.00 (ref.)	1.00 (ref.)
Overweight	3844	647	1.47 (1.33-1.64)	1.54 (1.39-1.72)	1.29 (1.16-1.44)
Obese	530	131	2.53 (2.10-3.05)	2.70 (2.24-3.26)	1.88 (1.55-2.28)
P for trend			< 0.001	< 0.001	< 0.001
BMI per 2 kg/m ²			1.19 (1.16-1.23)	1.21 (1.18-1.25)	1.12 (1.09-1.16)

Effect estimates are presented as HR (95% CI), n=16104, Model 1: adjusted for age and sex, Model 2: adjusted for age, sex, BMI/LTPA and daily smoking, Model 3: adjusted for age, sex, BMI/LTPA, systolic blood pressure, diastolic blood pressure, total cholesterol, diabetes, hypertension treatment and daily smoking

Table 3: Joint associations of LTPA and BMI on MI risk. The Tromsø Study 1979-2013

BMI-LTPA category	n	MI events	Model 1	Model 2	Model 3
Normal weight/active	9390	643	1.00 (ref.)	1.00 (ref.)	1.00 (ref.)
Normal weight/inactive	2340	192	1.37 (1.16-1.61)	1.20 (1.02-1.41)	1.17 (1.00-1.38)
Overweight/active	3003	489	1.50 (1.33-1.69)	1.58 (1.40-1.78)	1.31 (1.16-1.48)
Overweight/inactive	841	158	1.87 (1.57-2.23)	1.77 (1.49-2.11)	1.45 (1.22-1.74)
Obese/active	395	93	2.51 (2.01-3.12)	2.74 (2.20-3.41)	1.92 (1.54-2.40)
Obese/inactive	135	38	3.34 (2.40-4.63)	3.20 (2.30-4.44)	2.10 (1.51-2.92)
P for trend			< 0.001	< 0.001	< 0.001

Effect estimates are presented as HR (95% CI), n=16104, Model 1: adjusted for age and sex, , Model 2: adjusted for age, sex and daily smoking, Model 3: adjusted for age, sex, systolic blood pressure, diastolic blood pressure, total cholesterol, diabetes, hypertension treatment and daily smoking

Table 4: Risk of MI in relation to LTPA stratified by BMI groups. The Tromsø Study 1979-2013

LTPA level	n	MI events	Model 1	Model 2	Model 3
Normalweight	11730	835			
Active	9390	643	1.00 (ref)	1.00 (ref)	1.00 (ref)
Inactive	2340	192	1.38 (1.18-1.62)	1.19 (1.01-1.40)	1.17(1.00 -1.37)
Overweight	3844	647			
Active	3003	489	1.00 (ref.)	1.00 (ref)	1.00 (ref)
Inactive	841	158	1.24 (1.03-1.48)	1.13 (0.94-1.36)	1.11 (0.93-1.34)
Obese	530	131			
Active	395	93	1.00 (ref.)	1.00 (ref)	1.00 (ref)
Inactive	135	38	1.30 (0.89-1.90)	1.19 (0.81-1.75)	1.10 (0.75-1.62)

Effect estimates are presented as HR (95% CI), n=16104, Model 1: adjusted for age and sex, , Model 2: adjusted for age, sex and daily smoking, Model 3: adjusted for age, sex, systolic blood pressure, diastolic blood pressure, total cholesterol, diabetes, hypertension treatment and daily smoking

Highlights

- Low physical activity (PA) and high BMI independently linked to heart attack risk
- Lower heart attack risk in active than inactive individuals within each BMI category
- PA attenuates, but does not eliminate risk of heart attack associated with high BMI

ACCEPTED MANUSCRIPT