

Table II.

Asymptomatic patients examined with routine CT KUB.

<b>Asymptomatic patients</b>	Total	Women	Men
<b>Patients, n (%)</b>	171 (90)	73 (92)	98 (89)
<b>Age, yr, median (mean)</b>	56 (55.7)	55 (55.6)	57 (55.9)
Range	18-89	19-89	18-84
<b>Duration of symptoms at referral, days, median (mean)<sup>a</sup></b>	1 (3.3)	2 (3.7)	1 (3.1)
Range	1-20	1-19	1-20
<b>Previous known urolithiasis, n (%)<sup>b</sup></b>	71 (41,5)	29 (40)	42 (43)
<b>Haematuria<sup>c</sup></b>	98 (57.3)	46 (63)	52 (53)
<b>Time from referral to CT, days, median (mean)</b>	17 (26.4)	21 (28.5)	16 (24.8)
Range	1-106	1-106	1-105
<b>Urolithiasis on CT KUB, n (%)</b>	40 (23)	12 (16)	28 (29)
<b>Surgical intervention, n (%)</b>	3 (1.8)	0 (0)	3 (3.1)

<sup>a</sup> Missing information in 48 (28%) of cases.

<sup>b</sup> Missing information in 64 (37%) cases, defined as not having a history of urolithiasis in calculation of percentage.

<sup>c</sup> Missing information in 52 (30%) cases, defined as not having haematuria in calculation of percentage.