



ELSEVIER

Contents lists available at [ScienceDirect](https://www.elsevier.com/locate/dib)

## Data in Brief

journal homepage: [www.elsevier.com/locate/dib](https://www.elsevier.com/locate/dib)

## Data Article

# Data on RT-qPCR assay of nuclear progesterone receptors (nPR), membrane progesterone receptors (mPR) and progesterone receptor membrane components (PGRMC) from human uterine endometrial tissue and cancer cells of the Uterine Cervix



Natalia Smaglyukova<sup>a</sup>, Elise Thoresen Sletten<sup>b,c,d</sup>, Anne Ørbo<sup>c,e</sup>, Georg Sager<sup>a,f,\*</sup>

<sup>a</sup> Research group for Experimental and Clinical Pharmacology, Department of Medical Biology, Arctic University of Norway, Tromsø, Norway

<sup>b</sup> Department of Gynecologic Oncology, Clinic for Surgery, Cancer and Women's Diseases, University Hospital of North Norway, Tromsø, Norway

<sup>c</sup> Research group for Gynecologic Oncology, Department of Medical Biology, Faculty of Health Sciences, Arctic University of Norway, Tromsø, Norway

<sup>d</sup> Department of Clinical Medicine, Faculty of Health Sciences, Arctic University of Norway, Tromsø, Norway

<sup>e</sup> Department of Clinical Pathology, University Hospital of North Norway, Tromsø, Norway

<sup>f</sup> Clinical Pharmacology, Department of Laboratory Medicine, University Hospital of North Norway, Tromsø, Norway

## ARTICLE INFO

## Article history:

Received 14 May 2020

Revised 20 June 2020

Accepted 22 June 2020

Available online 25 June 2020

## Keywords:

RT-qPCR

mRNA

Progesterone

Nuclear receptors

Membrane receptors

Receptor membrane components

## ABSTRACT

A previous investigation showed that the endometrium normalized in women with endometrial hyperplasia after three months treatment with high dose levonorgestrel IUS (intrauterine system) [1]. The effect was maintained even if immunohistochemical analyses of the endometrium showed that nuclear progesterone receptors (nPRs) were completely downregulated. These observations indicated that some type of non-genomic effect existed [2]. We conducted new investigations of endometrial hyperplasia, now with 6 months low dose levonorgestrel IUS treatment. Again, the growth disturbances were reversed with normalization of the en-

DOI of original article: [10.1016/j.jsbmb.2020.105701](https://doi.org/10.1016/j.jsbmb.2020.105701)

\* Corresponding author.

E-mail address: [georg.sager@uit.no](mailto:georg.sager@uit.no) (G. Sager).

<https://doi.org/10.1016/j.dib.2020.105923>

2352-3409/© 2020 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license.

(<http://creativecommons.org/licenses/by/4.0/>)

ometrium [3,4]. In the context of these studies, RT-qPCR analyses of the endometrium were performed before and after treatment, to determine expression of nuclear progesterone receptors (nPRA+B and nPRB), membrane progesterone receptors (mPR,  $\alpha$ -,  $\beta$ - and  $\gamma$ -subtypes) and progesterone receptor membrane components (PGRMC1 and PGRMC2). The human cervical cell line (C-4 I) [5] with no detectable nPRs [6,7], was included in the investigation as biological control. The gene expression of nPRs, mPRs and PGRMCs was determined in the logarithmic growth phase. Tissue and cellular mRNA was determined with RT-qPCR and used as a surrogate marker for receptor (protein) expression. The present data are connected to the related article entitled "Expression of nuclear progesterone receptors (nPRs), membrane progesterone receptors (mPRs) and progesterone receptor membrane components (PGRMCs) in the human endometrium after 6 months levonorgestrel low dose intrauterine therapy" [8].


© 2020 The Authors. Published by Elsevier Inc.  
This is an open access article under the CC BY license.  
(<http://creativecommons.org/licenses/by/4.0/>)

## Specifications Table


Subject	Molecular Biology
Specific subject area	mRNA and gene expression of progesterone receptors and receptor membrane components
Type of data	Table Chart Graph
How data were acquired	RT-qPCR (Reverse transcription polymerase chain reaction) Instruments: Experion™ electrophoresis system (Bio-Rad, Hercules, CA, USA), NanoDrop 2000c spectrometer (Thermo Scientific, Wilmington, DE, USA), CFX96 real-time PCR detection system (Bio-Rad, Hercules, CA, USA), Software: qBase, geNorm
Data format	Raw and analyzed
Parameters for data collection	Endometrial biopsies from women with endometrial hyperplasia were obtained before and after treatment with levonorgestrel intrauterine device. The human cervical cell line (C-4 I) were expanded and harvested during logarithmic growth. Expression of genes for nuclear progesterone receptors, membrane progesterone receptors and progesterone receptor membrane components were evaluated.
Description of data collection	Total RNA were extracted from human endometrium and human C-4I cells to perform RT-qPCR
Data source location	UiT - The Arctic University of Norway, Tromsø, Norway
Data accessibility	The data are hosted in a public repository (Mendeley Data): <a href="https://data.mendeley.com/datasets/4fwxby52h9/1">https://data.mendeley.com/datasets/4fwxby52h9/1</a>
Related research article	E.T.Sletten, N. Smaglyukova, A.Ørbo and G.Sager, Expression of nuclear progesterone receptors (nPR), membrane progesterone receptors (mPR) and progesterone receptor membrane components (PGRMC) in the human endometrium after 6 months levonorgestrel low dose intrauterine therapy, Journal of Steroid Biochemistry and Molecular Biology <a href="https://doi.org/10.1016/j.jsbmb.2020.105701">https://doi.org/10.1016/j.jsbmb.2020.105701</a>

## Value of the Data

- The regulation of receptors that activate signal pathways of progesterone and gestagens are incompletely understood
- The present data will be useful for scientists in the basic and clinical research of progesterone biochemistry, physiology and pharmacology



**Fig. 1.** Experimental design for RT-qPCR analysis of nPR (nuclear progesterone receptors), mPr (membrane progesterone receptors) and PGRMC (progesterone receptors membrane components) gene expression (mRNA).



**Fig. 2.** Stability of reference genes tested for endometrial biopsies. B2M (Beta-2-Microglobulin), POLR2A (RNA Polymerase II Subunit A), GAPDH (Glyceraldehyde-3-Phosphate Dehydrogenase), HPRT1 (hypoxanthine phosphoribosyltransferase 1) and PKG1 (protein kinase cGMP-dependent 1).


- The present data represent a potential tool for scientists to characterize the complex interplay between nuclear progesterone receptors (nPRs), membrane progesterone receptors (mPRs) and progesterone receptor membrane components (PGRMCs) [9,10]
- The data can lead to development of new experiments to extend the knowledge of the regulation of nuclear as well as membrane progesterone receptors under various physiological conditions and under influence of pharmacological substances *in vivo* and *in vitro*

## 1. Data Description

The present data are a result of the investigation of nPR, mPR and PGRMC gene expression in human endometrium and in a human cancer cell line from the uterine cervix (C-41) and obtained according to the flowchart in Fig. 1. Table 1 shows the gene symbols, NCBI genBank accession numbers gene names, primer sequences, amplicon size and PCR efficiency. Fig. 2 shows the difference in stability of reference genes tested for the analysis of endometrial biopsies. The order of stability was  $PKG1 \geq HPRT1 \geq GAPDH > POLR2A > B2M$ . The three most stable (GAPDH,

**Table 1**  
List of primers for reference genes employed in qPCR-analysis. For the cultured C-4I cells, a commercially available reference gene primer sets CYC 1, ATP5B, GAPDH, 18S, ACTB and UBC were used (Primerdesign Ltd., Southampton, UK).

Gene symbol	NCBI GenBank Accession No.	Gene Name	Primer sequence 5'–3' (F,R)	Amplicon size (bp)	PCR efficiency%
GAPDH	NM_001289745	Glyceraldehyde-3-phosphate dehydrogenase	5'GAGCGAGATCCCTCCAAAT3' 5'AAATGAGCCCCAGCCTTCT3'	101	90.8
HPRT1	NM_000194	Hypoxanthine phosphoribosyltransferase 1	5'CTAATTATGGACAGGACTGAAC3' 5'AGCAAAGAATTTATAGCCCC3'	108	99.1
PGK1	NM_000291	Phosphoglycerate kinase 1	5'CTAAGCAGATTGTGTGGAATG3' 5'CTCACATGGCTGACTTTATC3'	187	94.9
POLR2A	NM_000937	Polymerase II subunit A	5'GAATACCTTCCACTATGCTG3' 5'AGAATATCCTTGCTCTCTC3'	162	–
B2M	NM_004048	Beta-2-microglobulin	5'TCATCCACCAGCAGAAATGGAA3' 5'TCTGAATGCTCCAGTTTTTCAA3'	126	–
CYC 1	NM_001916	Cytochrome-c1			95.8
ATP5B	NM_001686	ATP synthase, H <sup>+</sup> transporting, mitochondrial F1 complex, beta polypeptide			93.7
GAPDH	NM_001289745	Glyceraldehyde-3-phosphate dehydrogenase			–
18S	X03205	18S ribosomal RNA			–
ACTB	NM_001101	Actin beta			–
UBC	NM_021009	Ubiquitin C			–



**Fig. 3.** Stability of reference genes tested for C-4I cells. UBC (ubiquitin C), ACTB (Actin Beta), 18S (18S ribosomal RNAs), GAPDH (Glyceraldehyde-3-Phosphate Dehydrogenase), CYC 1 (Cytochrome C1) and ATP5B (ATP Synthase F1 Subunit Beta).

HPRT1 and PGK1) were used for normalization (Table 1). Fig. 3 shows the difference in stability of reference genes tested for the analysis of C-4I cells. The order of stability was ATP5B = CYC 1 = GAPDH > 18S > ACTB > UBC. In the analyses CYC 1 and ATP5B were selected as reference genes for normalization control of C-4I cells (Table 1).

Table 2 shows the primers used for detection of PRA+PRB, PRB, mPR $\alpha$ , mPR $\beta$ , mPR $\gamma$ , PGRMC1, PGRMC2 by RT-qPCR, in addition to gene symbols, NCBI genBank accession numbers, gene names, primer sequences, amplicon size and PCR efficiency. The corrected normalized relative quantity (CNRQ) was calculated for each gene and scaled to the average value.

## 2. Experimental design, materials, and methods

### 2.1. Experimental design

Fig. 1 shows a flowchart for the experimental design of RT-qPCR analysis of nPRs (nuclear progesterone receptors), mPRs (membrane progesterone receptors) and PGRMCs (progesterone receptor membrane components) gene expression (mRNA).

### 2.2. Materials

Endometrial biopsies with analyzable tissue samples were obtained from 42 women. A group of women ( $n=61$ ) was recruited to a prospective, multicenter pilot investigation to assess the efficacy of LNG-IUS 13.5 mg (Jaydess<sup>TM</sup>, Bayer Pharmaceuticals, Berlin, Germany) for treatment of endometrial hyperplasia. Only those women ( $n=49$ ) with a completed treatment period of six months were included in the present work. In seven of the 49 women, endometrial biopsy material was insufficient for qPCR analysis. Endometrial biopsies were obtained by an endometrial suction cuvette (Pipelle<sup>TM</sup>, Laboratoire CCD, Paris, France) or dilatation and curettage or hysteroscopic transcervical resection. All endometrial tissue specimens (baseline and post therapy biopsies) were conserved in RNAlater stabilization solution (Ambion RNAlater<sup>TM</sup>, ThermoFisher Scientific, Waltham, MA, USA, code AM7021) and immediately frozen and kept at -18 °C until further analysis. Details of the clinical investigations are reported earlier [3,4].

**Table 2**List of primers used for detection of PRA+PRB, PRB, mPR $\alpha$ , mPR $\beta$ , mPR $\gamma$ , PGRMC1, PGRMC2 by RT-qPCR.

Gene symbol	NCBI GenBank Accession No.	Gene Name	Primer sequence 5'–3' (F,R)	Amplicon size (bp)	PCR efficiency%
PRA+PRB	NM_001202474	Homo sapiens progesterone receptor (PGR), transcript variant 1, mRNA	5'AGGTCTACCCGCCCTATCTC3' 5'TCCCACAGGTAAGGACACCA3'	150	93.5
PRB	NM_000926	Homo sapiens progesterone receptor (PGR), transcript variant 2, mRNA	5'GCACGAGTTTGATGCCAGAGA3' 5'CTGCGACGGCAATTTAGTGA3'	69	98.4
mPR $\alpha$	NM_178,422	Progesterin and adipoQ receptor family member 7 (PAQR7)	5'CGCTCTCTGGAAGCCGTACATCTATG3' 5'CAGACGGTGGTCCAGACATTCAC3'	121	110.6
mPR $\beta$	NM_133,367	Progesterin and adipoQ receptor family member 8 (PAQR8)	5'GTCCATCTGTACGCTCTCCC3' 5'GCAGGCCATGTGGACAGATA3'	106	95.9
mPR $\gamma$	NM_017705	Progesterin and adipoQ receptor family member 5 (PAQR5)	5'CAGCTGTTTCACGTGTGTGATCCTG3' 5'GGACAGAAGTATGGCTCCAGCTATCTGAG3'	144	99.8
PGRMC1	NM_006667	Progesterone receptor membrane component 1	5'TGACCTTCTGACCTCACTGC3' 5'GCCACGTGATGATACTTGA3'	85	93.4
PGRMC2	NM_006320	Progesterone receptor membrane component 2	5'TCGAGAAATGGAAATGCAG3' 5'TTGTGATCCTTGATCTCTTCA3'	111	92.8

The human cell line C-4I (ATCC, CRL-1594<sup>TM</sup>) was derived from a squamous carcinoma of the uterine cervix [5] and obtained from the American Type Culture Collection (Rockville, MD, USA). The cells were cultured in RPMI-1640 (Sigma-Aldrich, St. Louis, MO, USA, code R8758) with 10% (v/v) fetal bovine serum (Sigma-Aldrich, St. Louis, MO, USA, code F7524), 1% Penicillin-Streptomycin with 10,000 units penicillin and 10 mg streptomycin per mL in 0.9% NaCl (Sigma-Aldrich, St. Louis, MO, USA, code P0781) at 37 °C in a humidified 5% CO<sub>2</sub> atmosphere in our laboratory. The cells were seeded in six well plates at a density of  $4 \times 10^4$  cells/mL in five parallels. The medium was changed every second day and the cells were harvested in the logarithmic phase of growth (five days after seeding).

To collect the cultures for RNA extraction, media was removed by aspiration, and the cells were brought into a suspension using 200  $\mu$ L 0.25% (w/v) trypsin with 0.2% (w/v) Na<sub>4</sub>EDTA, (Sigma-Aldrich, St. Louis, MO, USA, code T4049), and the trypsin activity was terminated by addition of 800  $\mu$ L of incubation media. The cell counts were obtained (Contess Automated Cell Counter<sup>TM</sup>, Invitrogen, ThermoFisher Scientific, Carlsbad, CA, USA) and the cells were conserved in RNA stabilization solution (Ambion RNAlater<sup>TM</sup>, ThermoFisher Scientific, Waltham, MA, USA, code AM7021), stored at 4 °C overnight and then transferred to -20 °C until further analysis.

### 2.3. Methods

The first step was RNA isolation and cDNA synthesis. Total RNA extraction from endometrial tissue samples and C-4I cells was performed using a commercial RNA extraction kit (Bio-Rad Aurum Total RNA Mini Kit, BIO-RAD, Hercules, CA, USA, code 732-6820). RNA was isolated according to manufacturer's instruction manual. Tissue samples (10–30 mg) were transferred to MagNA Lyser Green Beads tubes (Roche Molecular Biochemicals, Mannheim, Germany). Lysis buffer (700  $\mu$ L) from RNA isolation kit, containing 1% (v/v)  $\beta$ -mercaptoethanol (BioRad, Hercules, CA, USA, code 1,610,710), was added to the tubes. The samples were homogenized in a Precellys<sup>TM</sup> 24 high-throughput homogenizer (Bertin Technologies, Rockville, MD, USA) at 6000  $\times$  g for 30 s and cooled on ice for 2 min. Afterwards, the samples were left at room temperature for 5 min. RNA was eluted and stored at -70 °C.

RNA quality was evaluated with the Experion<sup>TM</sup> electrophoresis system (Bio-Rad, Hercules, CA, USA) using the Experion<sup>TM</sup> RNA StdSens Analysis Kit (Bio-Rad Laboratories, Hercules, CA USA, code 7,007,104). RNA samples isolated from patient biopsies and from cultured cells were examined for concentration, purity and integrity. The concentrations and the purity of RNA were determined using NanoDrop 2000c spectrometer (Thermo Scientific, Wilmington, DE, USA). The endometrial RNA samples showed A260/280 ratio of  $2.1 \pm 0.023$  (mean  $\pm$  SD) and a RNA integrity of  $8.5 \pm 0.9$  (mean  $\pm$  SD). The A260/280 ratio of the C-4I cell RNA samples was  $2.1 \pm 0.012$  (mean  $\pm$  SD) and with a RNA integrity of  $9.9 \pm 0.1$  (mean  $\pm$  SD).

The concentration of RNA was measured using NanoDrop 2000c spectrophotometer (Thermo Scientific, Wilmington, DE, USA). Reverse transcription was performed using iScript<sup>TM</sup> cDNA Synthesis Kit (Bio-Rad, Hercules, CA, USA, code 170-8891). A total of 500 ng RNA was reverse transcribed in a final volume of 20  $\mu$ L. After the synthesis, cDNA was diluted 10 x with nuclease free water (Promega, Madison, WI, USA, code P1195). The cDNA was stored at -20 °C for further use.

The next step was analysis of reference gene stability, selection of primers and reaction efficiency. The most stable reference genes were chosen for normalization calculations in qBase program. Predesigned reference genes were obtained from Sigma-Aldrich (Haverhill, UK) and Primerdesign Ltd (Southampton, UK). The geNorm program (<https://genorm.cmgg.be>) was used to find the most stable gene transcript of reference genes for normalization controls. In the qPCR analyses, several reference genes belonging to different functional classes were evaluated by the geNorm software. For the endometrial biopsies five reference genes (B2M, POLR2A, GAPDH, HPRT1 and PGK1) were examined. Six genes (UBC, ACTB, 18S, GAPDH, CYC1 and ATP5B) were evaluated as reference genes for C-4I cells. Pairwise variation ( $V_n/n+1$ ) geNorm V analysis was carried out to determine the number of reference genes required for normalization. In the case of C-4I cells, the optimal number was two reference genes, Cyc1 and ATP5B.

PRA+PRB, PRB and mPR $\beta$  primers were designed using Primer-blast on the NCBI website (<http://blast.ncbi.nlm.nih.gov/>). The design of mPR $\alpha$  and mPR $\gamma$  primers was performed as described by Dressing and Thomas [11] and those for PGRMC1 and PGRMC2 as described by Causey et al. [12]. The sequences of primers are listed in Table 2.

A standard curve from a twofold serial dilution of pooled cDNA was made and gave a high RT-qPCR reaction efficiency of the primer pairs, shown in Table 1 and Table 2, respectively. A gene-specific amplification was confirmed by a single peak in the melting curve analysis.

Real-time qPCR was carried out using the CFX96 real-time PCR detection system (Bio-Rad, Hercules, CA, USA) and SsoAdvance Universal SYBR Green Supermix (BIO-RAD, Hercules, CA, USA, code 172–5271) and human SYBRgreen reference gene detection kit. 900 reaction (Primerdesign Ltd., Southampton, UK, code HK-SY-hu-900). The assay was run with 2.5  $\mu$ L of 10  $\times$  diluted cDNA in a 10  $\mu$ L total reaction supermix. The thermal profile for SYBR real-time qPCR included an initial heat-denaturing step at 95  $^{\circ}$ C for 30 s, 40 cycles at 95  $^{\circ}$ C for 5 s and 60  $^{\circ}$ C for 30 s. Following amplification, a melt-curve analyses of the PCR products were performed from 65  $^{\circ}$ C to 95  $^{\circ}$ C to determine the specificity of amplification. Each sample was run in duplicate. A non-template control and an inter-run calibrator were added to each run. Data acquisition and subsequent data analyses were performed using the CFX Manager software (Bio-Rad, Hercules, CA, USA). Gene expression was analyzed with qBase software as described by Hellemans et al. [13]. The program employs a modified delta-Ct method with the possibility to adjust for PCR efficiency and to use multiple reference genes for normalization. The corrected normalized relative quantity (CNRQ) was calculated for each gene and scaled to the average value.

## Declaration of Competing Interest

None.

## Acknowledgment

We thank the Regional Research Board of Northern Norway (Helse Nord, Tromsø, Norway) and UiT – The Arctic University of Norway (Tromsø, Norway) for funding.

## Ethics statement

The current data were a second gain of two clinical investigations [3,4]. The investigations were approved by the Regional Committees for Medical and Health Research Ethics (2015/381) and by the Norwegian Medicines Agency (EUDRACT nr 2015–000612–17). Study protocol is available at EU Clinical Trials Register. Written informed consent was mandatory. Insurance for coverage of pharmaceutical injuries was signed for all study participants.

## References

- [1] A.B. Vereide, T. Kaino, G. Sager, M. Arnes, A. Orbo, Effect of levonorgestrel IUD and oral medroxyprogesterone acetate on glandular and stromal progesterone receptors (PRA and PRB), and estrogen receptors (ER-alpha and ER-beta) in human endometrial hyperplasia, *Gynecol.Oncol.* 101 (2006) 214–223.
- [2] R.M. Losel, E. Falkenstein, M. Feuring, A. Schultz, H.C. Tillmann, K. Rossol-Haseroth, M. Wehling, Nongenomic steroid action: controversies, questions, and answers, *Physiol.Rev.* 83 (2003) 965–1016.
- [3] E.T. Sletten, M. Arnes, A.B. Vereide, A. Orbo, Low-dose LNG-IUS as therapy for Endometrial Hyperplasia. a prospective Cohort pilot study, *Anticancer Res.* 38 (2018) 2883–2889.
- [4] E.T. Sletten, M. Arnes, A.B. Vereide, A. Orbo, Intrauterine Progestin therapy as a new approach to Premalignant Endometrial Polyps: a prospective observational study, *Anticancer Res.* 39 (2019) 4897–4903.
- [5] N. Auersperg, A.P. Hawryluk, Chromosome observations on three epithelial-cell cultures derived from carcinomas of the human cervix, *J.Natl.Cancer Inst.* 28 (1962) 605–627.
- [6] G. Sager, A. Orbo, R. Jaeger, C. Engstrom, Non-genomic effects of progestins-inhibition of cell growth and increased intracellular levels of cyclic nucleotides, *J.Steroid.Biochem.Mol.Biol.* 84 (2003) 1–8.

- [7] E. Berthelsen, P. Endresen, A. Orbo, G. Sager, Non-genomic cell growth inhibition by progesterone. Cell cycle retardation and induction of cell death, *Anticancer Res.* 24 (2004) 3749–3756.
- [8] E.T. Sletten, N. Smaglyukova, A. Orbo, G. Sager, Expression of nuclear progesterone receptors (nPR), membrane progesterone receptors (mPR) and progesterone receptor membrane components (PGRMC) in the human endometrium after 6 months levonorgestrel low dose intrauterine therapy, *J. Steroid.Biochem.Mol.Biol.* 202 (2020) 1057012020.
- [9] B. Gellersen, M.S. Fernandes, J.J. Brosens, Non-genomic progesterone actions in female reproduction, *Hum Reprod Update* 15 (2009) 119–138.
- [10] P. Thomas, Y. Pang, J. Dong, Enhancement of cell surface expression and receptor functions of membrane progesterin receptor alpha (mPRalpha) by progesterone receptor membrane component 1 (PGRMC1): evidence for a role of PGRMC1 as an adaptor protein for steroid receptors, *Endocrinology* 155 (2014) 1107–1119.
- [11] G.E. Dressing, P. Thomas, Identification of membrane progesterin receptors in human breast cancer cell lines and biopsies and their potential involvement in breast cancer, *Steroids* 72 (2007) 111–116.
- [12] M.W. Causey, L.J. Huston, D.M. Harold, C.J. Charaba, D.L. Ippolito, Z.S. Hoffer, T.A. Brown, J.D. Stallings, Transcriptional analysis of novel hormone receptors PGRMC1 and PGRMC2 as potential biomarkers of breast adenocarcinoma staging, *J.Surg.Res.* 171 (2011) 615–622.
- [13] J. Hellemsans, G. Mortier, P.A. De, F. Speleman, J. Vandesompele, qBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data, *Genome Biol.* 8 (2007) <https://doi.org/10.1186/gb-2007-8-2-r19>.