

Influence of reproductive output on divorce rates in polar seabirds

Guillaume Mercier^{1,2} | Nigel G. Yoccoz² | Sébastien Descamps¹ 

¹Norwegian Polar Institute, Fram Centre, Tromsø, Norway

²Department of Arctic and Marine Biology, UiT The Arctic University of Norway, Tromsø, Norway

Correspondence

Sébastien Descamps, Norwegian Polar Institute, Fram Centre, Tromsø 9296, Norway.
Email: sebastien.descamps@npolar.no

Funding information

SEAPOP; MOSJ

Abstract

The high occurrence of social monogamy in birds has led to questions about partner fidelity, or the perennial nature of monogamy from one breeding season to another. Despite the evolutionary advantages of partner fidelity, divorce occurs among 95% of bird species. We aimed to describe patterns of divorce and partner fidelity in five seabird species breeding in Arctic and Antarctic regions and investigated the influence of breeding status on pair bond maintenance. For four out of the five species considered, we observed low divorce rates (respectively 1.9%, 3.3%, 2.5%, and 0.0% for Brünnich's guillemot, glaucous gull, Antarctic petrel, and south polar skua), while the divorce rate was much higher (19.1%) for the black-legged kittiwake. For kittiwakes, the divorce rate was lower for pairs that managed to raise their chick to 15 days of age, while the effect of breeding success on divorce in the four other species could not be tested due to the rareness of divorce events. Our results emphasize the potentially large temporal (interannual) variations that should be taken into account in understanding divorce and partner fidelity in seabirds.

KEYWORDS

Antarctica, breeding status, divorce, partner fidelity, Svalbard

1 | INTRODUCTION

The existence of monogamous mating systems in the animal kingdom has long been a topic of interest in evolutionary and behavioral ecology (Reichard, 2003). Defined as systems with an exclusive social relationship between one adult female and one adult male during a given reproductive event, socially monogamous mating systems are globally poorly represented in animals (Klug, 2018; Kvarnemo, 2018). They are almost absent in invertebrates (Mathews, 2002; McKeown & Shaw, 2008) and occur in few species of amphibians (Gillette et al., 2000; Tumulty et al., 2014), fish (Whiteman & Côte, 2004), and mammals (Lukas & Clutton-Brock, 2013). However, with about 80% of species considered as socially monogamous, birds are exceptions to these global patterns (Black, 1996; Cockburn, 2006).

Pair bonds in birds have been described as typical examples of cooperative behavior in action (Black, 1996; Cockburn, 2006). Social monogamy has been shown to be favored in systems where the sharing of parental care within a pair (egg incubation, chick feeding, and defense), the lack of ability for individuals to sustain multiple partners during a breeding season (in terms of resources and territory) and the occurrence of active defensive behaviors to maintain unique access to a single partner occur (Brotherton & Komers, 2003; Gowaty, 1996; Grønseth, 2018; Klug, 2018; Møller, 2003). Social monogamy is nevertheless not a monolithic term and finds one of its sources of variation in pair bond duration (Gowaty, 1996). The reunion of the two individuals forming a pair from one breeding event to another, called partner fidelity or perennial monogamy, occurs heterogeneously in monogamous bird species (Black, 1996;

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd.

Griffith, 2019). While the ending of a pair bond may logically be induced by the death of one partner (widowing), it can also happen through a divorce when two birds forming a couple are still alive and pair with a new partner. Divorce has been recorded in 95% of socially monogamous bird species (Black, 1996; Choudhury, 1995; Culina et al., 2015; Ens et al., 1996).

From an evolutionary perspective, the maintenance of a pair bond throughout the years may be advantageous in terms of familiarity between mates, and enhancing the coordination and cooperation of the two members of a couple (Black, 1996; Bried & Jouventin, 2002; Choudhury, 1995; Sánchez-Macouzet et al., 2014). Remaining faithful to the same partner also allows individuals to save energetic resources that would otherwise be allocated to obtaining a new mate without the risk of missing a breeding season (Bried & Jouventin, 2002). On the other hand, when a pair bond results in poor breeding success, the fitness benefits of divorce may exceed those of partner fidelity if divorce allows partner(s) to find a better mate and potentially achieve higher breeding success (Black, 1996; Bried & Jouventin, 2002; Choudhury, 1995; McNamara & Forslund, 1996). Therefore, divorce can be considered as an adaptive mechanism to correct for suboptimal partnerships associated with poor reproductive performance, with breeding success being a proxy used by birds to assess their partner's quality. The effect of breeding status on partner fidelity was analyzed in two meta-analyses, based on 35 and 64 bird species, respectively, and an overall significant pattern of pairs with low breeding success having higher divorce rates was found (Culina et al., 2015; Dubois & Cézilly, 2002).

Besides these general patterns observed in meta-analyses, responses can vary greatly among species, depending on life-history traits such as longevity (Dubois et al., 1998; Ens et al., 1996; Jeschke & Kokko, 2008). Long-lived species can indeed capitalize on mate fidelity and take advantage of the familiarity effect within a pair bond (Bried et al., 2003; Bried & Jouventin, 2002; Jeschke & Kokko, 2008) so that divorce is expected to be less beneficial for long-lived species. The ecology of birds can also potentially add other constraints on partner fidelity and divorce patterns. Seabirds, for example, rely on marine resources for their food supplies but breed on land, so they must often travel considerable distances to find food for their chicks. In such systems, sharing parental care and coordination between partners is an important condition for successful reproduction and the choice of the reproductive partner is crucial (Bried & Jouventin, 2002). For birds breeding at high latitudes, the very short breeding season and the need to fulfill the breeding cycle in a shorter period of time can add another constraint on divorce and favor partner fidelity. Indeed, searching for a new mate may delay the initiation of breeding activities, which could be very costly in polar environments where breeding phenology is an important determinant of breeding success (Burr et al., 2016; Ens et al., 1996; Ritz et al., 2005). Furthermore, the familiarity between faithful mates may also be an important factor to achieve a successful breeding and raise offspring in these harsh and stochastic environments (Halimubieke et al., 2020).

The aim of this present study was to describe partner fidelity and divorce patterns of five seabird species breeding in polar environments (Arctic and Antarctic). We predicted low divorce rates for these long-lived species breeding at high latitudes. We then investigated the influence of breeding success on divorce, testing the prediction that divorce should be higher following a breeding failure.


2 | MATERIAL AND METHODS

2.1 | Study system and species

We based our study on five seabird species: the black-legged kittiwake (*Rissa tridactyla*), the Brünnich's guillemot (*Uria lomvia*), the glaucous gull (*Larus hyperboreus*), the Antarctic petrel (*Thalassoica antarctica*), and the south polar skua (*Stercorarius maccormicki*). Data from the first three species were from colonies in two fjords of the high Arctic Svalbard Archipelago (Figure 1). The kittiwake colony was located in Isfjorden (Grumantbyen) at an abandoned human settlement where buildings are used as a nesting ground for approximately 45 pairs. Situated in Kongsfjorden, the Brünnich's guillemot colony (Ossian Sarsfjellet) consisted of a bird cliff used as a nesting ground by approximately 1,000 pairs. Also breeding in Kongsfjorden, glaucous gull nests were distributed across the entire fjord (ca. 100 nests in total in the fjord). The two other species, the Antarctic petrel and the south polar skua, were breeding at Svarthamaren, an ice-free area (nunatak) located ca. 200 km inland in Dronning Maud Land, Antarctica. With between 100,000 and 250,000 breeding pairs, this colony is one of the largest Antarctic petrel colonies (Descamps et al., 2016; Schwaller et al., 2018; Van Franeker et al., 1999). Svarthamaren also hosts 100–150 skua breeding pairs nesting in the lower flat parts, relying exclusively on petrel eggs and chicks as their food resource during the breeding season (Busdieker et al., 2020).

These five species are seabirds, which breed on land and remain at sea during the internuptial period (Coulson, 2011; del Hoyo et al., 1992; Delord et al., 2020; Frederiksen et al., 2012, 2016; Gaston & Jones, 1998; Weimerskirch et al., 2015; Weiser & Gilchrist, 2020). They are all characterized by high adult survival rates and thus long lifespans (adult survival rate: 0.85 for *R. tridactyla*, 0.88 for *U. lomvia*, 0.85 for *L. hyperboreus*, 0.91 for *T. antarctica* and 0.91 for *S. maccormicki*, Anker-Nilssen et al., 2020; Descamps et al., 2016; Fluhr et al., 2017; S. Descamps, unpublished data). Clutch size varies between species but is low (1 egg for Brünnich's guillemot and Antarctic petrel, 1–2 eggs for black-legged kittiwake breeding on Svalbard, 1–2 eggs for south polar skuas, and 1–3 eggs for glaucous gulls). The amount of time individuals allocate to chick parental care fluctuates between ca. two weeks for the Brünnich's guillemot (for female parental care only as males stay longer with their chick), 5 weeks for the south polar skua, 6 weeks for the Antarctic petrel, 5–7 weeks for the black legged kittiwakes, and up to seven weeks for the glaucous gull (del Hoyo et al., 1992). These five species are socially monogamous and partners share breeding duties (nest building, egg brooding, and parental care) until the departure from

FIGURE 1 Study sites in (a) Svalbard, Arctic and (b) Dronning Maud Land, Antarctica. Colonies in Svalbard were located in two fjords: Isfjorden ($78^{\circ}18'N$, $15^{\circ}10'E$) for black-legged kittiwakes and Kongsfjorden ($78^{\circ}56'N$, $12^{\circ}25'E$) for Brünnich's guillemot and glaucous gull. Svarthamaren ($71^{\circ}53'S$, $5^{\circ}10'E$) in Dronning Maud Land was located ca. 200 km inland and was used as a nesting site by the south polar skua and the Antarctic petrel



the colony at the end of the breeding season (del Hoyo et al., 1992), except for Brünnich's guillemot, for which males stay longer with the chicks after they leave the nest at an age of 15–30 days (Gaston & Hipfner, 2020; Young et al., 2013). Males and females may have different wintering strategies (e.g., Bogdanova et al., 2011 for *R. tridactyla*; Frederiksen et al., 2016 for *U. lomvia*), but the consequences on the pair bond are unknown. The high average resighting rates of all species in the study sites indicate high site fidelity and low emigration rates (average resighting rates obtained from capture–mark–recapture data modeling: 0.73 for *L. hyperboreus* and *S. maccormicki*, 0.74 for *R. tridactyla*, 0.79 for *T. antarctica*, and a 0.90 for *U. lomvia*; unpublished data).

2.2 | Divorce data

Colonies were monitored from 3 to 11 years from 2009 to 2020 (Table 1). Fieldwork and data collection took place annually from early or mid-incubation to mid- or late chick rearing (early/mid-June to the end of July/beginning of August in Svalbard and from early December to mid-February at Svarthamaren). Every year, a sample of breeding adults nesting at each colony were caught with a nylon loop attached to a telescopic pole while at the nest and ringed with a metal ring and a coded plastic ring for identification at a distance. Blood or feathers were sampled to determine the sex of the individuals using molecular analyses (details about the sexing procedure

TABLE 1 Study period and number of pairs whose status and breeding success were identified for each species (D, divorce; Fi, fidelity; NA, status not used for the analyses)

	<i>Rissa tridactyla</i>	<i>Uria lomvia</i>	<i>Larus hyperboreus</i>	<i>Stercorarius maccormicki</i>	<i>Thalassoica antarctica</i>
Study period	2009–2020	2011–2020	2011–2020	2011–2014	2011–2014
Years of monitoring	11	9	9	3	3
Number of couples with known pair bond status	118 (76 Fi, 18 D, 24 NA)	175 (152 Fi, 3 D, 20 NA)	32 (29 Fi, 1 D, 2 NA)	43 (38 Fi, 5 NA)	46 (40 Fi, 1 D, 5 NA)
Number of couples with known pair bond and breeding status	89	146	29	37	25

in Harris et al., 2020 and Tarroux et al., 2020). Colonies were visited regularly during the breeding season (every two to four days on average), and during each visit, the identity of birds occupying the nests and the breeding status (number of eggs or chicks, hatching of eggs) were recorded. These observations allowed the determination of the breeding pairs and their breeding status for a sample of nests every year (Table 1). Several visits may have been necessary to identify breeding pairs, which led to annual average breeding success being potentially over-estimated (as pairs failing very early in the season were less likely to be identified and thus to be included in the study). Accordingly, pair bond status from one year to the next was determined and classified as (a) fidelity when two individuals of a pair at year t were observed at least once in the same nest in year $t + 1$; (b) divorce when two individuals of a pair in year t were still alive (observed at the colony) but not forming a pair during year $t + 1$. Furthermore, the hatching success (HS) and chick survival 15 days after hatching (CS15d) were used to reflect the early and late breeding success of a pair, respectively. Due to the difficulty in monitoring the survival of glaucous gull chicks after hatching, only hatching success was used for this species. Ultimately, only pairs for which we were able to retrieve breeding and pair bond status in years t and $t + 1$ were used for testing the effect of divorce on breeding success (Table 1).

2.3 | Statistical analyses

2.3.1 | Influence of breeding status on pair bond status

Statistical analyses were carried out with R software (R Core Team, 2019) using generalized linear mixed models under a Bayesian framework with the *rstanarm* package (Gelman et al., 2020). Our initial goal was to test for a relationship between divorce and breeding success for all five species, but due to the very low number of divorces in most species (see results), we could only test this for the black-legged kittiwake. Partner fidelity from year t to $t + 1$ (response variable) was modeled as a binary variable (0 = fidelity, 1 = divorce) with a binomial distribution and a logit link function. Three variables (fixed effects) reflecting the different stages of breeding were alternatively used to test the effect of breeding status on the response variable: the hatching status (HS, failure, or success), the chick survival 15 days after hatching (CS15d, failure, or success), and the overall breeding (OB, failure or success), that is, the product of HS and CS15d. The breeding success or failure was defined by the presence or absence of at least one egg or chick at the nest. Three other variables were included in the models as random effects, as they structure the study design and potentially influence the response variable. They consist of the year, used as a proxy of average annual abiotic (environmental conditions, climatic fluctuations) and biotic (predation, prey availability) pressures (Botero & Rubenstein, 2012; Christensen-Dalsgaard et al., 2018; Descamps et al., 2015, 2016), the pair identity, assumed to reflect the intrinsic quality of the pairs

(Bried & Jouventin, 2002) and the nest identity, illustrating the potential quality difference among nesting places across the colony (Massaro et al., 2001; Varpe & Tveraa, 2005). In our systems, even though a given pair often used the same nest, this was not always the case and both variables were not equivalent. Models computed were of the following form: $\text{logit}(\text{pair bond status}_{t+1}) = \alpha + \beta \times \text{breeding status}_t + (1|\text{year}_t) + (1|\text{nest}_t) + (1|\text{couple}_t)$, with α corresponding to the intercept (global mean response), β the breeding status effect (HS or CS15d or OB), and $1|\text{year}_t$, $1|\text{nest}_t$, and $1|\text{couple}_t$ the random effects. We used the `stan_glmr` function with five chains of 30,000 iterations for each model (Stan Development Team, 2020). Default weakly informative prior distribution for the Bayesian model was implemented to reduce posterior uncertainty and stabilize computations (Muth et al., 2018). Convergence of chains was assessed following the procedure described by Muth et al. (2018) using the `shinystan` function (Stan Development Team, 2017). The leave-one-out cross-validation (hereafter, LOOIC), an information criterion adapted to a mixed models approach, was used to compare models and determine their relative goodness of fit (Vehtari et al., 2017).

3 | RESULTS

3.1 | Pair bond and breeding status observations

Breeding success of black legged kittiwakes showed large inter-annual variability with a total breeding failure in 2013 (Figure 2). Divorce rates varied among years and divorces were observed in five out of the 11 study years (2014, 2015, 2016, 2017, 2019, 2020; Figure 2). Kittiwake divorce rates ranged from 13.3% to 50% and was equal to 19.1% for the studied population (Figure 2). For the two other species where long-term data were available, Brünnich's guillemot and the glaucous gull, interannual variability in breeding success was low (Figure 2). Hatching success averaged 87% and 93% for these two species, respectively, and 82% of the chicks survived up to 15 days after hatching for Brünnich's guillemot. These two species experienced low divorce rates over the study period (1.9% for Brünnich's guillemot, 3.3% for glaucous gull; Figure 2). Breeding success of the Antarctic petrel varied during the two years of monitoring and only one divorce was observed for this species, leading to a divorce rate of 2.5% (Figure 2). For the south polar skua, breeding success was generally high (96.6% in average) and no divorce was observed during the study period (among 43 pairs; Figure 2).

For black-legged kittiwakes, the proportion of couples divorcing after a breeding failure was higher than after a breeding success (30.8% vs. 7.1% overall breeding; Figure 3). Furthermore, the probability for a breeding pair to divorce decreased the later the breeding failure occurred (30.8% divorce for early breeding failure, 25.5% for late breeding failure, Figure 3). These results were partly supported by our model selection (Table 2). Indeed, the model with the lowest LOOIC included the chick survival variable (CS15d; Table 2) but the difference in LOOIC with the null model was small ($\Delta\text{LOOIC}=0.6$). For the Antarctic petrels and glaucous gulls, the only divorce was

observed for a pair that was successful in the previous year. For Brünnich's guillemot, two of the three divorces occurred after a successful breeding.


4 | DISCUSSION

The high occurrence of social monogamy in birds in comparison to other animals has raised questions about partner fidelity and divorce. In our study, based on five seabird species breeding in polar regions, we assessed divorce occurrences and their connection with previous breeding success. For four out of the five species considered, we observed very low divorce rates (respectively 1.9%, 3.3%, 2.5%, and 0.0% for Brünnich's guillemot, glaucous gull, Antarctic petrel, and south polar skua) while divorce was much higher (19.1%) for the black-legged kittiwake and negatively associated with previous breeding success.




4.1 | Divorce and partner fidelity in seabirds

For Brünnich's guillemot and the Antarctic petrel, our results constitute the first numerical values of divorce rate and partner fidelity. In Brünnich's guillemot, partner fidelity has been suggested to be very high, and thus, the divorce rate very low, as observed in other Alcid species sharing a range of ecology and life-history traits (divorce rates of 9.2%–11.7% for Common guillemots (*Uria aalge*); 5.7% for razorbills (*Alca torda*); and 5.7%–16.0% for Atlantic puffins (*Fratercula arctica*); Ashcroft, 1979; Ens et al., 1996; Gaston & Hipfner, 2020; Harris & Wanless, 1989). For Procellariiforms, to which the Antarctic petrel belongs, generally low divorce rates were also found (median divorce rate of 8.8% for 31 species, Bried et al., 2003). For the three other species, the divorce rates obtained in this study were in line with those observed previously, despite some intraspecific variation (see details below). Indeed, the average divorce rate of the kittiwake on Svalbard (19.1%, this study) was similar those observed in Alaska (19.3%, Hatch et al., 1993), France (26.0%, Naves et al., 2006), and the UK (26.1%, Coulson & Thomas, 1980). In another Svalbard colony (Kongsfjorden), based on 32 pairs over three years, a much higher divorce rate (45.7%) was observed (Angelier et al., 2007). Concerning the glaucous gull, one previous study in Canada had estimated their divorce rate at 9% (Gaston et al., 2009), which is higher but in the same order of magnitude as our observations on Svalbard (3%). For the south polar skuas, the fact that no divorce occurred during our study contrasts with some results obtained earlier (divorce rates of 1.5%, 9.1%, and 15%, Ainley et al., 1990; Bried & Jouventin, 2002; Pietz & Parmelee, 1994). The differences in monitoring effort and numbers of couples identified (e.g., 23 years of monitoring in a colony of 1,000 breeding pairs and a significant ringing effort for Ainley et al., 1990; 34 couples over four breeding seasons for Pietz & Parmelee, 1994) may explain these variations. A possible underestimation of divorce rates in our study might also explain the differences observed compared to other studies. Indeed, in our approach,




Open Access





R. tridactyla






U. lomvia





L. hyperboreus



T. antarctica



S. maccormicki



Year

○ Hatching success
● Chick survival 15 days

Year

Light colours: Fidelity
Dark colours: Divorce

FIGURE 2 Breeding success and pair bond status for five polar seabirds. Left panels: breeding success with average hatching success (HS, open circles) and chick survival 15 days after hatching (CS15d, closed circles). Right panels: number of pairs with known bond and breeding status (n : total number of pairs and DR: average divorce rate). DR was calculated using all individuals with known pair bond status (including those with unknown breeding status, see Table 1). Light colors represent the number of pairs remaining faithful, while dark colors represent the number of pairs that divorced. Influence of breeding status on divorce




FIGURE 3 Influence of breeding status on divorce rates for the black-legged kittiwake (*Rissa tridactyla*). Different variables reporting breeding status of birds are presented in the three panels, hatching status (left panel), CS15d (chick survival 15 days after hatching, middle panel), and the overall breeding (early failure: hatching failure, late failure: hatching success but death of the chick(s) within the first 15 days, overall success: hatching success and chick survival 15 days after hatching; right panel)

a couple was described as divorced when both partners were observed alive at the colony. However, it is possible that in some cases, one of these partners had not been observed at the colony even if present, which could thus have led to an underestimation of the total number of divorces.

Among our studied species, divorce was thus generally low and there was a general tendency to maintain pair bonds through time. This corresponds to general observations in seabirds and in birds in general. Indeed, when comparing our results to the 209 species of monogamous birds for which we were able to retrieve a measure of partner fidelity in the literature, we observed that divorce occurs in 95% of the species, and half of the divorce rates were below 15% (Appendix S1). Large differences between species are, however, observed, ranging from birds repairing with a new partner every season (100% divorce, e.g., great blue and gray herons [*Ardea herodias*] and [*Ardea cinerea*], common house martin [*Delichon urbicum*] to strict partner fidelity (0% divorce, e.g., Eurasian nuthatch [*Sitta europaea*], and common pigeon [*Columba livia*, Appendix S1]). Similar patterns were observed for seabirds specifically, with a median divorce rate of 13.8% (Appendix S1). While phylogeny can be an important driver of these evolutionary patterns, different life-history

and ecological traits emerge in the literature to explain interspecies variations such as longevity (Jeschke & Kokko, 2008), degree of coloniality (Dubois et al., 1998), or whether birds are migratory or resident (Ens et al., 1996). Among our study species, all colonial breeders and migratory birds, a lower adult survival rate (85%) was observed for the black-legged kittiwake, compared to 88 to 91% for the Brünnich's guillemot, Antarctic petrel, and south polar skua, which could partly explain their higher divorce rate (Anker-Nilssen et al., 2020; Descamps et al., 2016; Fluhr et al., 2017; unpublished data). However, the adult survival of Svalbard glaucous gull was the same as kittiwake and its divorce rate was much lower as well (unpublished data). This indicates that the among-species variation in divorce and partner fidelity likely depends on a suite of life-history traits (Culina et al., 2015) and cannot be explained by the species longevity alone.

Genetic monogamy may be important to consider in understanding divorce rates and its among-species variation. Beyond socially monogamous mating systems, birds can indeed be involved in extrapair paternity (Griffith et al., 2002). Such situations could allow individuals to maintain a socially monogamous pair bond, leading then to low divorce rates in the population, while increasing their

Model	Number of parameters	Variables (fixed and random)	Estimates	SD/SE	LOOIC	Δ LOOIC
Null model	3	Intercept	1.9	0.6 SE	80.5	0.6
		1 couple		1.0 SD		
		1 nest		0.9 SD		
		1 year		1.5 SD		
HS	4	Intercept	1.3	1.2 SE	82.5	2.6
		HS	0.8	1.2 SE		
		1 couple		1.2 SD		
		1 nest		0.9 SD		
		1 year		1.4 SD		
CS15d	4	Intercept	1.3	0.6 SE	79.9	0.0
		CS15d	1.8	1.0 SE		
		1 couple		1.0 SD		
		1 nest		0.8 SD		
		1 year		1.1 SD		
OB	4	Intercept	-0.3	1.4 SE	80.6	0.7
		OB	1.0	0.6 SE		
		1 couple		1.1 SD		
		1 nest		0.9 SD		
		1 year		1.1 SD		

The model in bold is the one with the lowest LOOIC.

reproductive values by mating with other partners. However, extra-pair paternities are likely rare for our studied species and represent only a very small amount of the chicks produced (7.0% for *S. mac-cormicki* and *T. antarctica*, Griffith et al., 2002; 0% for *R. tridactyla*, Helfenstein et al., 2004; low occurrence in Larid and Alcid species, Anker-Nilssen et al., 2008, 2010).

4.2 | Temporal and spatial variations in divorce rates

Divorce may be dependent on environmental variations and vary both spatially and temporally. Indeed, in the course of our study, we observed important interannual variations in divorce rate for the species for which we had long-term data (9–11 years for Arctic species). While these variations in divorce rates may represent random changes due to small sample sizes, they could also be attributed to fluctuation in yearly environmental conditions (e.g., weather, Botero & Rubenstein, 2012) or population structure (e.g., sex ratio, Liker et al., 2014). Such interannual variations could explain the differences in divorce rates observed for some species among the different studies (e.g., 19.1% divorce for kittiwakes in our study vs. 45.7% in Angelier et al., 2007). This emphasizes the importance of considering multiple years and ideally multiple sites when studying divorce or partner fidelity in a given species, and divorce rates estimated from a single year and/or single colony (including the divorce rates estimated for Antarctic petrel and south polar skua in our study) should be interpreted with caution.

TABLE 2 Output of the generalized linear mixed models testing the effect of breeding status on divorce for Svalbard black-legged kittiwakes

The breeding environment can also constitute an important factor for bird partnerships. At high latitudes, the phenological window to fulfill breeding duties is shorter, which can eventually affect mate fidelity patterns (Bried & Jouventin, 2002; Burr et al., 2016) and, more specifically, may increase the fitness costs of divorce. Indeed, in polar environments, reproduction must be initiated early enough to produce chicks during the short summers and delaying the onset of breeding may lower breeding success (Descamps, 2019; Descamps et al., 2011; Dunn, 2004; Sauve et al., 2019). As looking for a new partner may take time and delay reproduction (Ens et al., 1996), it may thus entail higher fitness costs at high latitudes as compared to nonpolar environments. This hypothesis could, at least partly, explain why we observed so few divorce events in four of the study species (Brünnich's guillemot, glaucous gull, south polar skua, Antarctic petrel) but additional studies are needed to confirm this.

4.3 | Does breeding failure affect the probability to divorce?

Breeding success is expected to affect divorce rates in birds and the probability of a divorce should increase following a breeding failure (Culina et al., 2015; Dubois & Cézilly, 2002). For black-legged kittiwakes, divorce is a physiologically and energetically costly process (Angelier et al., 2007; Chardine, 1987). Due to the lack of coordination in a newly formed couple, higher baseline corticosterone levels, indicating prolonged stress levels, have been measured for divorced

individuals (Angelier et al., 2007). Moreover, newly formed couples undergo a diminution of time off-duty and an augmentation of reproductive effort (increase in greeting ceremonies within the couple, longer copulations) (Chardine, 1987). Such an increase in reproductive effort occurs at the cost of other activities (e.g., feeding activities, maintenance of body conditions), which may ultimately impact individual fitness. Accordingly, our results indicated that the divorce rate was 17% higher following a breeding failure (i.e., failed hatching or death of the chick within the first 15 days) than following a breeding success in Svalbard black-legged kittiwakes, though the statistical support for such an increase was not strong. This increase was relatively low compared to other studies on black-legged kittiwakes that showed an increase of 34% (Cap Sizun, France, Naves, 2005) and 32% (Shiefields, UK, Coulson, 1972) in divorce rate after a breeding failure. The low size of our study colony (45 breeding pairs) possibly restricted the probability of repairing after divorce and consequently inhibited divorce mechanisms (i.e., the fitness cost of divorcing would be too high considering the low probability to find a new partner). The breeding failures observed in this colony may also have been caused by different factors (e.g., nest falling from the ledge, predation by glaucous gulls, starvation), not all necessarily associated with bird quality. It may thus be important to assess the drivers of breeding failures to better understand the relationships between breeding output and divorce. Finally, the age-structure in our study colony may also differ from those in these other kittiwake colonies (Cap Sizun, France, Naves, 2005 and Shiefields, UK, Coulson, 1972). Age is known to affect breeding success and divorce (Ens et al., 1996), but was unfortunately unknown for the individuals included in our study.

5 | CONCLUSION

Our results complement previous studies by studying mating systems and partner fidelity for species in polar environments. While divorce was rare in four of the species studied, it was more frequent for black-legged kittiwakes and was higher after a breeding failure. Moreover, we found large interannual variations in pair bond status, stressing the importance of multi-year studies and the caution needed when interpreting results from short-term studies. In Arctic and Antarctic regions, the environment is rapidly changing (Barros et al., 2014; van der Bilt et al., 2019), but what this means for partner fidelity and divorce patterns remains to be elucidated.

ACKNOWLEDGMENTS

This research was funded by the programs SEAPOP (<http://www.seapop.no/en/>) and MOSJ (<http://www.mosj.no/en/>). We thank Harald Steen for initiating the monitoring of black-legged kittiwakes at Grumantbyen, Connie Sullivan for editing the English and all the field workers that helped in collecting the data, in particular Delphin Ruché, Saga Svavarsdóttir, Iñigo López Sarasa, Antonio Vilches, and Benjamin Merkel for their long-term involvement.

CONFLICT OF INTEREST

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

Guillaume Mercier: Conceptualization (equal); Formal analysis (lead); Methodology (equal); Writing-original draft (lead). **Nigel G. Yoccoz:** Formal analysis (equal); Methodology (equal); Supervision (supporting); Validation (equal); Writing-review & editing (supporting). **Sébastien Descamps:** Conceptualization (lead); Funding acquisition (lead); Project administration (lead); Supervision (lead); Writing-original draft (supporting); Writing-review & editing (equal).

DATA AVAILABILITY STATEMENT

The data associated with this manuscript are available from the Norwegian Polar Institute Datacenter (<https://data.npolar.no/dataset/d506b9fa-5bf4-4b51-8f83-1589018ec311>, <https://doi.org/10.21334/npolar.2021.d506b9fa>).

ORCID

Sébastien Descamps  <https://orcid.org/0000-0003-0590-9013>

REFERENCES

- Ainley, D. G., Ribic, C. A., & Wood, R. C. (1990). A demographic study of the south polar skua *Catharacta maccormicki* at Cape Crozier. *The Journal of Animal Ecology*, 59(1), 1. <https://doi.org/10.2307/5155>
- Angelier, F., Moe, B., Clement-Chastel, C., Bech, C., & Chastel, O. (2007). Corticosterone levels in relation to change of mate in Black-legged kittiwakes. *The Condor*, 109, 668–674.
- Anker-Nilssen, T., Barrett, R., Christensen-Dalsgaard, S., Dehnhard, N., Descamps, S., Systad, G. H. R., Moe, B., Reiertsen, T. K., Bustnes, J. O., Erikstad, K.-E., Follestad, A., Hanssen, S. A., Langset, M., Lorentsen, S.-H., Lorentzen, E., & Strøm, H. (2020). *Key-site monitoring in Norway 2019, including Svalbard and Jan Mayen*. SEAPOP Short Report 1-2020: p. 15.
- Anker-Nilssen, T., Kleven, O., Aarvak, T., & Lifjeld, J. T. (2008). No evidence of extra-pair paternity in the Atlantic Puffin *Fratercula arctica*. *Ibis*, 150(3), 619–622. <https://doi.org/10.1111/j.1474-919X.2008.00810.x>
- Anker-Nilssen, T., Kleven, O., Aarvak, T., & Lifjeld, J. (2010). Low or no occurrence of extra-pair paternity in the Black Guillemot *Cephus grylle*. *Journal of Ornithology*, 151, 247–250. <https://doi.org/10.1007/s10336-009-0450-y>
- Ashcroft, R. E. (1979). Survival rates and breeding biology of puffins on Skomer Island, Wales. *Ornis Scandinavica*, 10, 100–110. <https://doi.org/10.2307/3676349>
- Barros, V. R., Field, C. B., Dokken, D. J., Mastrandrea, M. D., Mach, K. J., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y. O., & Genova, R. C. (2014). Polar Regions. In B. Part (Ed.), *Climate change 2014: Impacts, adaptation, and vulnerability. Part B: Regional aspects. Contribution of working group II to the fifth assessment report of the intergovernmental panel of climate change* (pp. 1567–1612). Cambridge, UK: New York, NY: Cambridge University Press.
- Black, J. M. (1996). Introduction: Pair bonds and partnerships. In J. M. Black (Ed.), *Partnerships in birds: The study of monogamy* (pp. 3–20). Oxford University Press.
- Bogdanova, M. I., Daunt, F., Newell, M., Phillips, R. A., Harris, M. P., & Wanless, S. (2011). Seasonal interactions in the black-legged kittiwake, *Rissa tridactyla*: Links between breeding performance and winter distribution. *Proceedings of the Royal Society B: Biological Sciences*, 278(1717), 2412–2418.

- Botero, C. A., & Rubenstein, D. R. (2012). Fluctuating environments, sexual selection and the evolution of flexible mate choice in birds. *PLoS One*, 7(2), e32311. <https://doi.org/10.1371/journal.pone.0032311>
- Bried, J., & Jouventin, P. (2002). Site and mate choice in seabirds: An evolutionary approach. In E. A. Schreiber & J. Burger (Eds.), *Biology of marine birds* (Chapter 9, p. 44). CRC Press.
- Bried, J., Pontier, D., & Jouventin, P. (2003). Mate fidelity in monogamous birds: A re-examination of the Procellariiformes. *Animal Behaviour*, 65(1), 235–246. <https://doi.org/10.1006/anbe.2002.2045>
- Brotherton, P. N. M., & Komers, P. N. (2003). Mate guarding and the social evolution of social monogamy in mammals. In *Monogamy: Mating strategies and partnerships in birds, humans, and other mammals* (p. 267). Cambridge University Press.
- Burr, Z. M., Varpe, Ø., Anker-Nilssen, T., Erikstad, K. E., Descamps, S., Barrett, R. T., Bech, C., Christensen-Dalsgaard, S., Lorentsen, S.-H., Moe, B., Reiertsen, T. K., & Strøm, H. (2016). Later at higher latitudes: Large-scale variability in seabird breeding timing and synchronicity. *Ecosphere*, 7(5), e01283. <https://doi.org/10.1002/ecs2.1283>
- Busdieker, K. M., Patrick, S. C., Trevail, A. M., & Descamps, S. (2020). Prey density affects predator foraging strategy in an Antarctic ecosystem. *Ecology and Evolution*, 10(1), 350–359. <https://doi.org/10.1002/ece3.5899>
- Chardine, J. W. (1987). The influence of pair-status on the breeding behaviour of the Kittiwake *Rissa tridactyla* before egg laying. *Ibis*, 129, 515–526.
- Choudhury, S. (1995). Divorce in birds: A review of the hypotheses. *Animal Behaviour*, 50(2), 413–429. <https://doi.org/10.1006/anbe.1995.0256>
- Christensen-Dalsgaard, S., May, R., Barrett, R., Langset, M., Sandercock, B., & Lorentsen, S. (2018). Prevailing weather conditions and diet composition affect chick growth and survival in the black-legged kittiwake. *Marine Ecology Progress Series*, 604, 237–249. <https://doi.org/10.3354/meps12744>
- Cockburn, A. (2006). Prevalence of different modes of parental care in birds. *Proceedings of the Royal Society B: Biological Sciences*, 273(1592), 1375–1383. <https://doi.org/10.1098/rspb.2005.3458>
- Coulson, J. C. (1972). The significance of the pair-bond in the kittiwake. In *Proceedings of the International Ornithological Congress* (Vol. 15, pp. 424–433).
- Coulson, J. C. (2011). *The Kittiwake*. Poyser.
- Coulson, J. C., & Thomas, C. (1980). A study of the factors influencing the duration of the pair-bond in the Kittiwake Gull *Rissa tridactyla*. In *17th Int. Ornithol. Congr.* (pp. 822–833).
- Culina, A., Radersma, R., & Sheldon, B. C. (2015). Trading up: The fitness consequences of divorce in monogamous birds. *Biological Reviews*, 90(4), 1015–1034. <https://doi.org/10.1111/brv.12143>
- del Hoyo, J., Elliott, A., Sargatal, J., & Cabot, J. (Eds.) (1992). *Handbook of the birds of the world*. Lynx Edicions.
- Delord, K., Kato, A., Tarroux, A., Orgeret, F., Cotté, C., Ropert-Coudert, Y., Cherel, Y., & Descamps, S. (2020). Antarctic petrels 'on the ice rocks': Wintering strategy of an Antarctic seabird. *Royal Society Open Science*, 7(4), 191429. <https://doi.org/10.1098/rsos.191429>
- Descamps, S. (2019). Breeding synchrony and predator specialization: A test of the predator swamping hypothesis in seabirds. *Ecology and Evolution*, 9(3), 1431–1436. <https://doi.org/10.1002/ece3.4863>
- Descamps, S., Forbes, M. R., Gilchrist, H. G., Love, O. P., & Bêty, J. (2011). Avian cholera, post-hatching survival and selection on hatch characteristics in a long-lived bird, the common eider *Somateria mollissima*. *Journal of Avian Biology*, 42(1), 39–48. <https://doi.org/10.1111/j.1600-048X.2010.05196.x>
- Descamps, S., Tarroux, A., Lorentsen, S.-H., Love, O. P., Varpe, Ø., & Yoccoz, N. G. (2016). Large-scale oceanographic fluctuations drive Antarctic petrel survival and reproduction. *Ecography*, 39(5), 496–505. <https://doi.org/10.1111/ecog.01659>
- Descamps, S., Tarroux, A., Varpe, Ø., Yoccoz, N. G., Tveraa, T., & Lorentsen, S.-H. (2015). Demographic effects of extreme weather events: Snow storms, breeding success, and population growth rate in a long-lived Antarctic seabird. *Ecology and Evolution*, 5(2), 314–325. <https://doi.org/10.1002/ece3.1357>
- Dubois, F., & Cézilly, F. (2002). Breeding success and mate retention in birds: A meta-analysis. *Behavioral Ecology and Sociobiology*, 52(5), 357–364. <https://doi.org/10.1007/s00265-002-0521-z>
- Dubois, F., Cézilly, F., & Pagel, M. (1998). Mate fidelity and coloniality in waterbirds: A comparative analysis. *Oecologia*, 116(3), 433–440. <https://doi.org/10.1007/s004420050607>
- Dunn, P. (2004). Breeding dates and reproductive performance. In A. Møller, W. Fiedler, & P. Berthold (Eds.), *Birds and climate change* (Vol. 35, p. 69–87). Academic Press. [https://doi.org/10.1016/S0065-2504\(04\)35004-X](https://doi.org/10.1016/S0065-2504(04)35004-X)
- Ens, B. J., Choudhury, S., & Black, J. M. (1996). Mate fidelity and divorce in monogamous birds. In *Partnerships in birds: The study of monogamy* (pp. 345–394). Oxford University Press.
- Fluhr, J., Strøm, H., Pradel, R., Duriez, O., Beaugrand, G., & Descamps, S. (2017). Weakening of the subpolar gyre as a key driver of North Atlantic seabird demography: A case study with Brünnich's guillemots in Svalbard. *Marine Ecology Progress Series*, 563, 1–11. <https://doi.org/10.3354/meps11982>
- Frederiksen, M., Descamps, S., Erikstad, K. E., Gaston, A. J., Gilchrist, H. G., Grémillet, D., Johansen, K. L., Kolbeinsson, Y., Linnebjerg, J. F., Mallory, M. L., McFarlane Tranquilla, L. A., Merkel, F. R., Montevecchi, W. A., Mosbech, A., Reiertsen, T. K., Robertson, G. J., Steen, H., Strøm, H., & Thórarinnsson, T. L. (2016). Migration and wintering of a declining seabird, the thick-billed murre *Uria lomvia*, on an ocean basin scale: Conservation implications. *Biological Conservation*, 200, 26–35. <https://doi.org/10.1016/j.biocon.2016.05.011>
- Frederiksen, M., Moe, B., Daunt, F., Phillips, R. A., Barrett, R. T., Bogdanova, M. I., Boulinier, T., Chardine, J. W., Chastel, O., Chivers, L. S., Christensen-Dalsgaard, S., Clément-Chastel, C., Colhoun, K., Freeman, R., Gaston, A. J., González-Solis, J., Goutte, A., Grémillet, D., Guilford, T., ... Anker-Nilssen, T. (2012). Multicolony tracking reveals the winter distribution of a pelagic seabird on an ocean basin scale: Winter distribution of Atlantic kittiwakes. *Diversity and Distributions*, 18(6), 530–542. <https://doi.org/10.1111/j.1472-4642.2011.00864.x>
- Gaston, A. J., Descamps, S., & Gilchrist, H. G. (2009). Reproduction and survival of Glaucous Gulls breeding in an Arctic seabird colony. *Journal of Field Ornithology*, 80, 135–145.
- Gaston, A. J., & Hipfner, J. M. (2020). Thick-billed Murre (*Uria lomvia*), version 1.0. In S. M. Billerman (Ed.), *Birds of the World*. Cornell Lab of Ornithology. <https://doi.org/10.2173/bow.thbmur.01>
- Gaston, A. J., & Jones, I. L. (1998). *The auks: Alcidae*. Oxford University Press.
- Gelman, A., Hill, J., & Vehtari, A. (2020). *Regression and other stories*. Cambridge University Press.
- Gillette, J. R., Jaeger, R. G., & Peterson, M. G. (2000). Social monogamy in a territorial salamander. *Animal Behaviour*, 59(6), 1241–1250. <https://doi.org/10.1006/anbe.2000.1437>
- Gowaty, P. A. (1996). Battles of the sexes and origins of monogamy. In J. M. Black (Ed.), *Partnerships in birds: The study of monogamy* (pp. 21–52). Oxford University Press.
- Griffith, S. C. (2019). Cooperation and coordination in socially monogamous birds: Moving away from a focus on sexual conflict. *Frontiers in Ecology and Evolution*, 7, 455. <https://doi.org/10.3389/fevo.2019.00455>
- Griffith, S. C., Owens, I. P. F., & Thuman, K. A. (2002). Extra pair paternity in birds: A review of interspecific variation and adaptive function. *Molecular Ecology*, 11(11), 2195–2212. <https://doi.org/10.1046/j.1365-294X.2002.01613.x>

- Grønstøl, G. (2018). Polygyny threshold model. In I. J. Vonk, & T. Shackelford (Eds.), *Encyclopedia of animal cognition and behavior* (pp. 1–8). Springer International Publishing. https://doi.org/10.1007/978-3-319-47829-6_460-1
- Halimubieke, N., Kupán, K., Valdebenito, J. O., Kubelka, V., Carmona-Isunza, M. C., Burgas, D., Catlin, D., St Clair, J. J. H., Cohen, J., Figuerola, J., Yasué, M., Johnson, M., Mencarelli, M., Cruz-López, M., Stantial, M., Weston, M. A., Lloyd, P., Que, P., Montalvo, T., ... Székely, T. (2020). Successful breeding predicts divorce in plovers. *Scientific Reports*, *10*(1), 15576. <https://doi.org/10.1038/s41598-020-72521-6>
- Harris, M. P., & Wanless, S. (1989). The breeding biology of Razorbills *Alca torda* on the Isle of May. *Bird Study*, *36*(2), 105–114. <https://doi.org/10.1080/00063658909477012>
- Harris, S. M., Descamps, S., Sneddon, L. U., Bertrand, P., Chastel, O., & Patrick, S. C. (2020). Personality predicts foraging site fidelity and trip repeatability in a marine predator. *Journal of Animal Ecology*, *89*(1), 68–79. <https://doi.org/10.1111/1365-2656.13106>
- Hatch, S. A., Bay, D. R., & Brian, S. F. (1993). Adult survival of Black-legged Kittiwakes *Rissa tridactyla* in a Pacific colony. *Ibis*, *135*(3), 247–254. <https://doi.org/10.1111/j.1474-919X.1993.tb02841.x>
- Helfenstein, F., Tirard, C., Danchin, E., & Wagner, R. H. (2004). Low frequency of extra-pair paternity and high frequency of adoption in black-legged kittiwakes. *The Condor*, *106*(1), 149. <https://doi.org/10.1650/7337>
- Jeschke, J. M., & Kokko, H. (2008). Mortality and other determinants of bird divorce rate. *Behavioral Ecology and Sociobiology*, *63*(1), 1–9. <https://doi.org/10.1007/s00265-008-0646-9>
- Klug, H. (2018). Why monogamy? A review of potential ultimate drivers. *Frontiers in Ecology and Evolution*, *6*, 30. <https://doi.org/10.3389/fevo.2018.00030>
- Kvarnemo, C. (2018). Why do some animals mate with one partner rather than many? A review of causes and consequences of monogamy. *Biological Reviews*, *93*(4), 1795–1812. <https://doi.org/10.1111/brv.12421>
- Liker, A., Freckleton, R. P., & Székely, T. (2014). Divorce and infidelity are associated with skewed adult sex ratios in birds. *Current Biology*, *24*(8), 880–884. <https://doi.org/10.1016/j.cub.2014.02.059>
- Lukas, D., & Clutton-Brock, T. H. (2013). The evolution of social monogamy in mammals. *Science*, *341*(6145), 526–530. <https://doi.org/10.1126/science.1238677>
- Massaro, M., Chardine, J. W., & Jones, I. L. (2001). Relationships between black-legged kittiwake nest-site characteristics and susceptibility to predation by large gulls. *The Condor*, *103*(4), 10. <https://doi.org/10.1093/condor/103.4.793>
- Mathews, L. M. (2002). Territorial cooperation and social monogamy: Factors affecting intersexual behaviours in pair-living snapping shrimp. *Animal Behaviour*, *63*(4), 767–777. <https://doi.org/10.1006/maeb.2001.1976>
- McKeown, N., & Shaw, P. (2008). Single paternity within broods of the brown crab *Cancer pagurus*: A highly fecund species with long-term sperm storage. *Marine Ecology Progress Series*, *368*, 209–215. <https://doi.org/10.3354/meps07634>
- McNamara, J. M., & Forslund, P. (1996). Divorce rates in birds: Predictions from an optimization model. *The American Naturalist*, *147*(4), 609–640.
- Møller, A. P. (2003). The evolution of monogamy: Mating relationships, parental care and sexual selection. In *Monogamy: Mating strategies and partnerships in birds, humans and other mammals* (pp. 29–41).
- Muth, C., Oravec, Z., & Gabry, J. (2018). User-friendly Bayesian regression modeling: A tutorial with rstanarm and shinystan. *The Quantitative Methods for Psychology*, *14*(2), 99–119. <https://doi.org/10.20982/tqmp.14.2.p099>
- Naves, L. C. (2005). La fidélité au partenaire : Stratégie ou contrainte? Le rôle de l'hétérogénéité individuelle chez la mouette tridactyle *Rissa tridactyla*. [Ecologie]. Université Pierre et Marie Curie - Paris VI.
- Naves, L. C., Monnat, J.-Y., & Cam, E. (2006). Breeding performance, mate fidelity, and nest site fidelity in a long-lived seabird: Behaving against the current? *Oikos*, *115*(2), 263–276. <https://doi.org/10.1111/j.2006.0030-1299.14883.x>
- Pietz, P. J., & Parmelee, D. F. (1994). Survival, site and mate fidelity in South Polar Skuas *Catharacta maccormicki* at Anvers Island, Antarctica. *Ibis*, *136*(1), 32–38. <https://doi.org/10.1111/j.1474-919X.1994.tb08128.x>
- R Core Team. (2019). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Reichard, U. R. (2003). Monogamy: Past and present. In U. H. Reichard & C. Boesch (Eds.), *Monogamy: Mating strategies and partnerships in birds, humans and other mammals* (pp. 3–25).
- Ritz, M., Hahn, S., & Peter, H.-U. (2005). Factors affecting chick growth in the South Polar Skua (*Catharacta maccormicki*): Food supply, weather and hatching date. *Polar Biology*, *29*, 53–60. <https://doi.org/10.1007/s00300-005-0027-z>
- Sánchez-Macouzet, O., Rodríguez, C., & Drummond, H. (2014). Better stay together: Pair bond duration increases individual fitness independent of age-related variation. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1786), 20132843. <https://doi.org/10.1098/rspb.2013.2843>
- Sauve, D., Divoky, G., & Friesen, V. L. (2019). Phenotypic plasticity or evolutionary change? An examination of the phenological response of an arctic seabird to climate change. *Functional Ecology*, *33*(11), 2180–2190.
- Schwaller, M. R., Lynch, H. J., Tarroux, A., & Prehn, B. (2018). A continent-wide search for Antarctic petrel breeding sites with satellite remote sensing. *Remote Sensing of Environment*, *210*, 444–451. <https://doi.org/10.1016/j.rse.2018.02.071>
- Stan Development Team. (2017). *shinystan: Interactive visual and numerical diagnostics and posterior analysis for Bayesian models* (R package version 2.3.0.) [Computer software]. <http://mc-stan.org/>
- Stan Development Team. (2020). *RStan: The R interface to Stan* (R package version 2.19.3) [Computer software]. <http://mc-stan.org/>
- Tarroux, A., Cherel, Y., Fauchald, P., Kato, A., Love, O. P., Ropert-Coudert, Y., Spreen, G., Varpe, Ø., Weimerskirch, H., Yoccoz, N. G., & Zahn, S. (2020). Foraging tactics in dynamic sea-ice habitats affect individual state in a long-ranging seabird. *Functional Ecology*, *34*(9), 1839–1856.
- Tumulty, J., Morales, V., & Summers, K. (2014). The biparental care hypothesis for the evolution of monogamy: Experimental evidence in an amphibian. *Behavioral Ecology*, *25*(2), 262–270. <https://doi.org/10.1093/beheco/art116>
- van der Bilt, W., Bakke, J., Smedsrud, L. H., Sund, M., Schuler, T., Westermann, S., Wong, W. K., Sandven, S., Simpson, M. J. R., & Skogen, M. D. (2019). *Climate in Svalbard 2100*.
- Van Franeker, J. A., Gavrilov, M., Mehlum, F., Veit, R. R., & Woehler, E. J. (1999). Distribution and abundance of the Antarctic Petrel. *Waterbirds*, *22*, 14–28. <https://doi.org/10.2307/1521989>
- Varpe, Ø., & Tveraa, T. (2005). Chick survival in relation to nest site: Is the Antarctic petrel hiding from its predator? *Polar Biology*, *28*(5), 388–394. <https://doi.org/10.1007/s00300-004-0695-0>
- Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, *27*(5), 1413–1432. <https://doi.org/10.1007/s11222-016-9696-4>
- Weimerskirch, H., Tarroux, A., Chastel, O., Delord, K., Cherel, Y., & Descamps, S. (2015). Population-specific wintering distributions of adult south polar skuas over three oceans. *Marine Ecology Progress Series*, *538*, 229–237. <https://doi.org/10.3354/meps11465>
- Weiser, E., & Gilchrist, H. G. (2020). Glaucous gull (*Larus hyperboreus*), version 1.0. In S. M. Billerman (Ed.), *Bird of the World*. Cornell Lab of Ornithology. <https://doi.org/10.2173/bow.glagul.01>
- Whiteman, E. A., & Côte, I. M. (2004). Monogamy in marine fishes. *Biological Reviews*, *79*(2), 351–375. <https://doi.org/10.1017/S1464793103006304>

Young, R. C., Kitaysky, A. S., Haussmann, M. F., Descamps, S., Orben, R. A., Elliott, K. H., & Gaston, A. J. (2013). Age, sex, and telomere dynamics in a long-lived seabird with male-biased parental care. *PLoS One*, 8(9), e74931. <https://doi.org/10.1371/journal.pone.0074931>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Mercier, G., Yoccoz N. G., & Descamps S. (2021). Influence of reproductive output on divorce rates in polar seabirds. *Ecology and Evolution*, 00, 1–12. <https://doi.org/10.1002/ece3.7775>