

Geophysical Research Letters



RESEARCH LETTER

10.1029/2021GL093103

Key Points:

- Seismic stratigraphic continental rise-to-shelf link from International Ocean Discovery Program (IODP) Expedition 379 drill sites on the Amundsen Sea rise
- Major prograding shelf growth in the Amundsen Sea Embayment by frequent grounded ice advances in the early Pliocene
- Extended period of ice sheet retreat in mid-Pliocene (4.2–3.2 Ma) from buried grounding zone wedges on shelf and core records from rise

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

K. Gohl,
karsten.gohl@awi.de












Citation:

Gohl, K., Uenzelmann-Neben, G., Gille-Petzoldt, J., Hillenbrand, C.-D., Klages, J. P., Bohaty, S. M., et al. (2021). Evidence for a highly dynamic West Antarctic Ice Sheet during the Pliocene. *Geophysical Research Letters*, 48, e2021GL093103. <https://doi.org/10.1029/2021GL093103>

Received 23 FEB 2021

Accepted 18 JUN 2021

Evidence for a Highly Dynamic West Antarctic Ice Sheet During the Pliocene

Karsten Gohl¹ , Gabriele Uenzelmann-Neben¹ , Johanna Gille-Petzoldt¹ , Claus-Dieter Hillenbrand² , Johann P. Klages¹ , Steven M. Bohaty³ , Sandra Passchier⁴ , Thomas Frederichs⁵ , Julia S. Wellner⁶ , Rachel Lamb¹ , German Leitchenkov⁷ , and IODP Expedition 379 Scientists⁸

¹Alfred Wegener Institute Helmholtz-Center for Polar and Marine Research, Division of Geosciences, Bremerhaven, Germany, ²British Antarctic Survey, Cambridge, UK, ³National Oceanography Centre Southampton, University of Southampton, Southampton, UK, ⁴Department of Earth and Environmental Studies, Montclair State University, Montclair, NJ, USA, ⁵Department of Geosciences, University of Bremen, Bremen, Germany, ⁶Department of Earth and Atmospheric Sciences, University of Houston, Houston, TX, USA, ⁷VNIIOkeangeologia and St. Petersburg State University, St. Petersburg, Russia, ⁸See Appendix A

Abstract Major ice loss in the Amundsen Sea sector of the West Antarctic Ice Sheet (WAIS) is hypothesized to have triggered ice sheet collapses during past warm periods such as those in the Pliocene. International Ocean Discovery Program (IODP) Expedition 379 recovered continuous late Miocene to Holocene sediments from a sediment drift on the continental rise, allowing assessment of sedimentation processes in response to climate cycles and trends since the late Miocene. Via seismic correlation to the shelf, we interpret massive prograding sequences that extended the outer shelf by 80 km during the Pliocene through frequent advances of grounded ice. Buried grounding zone wedges indicate prolonged periods of ice-sheet retreat, or even collapse, during an extended mid-Pliocene warm period from ~4.2–3.2 Ma inferred from Expedition 379 records. These results indicate that the WAIS was highly dynamic during the Pliocene and major retreat events may have occurred along the Amundsen Sea margin.

Plain Language Summary Collapses of the West Antarctic Ice Sheet (WAIS) during past warm times are suggested to begin in the Amundsen Sea sector. During a drilling expedition of the International Ocean Discovery Program (IODP), deep-sea sediments were retrieved from the Amundsen Sea. These sediment cores contain records of colder and warmer periods in the Pliocene (5.3–2.6 million years ago) which relate to the behavior of the WAIS. By analyzing seismic images that allow correlation of sediment layers from the drill sites to the continental shelf, we show that the shelf grew oceanward by 80 km due to the erosion of sediments below the WAIS and their deposition at the shelf break as the result of frequent Pliocene advances of grounded ice across the shelf. The shelf sediment layers contain grounding zone wedges predominantly formed during ice sheet retreat. The preservation of these grounding zone wedges testifies that they were not eroded by subsequent ice advances, with their burial requiring periods of prolonged WAIS retreat during warm intervals. This interpretation is consistent with our IODP drill core observations of a prominent decrease in terrigenous sedimentation between 4.2 and 3.2 million years ago, which indicates a highly dynamic WAIS during the Pliocene.

1. Introduction

Warm periods in the Pliocene (5.3–2.6 Ma) are considered as analogs to present and near-future climate conditions with atmospheric carbon dioxide concentrations of 350–450 ppm, global mean temperatures 2–3°C higher compared to pre-industrial levels, and a global mean sea level 10–30 m higher than present (e.g., Dutton et al., 2015; Gasson et al., 2016; Pagani et al., 2010; Westerhold et al., 2020). The behavior of Antarctic ice sheets during such warm phases is important, as the ice sheets modulate the intensity of global sea level changes. Ice sheet models indicate a partial to complete loss of marine-based sectors of both West and East Antarctic ice sheets during warm Plio- and Pleistocene super interglacials, when incursions of warm deep water into sub-ice shelf cavities near grounding zones caused melting of the buttressing ice shelves (de Boer et al., 2015; Gollledge et al., 2017; Pollard & DeConto, 2020). The Amundsen Sea Embayment (ASE) in West Antarctica (Figure 1) is a key region for monitoring ice-sheet stability, since ice on this

© 2021. The Authors.

This is an open access article under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.




Figure 1. Amundsen Sea sector with International Ocean Discovery Program (IODP) Expedition 379 drill sites on Resolution Drift (red dots) and seismic lines (black lines). Seismic lines used in this study are in orange and yellow. Thwaites (TG) and Pine Island glaciers (PIG) formed Pine Island Trough (PIT) which divides into Pine Island Trough West (PITW) and Pine Island Trough East (PITE) on the outer shelf. Other glacial troughs are Dotson-Getz Trough (DGT) and Abbot-Cosgrove Trough (ACT). White dot marks DSDP Leg 35 Site 324. Bathymetry is from Arndt et al. (2013). Present ice divides (black lines) and flow pattern (blue arrows) are simplified from Rignot et al. (2011). Abbreviations denote West Antarctica (WA), East Antarctica (EA), Amundsen Sea Embayment (ASE), Antarctic Peninsula (AP), Weddell Sea (WS), Ross Sea (RS), Prydz Bay (PB), and Transantarctic Mountains (TAM).

margin is currently being lost at an accelerated pace, leading to rapid retreat of Pine Island, Thwaites, and neighboring glaciers (e.g., Rignot et al., 2019; Scambos et al., 2017; Smith et al., 2020). The Amundsen Sea sector is hypothesized as a precursor for major West Antarctic Ice Sheet (WAIS) retreat or even collapse during most intense warm periods of the Plio-Pleistocene over the past 5 Myr (DeConto & Pollard, 2016).

In order to address past ice-sheet dynamic processes in the Amundsen Sea sector, we analyzed seismic lines across the ASE shelf with direct connection to the two drill sites of the International Ocean Discovery Program (IODP) Expedition 379 (Gohl et al., 2021) on the continental rise (Figure 1). The ASE shelf is characterized by large cross-shelf paleo-ice stream troughs, such as Pine Island Trough (PIT) with its outer shelf western (PITW) and eastern (PITE) branches, Dotson-Getz Trough (DGT), and Abbot-Cosgrove Trough (ACT). The outer shelf shows sequences of prograding deposits, which have been identified at several locations around Antarctica and are characteristic for polar continental shelves (e.g., Anderson & Bartek, 1992; Bart & De Santis, 2012; Dowdeswell et al., 2006; Hochmuth & Gohl, 2019; Nitsche et al., 2000). Antarctic ice sheet advances across the shelves during glacial times added substantial volumes of glacially eroded sediments to the outer shelf, thereby expanding the continental shelf on average by about 7% since the onset of major glaciations (Hochmuth & Gohl, 2019). The amount of glacially induced shelf growth, however, varies

strongly along the Antarctic continental margins as it relates to the location of major outlet glacier sectors and is affected by the regional tectonic architecture (Hochmuth & Gohl, 2019). Outer shelf progradation caused a ~25% shelf growth in the ASE (Gohl et al., 2013; Hochmuth & Gohl, 2019).

IODP Expedition 379 drill sites U1532 and U1533 are located on an elongated sediment drift, informally named “Resolution Drift,” stretching ~300 km northward from the base of the continental slope to the abyssal plain (Figure 1 and S1). This drift and neighboring parallel drifts are flanked by deep-sea channels (Figure S2) that, predominantly during glacial periods, acted as pathways for detritus transported from the shelf down to the slope and then further across the continental rise toward the deep sea. On the continental rise, particles still in suspension were captured by eastward flowing bottom currents before they accumulated on the eastern channel flanks to form the drifts (Dowdeswell et al., 2006; Nitsche et al., 2000). Such sediment drifts are frequently observed along the Pacific margin of Antarctica (e.g., Rebesco et al., 2002; Uenzelmann-Neben & Gohl, 2012, 2014). In this type of depositional system, the highest supply of detritus occurred during glacial periods, when the grounded WAIS bulldozed large volumes of glacial debris to the outer shelf, from where it was redeposited down to the rise. In contrast, interglacial periods of ice-sheet retreat have a reduced time-integrated flux of downslope transported terrigenous material, which then allows enrichment of biogenic components in the drift sediments. The sediments at sites U1532 and U1533 contain unique records to study the cyclicity of WAIS advance and retreat. In particular, Site U1532 contains a sequence of Pliocene sediments with an almost complete paleomagnetic record for constraining very high resolution, suborbital scale climate variability, thus providing an opportunity to better develop climate records for the sparsely sampled pre-Pleistocene interval in the Amundsen Sea (Gohl et al., 2021).

2. Drill Site Description and Seismic Data

Prior to investigation with scientific drilling, previous models of the stratigraphic architecture and ages of shelf and rise sedimentary sequences in the Amundsen Sea (Gohl et al., 2013; Uenzelmann-Neben & Gohl, 2012, 2014) were based only on long distance or jump correlation of regional seismic reflection profiles to the seismic stratigraphy of the Ross Sea tied to Deep Sea Drilling Project (DSDP) Leg 28 sites (e.g., De Santis et al., 1999) as well as to DSDP Leg 35 and Ocean Drilling Program (ODP) Leg 178 sites west of the Antarctic Peninsula (Barker et al., 2002; Tucholke et al., 1976). Recently, shelf sediments in the ASE were drilled with the MARUM-MeBo70 sea-bed drilling device in 2017, recovering sedimentary rocks of Cretaceous to Oligocene-Miocene age (Gohl et al., 2017; Klages et al., 2020). The almost continuous sequences at IODP Expedition 379 Sites U1532 and U1533 on the continental rise (Gohl et al., 2021; Wellner et al., 2021a, 2021b) span the latest Miocene to Holocene with high core recovery rates of 90% and 70%, respectively (Figure 1). Site U1532 is located near the crest of Resolution Drift (Figure S1) and penetrated down to 794 m depth below the seafloor, exhibiting sedimentation rates of up to 61 cm kyr⁻¹ within the Pliocene. Site U1533 penetrated 383 m into the lowermost western flank of this drift, in proximity to a deep-sea channel (Figures S1 and S2), and shows a more condensed sequence with sedimentation rates up to 22 cm kyr⁻¹ in the early Pliocene. Preliminary investigations suggest that sediments at both sites are predominantly terrigenous, but are intercalated by pelagic or hemipelagic deposits that, in some cases, contain microfossils. Chronological control was achieved by both high-resolution magnetostratigraphy and biostratigraphy (Wellner et al., 2021a, 2021b).

Site U1532 is located at the northwestern end of seismic profile AWI-20100130, while Site U1533 is situated on profile TH86003B (Figures 2a and 2b; Figures S1 and S2). We performed correlation of horizons and unit characteristics between the seismic records of the drill sites and the continental shelf through a series of multichannel seismic profiles (Figure 1) (TH86 lines; Scheuer et al., 2006; Yamaguchi et al., 1988; AWI lines: Gohl et al., 2013; Uenzelmann-Neben & Gohl, 2012, 2014). Data acquisition and processing methods for the AWI lines are described in Gohl et al. (2013) and Uenzelmann-Neben and Gohl (2012, 2014). The single-channel seismic line NBP9902-11 on the shelf (Figure S3) displays high-quality data from the seafloor to its first multiple (Lowe & Anderson, 2002). Correlation of major reflectors from the drill sites across the continental rise to the lower slope was relatively straightforward, whereas correlation across the middle and upper slope was more challenging due to the thinning nature of seismic units with increasing slope angle and the presence of slumps and seafloor multiples (Figure S4). Careful reprocessing of line AWI-20100132, however, solved those issues, so that we were able to correlate within an uncertainty of 2–3 reflection wave-




Figure 2. (a) Crossing seismic lines in area of International Ocean Discovery Program (IODP) Expedition 379 Sites U1532 and U1533 near the crest and at the lowermost flank of Resolution Drift, respectively. (b) Seismic transect from Site U1532 on the rise to the outer shelf with interpreted zones of stratigraphic ages and the basement. Shelf unconformities ASS-u2 to ASS-u5 are modified from Gohl et al. (2013). Lines AWI-20100130 and AWI-20100131 in (a) and AWI-20100130 in (b) are rotated for pseudo-3D impression. (c) Section across prograding sequences of outer continental shelf. Circles mark identified shelf breaks in the Pliocene and Pleistocene. bGZW annotates buried grounding zone wedges. Enlarged seismic image shows middle to late Pliocene zone with bGZWs. Vertical black bars mark thicknesses converted from time. TWT is two-way travel-time.

lets up or down resulting in a two-way travel-time uncertainty of ± 100 ms. A further challenge was the extremely strong seafloor multiples in the seismic records from the outer shelf at water depths ≤ 550 m, in particular those of line AWI-20100132 (Figures 2b and 2c). Here, numerous iceberg scours carved into overcompacted seabed sediments (Graham et al., 2010; Lowe & Anderson, 2002). They generated high-amplitude diffractions, which remained present in the multiples despite the long 3000-m streamer, and thus, in conjunction with the relatively small airgun source, prevented a complete suppression of these multiples that are superposed on the lower part of the progradational reflector sequences, despite applying frequency-wavenumber (f-k) and Radon transform filtering.

3. Shelf-to-Rise Sedimentation

The seismic transect for this study is located in the eastern ASE, along the main path of grounded ice advancing from the present termini of Pine Island and Thwaites glaciers toward the outer shelf (Fig. 1; Fig. 2b). The top-of-Pliocene and top-of-Miocene seismic horizons, as identified from drill sites U1532 and U1533 on Resolution Drift (Figures 2a and S1), can be traced across the continental rise, parallel to the elongated drift, and onto the slope and shelf (Figure 2b).

In our re-evaluation of the previous seismic stratigraphic model of the ASE shelf (Gohl et al., 2013), the vast amount of sediment volume deposited within a ~ 80 -km wide zone of prograding foresets was supplied by WAIS advances during the Pliocene (Figures 2b and 2c). Most of the sequences stratigraphically lying below the Plio-Pleistocene units and presumably being of Miocene age were aggradationally deposited, indicating that grounded ice rarely reach the shelf break, while Pleistocene contribution to progradational shelf growth was relatively small compared to that of the Pliocene (cf. Dowdeswell et al., 2006). It should be noted that magnitude of shelf growth varies in the Amundsen Sea (Gohl et al., 2013; Hochmuth & Gohl, 2013; Nitsche et al., 1997), suggesting that tectonic architecture beneath the outer shelf played a significant role for shelf growth (Hochmuth & Gohl, 2019). Seismic lines AWI-20100132 (Figure 2c) and NBP9902-11 (Figure S3) cross the outer shelf just west of the main PITE (Figure 1). Unlike the other glacial troughs (DGT, PITW, ACT), the outer PITE shows a subtle modern and paleo-bathymetric difference (< 100 m) over a




Figure 3. Conceptual model illustrating outer shelf sequences with buried grounding zone wedges (bGZW) that are preserved due to prolonged intervals of sedimentation unaffected by grounded ice advances. GL: grounding line; MSGL: mega-scale glacial lination.

broad outer shelf area (Graham et al., 2010), which indicates that the ice stream widened on the outer shelf (Wellner et al., 2001).

The Pliocene and Pleistocene prograding wedge (Figures 2c and S3) contains distinct paleo-shelf break features from seismic interpretation, indicating a minimum of 14 glacial advances across the outer shelf throughout this period. As grounded ice repeatedly eroded some of the shelf sediments and their depositional shelf-break signature, this is a minimum estimate (cf. Bart et al., 2007), and the realistic number of outer shelf growth events is likely higher. Notably, in the seismic record of the outer shelf, a number of buried grounding zone wedges (bGZWs) is observed at or near the upward doming part of the sequences (Figure 2c). Tracing the bottom and top reflections of the bGZW zone onto the middle shelf south of the oldest onset of progradation, where the aggradational Pliocene layer reaches its full thickness, places this zone approximately into the middle to late Pliocene range. Grounding zone wedges (GZWs) are glaciomorphological depositional features interpreted to result from local accumulation of sediment during a temporary halt in grounding-line retreat of an ice stream (e.g., Batchelor & Dowdeswell, 2015; Klages et al., 2015). Such GZWs are frequently observed on the modern seafloor of polar continental shelves and mark the maximum advance of grounded ice during the last glacial period or phases of grounding-line stillstands during subsequent ice retreat usually within cross-shelf troughs formed by fast-flowing ice streams (e.g., Bart & Cone, 2012; Bart et al., 2017; O'Brien et al., 1999). A number of GZWs have been identified on the middle and outer shelf along the PITE, indicating retreat stillstands since the last glacial maximum (Graham et al., 2016; Jakobsson et al., 2012). The preservation of the Pliocene bGZWs on the ASE shelf indicates that their formations were followed by periods of hemipelagic deposition in a presumably glaciomarine environment without erosive glacial advances (Figure 3). The observed bGZW-2, -3 and -4 are 3–4 km in along-flow direction with crests 40–50 m high (based on a seismic interval velocity of 2,100 m/s derived from stacking velocities) and have the typical asymmetric wedge shape with a steep seaward flank downstream of the paleo-ice flow direction (Figure 2c). The lowermost imaged wedge (bGZW-1) shows an atypical, slightly steeper upstream flank. However, it is possible that this feature (a) is imaged obliquely by the corresponding seismic line, or (b) represents a paleo-shelf break bGZW, or (c) a lateral shear moraine (Batchelor & Dowdeswell, 2015; Dowdeswell & Fugelli, 2012).

Undisturbed sequences between the bGZWs can be used to estimate time periods without ice grounding events occurring in this area. Such an undisturbed sedimentary layer is observed between bGZW-2/-3 and bGZW-4 with a thickness of 50–70 m (Figure 2c), which is substantially thicker than the ~20-m vertical resolution limit of the seismic record. As drill core records from the outer ASE shelf do not exist to enable precise determination of age and duration of this sedimentary unit, we estimate an average sedimentation rate of 22 cm kyr⁻¹ from the total Pliocene (2.75-myrr timespan) layer thickness of 600 m south of the onset

of shelf progradation (Figure 2c). This is a minimum rate for the entire Pliocene, because sequences eroded by grounded ice cannot be accounted for. The Pliocene sedimentation rate is higher than at other Antarctic outer shelf drill sites, for example, 8.6 cm kyr⁻¹ for the early Pliocene (5.12–4.25 Ma) on the western Antarctic Peninsula (Bart & Iwai, 2012) and a minimum rate of 5.4 cm kyr⁻¹ for the mid to late Pliocene in the Ross Sea (McKay et al., 2019). Our estimated sedimentation rate for the outer ASE shelf suggests that deposition of the 50–70 m thick undisturbed layer between bGZW-2/-3 and bGZW-4 took about 227–318 kyr, an interval corresponding to 5–7 obliquity cycles in Earth's orbit. Ross Sea drill records indicate that during the Miocene and early Pliocene obliquity acted as the dominant orbital forcing of glacial/interglacial changes archived in sedimentary facies (Naish et al., 2009).

For comparing our seismic observations from the ASE outer shelf with IODP Expedition 379 drill records on the rise, we consider sedimentation rates and variations in the distribution of biogenic material in the drift sediments. We take into account that drift sedimentation is significantly controlled by ocean-bottom currents and not only by direct sediment supply from the nearby shelf and surface waters. Both drill sites (Figure 1) predominantly contain thinly laminated silty clay of variable biogenic content, with the occasional presence of dispersed sands and gravel. For this study, we consider only the Site U1532 record because of its higher linear sedimentation rate and higher temporal resolution.

Most Site U1532 cores contain lithological facies dominated by terrigenous detritus, suggesting deposition mainly during phases of colder climate, when plankton productivity was suppressed due to (nearly) permanent sea-ice cover (Hepp et al., 2006). Results show that terrigenous sedimentation dominated from the late Miocene to about 5.0 Ma in the early Pliocene. From 5.0 to 4.2 Ma, some sequences comprise a low to common diatom presence, potentially associated with more favorable climate conditions. With the exception of one interval, sedimentation rates were relatively high in the early Pliocene with a maximum of 61 cm kyr⁻¹ between 4.63 and 4.49 Ma. Some sediment intervals contain higher amounts of biosiliceous material, indicating a more pelagic and hemipelagic sedimentation, likely associated with phases of lower supply of glacial debris from the shelf. These sediments often contain pebbles indicating increased concentration of iceberg-rafted debris (IRD). The drill depth interval from 90 to 240 m spans the stratigraphic age range from 4.2 to 3.2 Ma and exhibits a relatively high abundance of diatoms (Figure 4). The interval from 4.2 to 3.2 Ma with the highest diatom abundance correlates with decreasing sedimentation rates to 18 cm kyr⁻¹, continuing to decrease throughout the late Pliocene and Pleistocene.

4. Discussion and Conclusions

Our integrated analysis of seismic stratigraphy, age constraints and general sedimentation characteristics from the Expedition 379 records links processes of major glacial progradation of the ASE shelf during the Pliocene to sediment deposition on a large drift on the continental rise about 280 km offshore from the shelf break. Bathymetric data show that the deep-sea channels starting at the slope base run seaward parallel to the elongated drift (Figure S2), suggesting that glacial detritus originating from the shelf provided most of the terrigenous drift sediments via gravity-driven transport running through the channels, with bottom-current capture and subsequent deposition of their suspended load on the drift. Although drift sediments can contain far-traveled components delivered by bottom currents (e.g., Hillenbrand & Ehrmann, 2002), initial shipboard analyses of the clay mineral provenance of the drift sediments showed that the assemblages at Sites U1532 and U1533 are characterized by the occurrence of the clay mineral kaolinite (Wellner et al., 2021a), indicating a predominant source on the ASE shelf (e.g., Ehrmann et al., 2011). While this rather rules out significant bottom-current delivery of far-traveled detritus, a clear separation of drift sediments into proximal components directly supplied from the nearby shelf and distal components supplied from the wider ASE shelf cannot be made at this stage.

The prograded wedge with lateral shelf growth of up to 80 km (Figures 2c, S3 and S4) formed in the early Pliocene (before ~4.2 Ma) and, to a smaller extent, in the late Pliocene (after ~3.2 Ma) and Pleistocene. We identified a minimum of eight early Pliocene, three late Pliocene and three Pleistocene ice-sheet advances associated with the seaward migration of the shelf-break (Figure 2c). The actual number of ice-sheet advances to the outer shelf is probably higher as not all may have reached the shelf edge (e.g., Klages et al., 2017), and some shelf-break formations were possibly eroded by a later grounded ice advance. Our




Figure 4. Lithology units, pebble abundance (visually counted number of clasts >4 mm per 10 cm core interval), relative diatom abundance, stratigraphic ages and sedimentation rates from shipboard analysis of drill cores from International Ocean Discovery Program (IODP) Expedition 379 Site U1532 (Gohl et al., 2021; Wellner et al., 2021a). Diatom abundance in smear slides is categorized as barren (B; 0%), trace (T; <2%), rare (R; 2%–5%), few (F; 5%–10%), common (C; 10%–20%), abundant (A; 20%–40%), and very abundant (V; 40%–60%). Gray shading of drill cores indicates core recovery. CSF-A is Core Depth Below Sea Floor-A.

observations are consistent with the hypothesis by Bart (2001) on major expansion of the Antarctic ice sheets in the early Pliocene. The high sedimentation rates on Resolution Drift during the early Pliocene documented by the Expedition 379 records are presumably related to major shelf progradation during this period, which resulted in high sediment supply to the rise.

Several bGZWs are observed in the Pliocene sequences of the outer shelf (Figure 2c). Such glaciomorphological features are only preserved when grounded ice does not override and erode them soon after their formation. The preservation of the bGZWs together with the thick sediment layer deposited during the time interval between formation of bGZW-2/-3 and bGZW-4 suggest deposition of this layer during a prolonged period, when the WAIS was receded and did not advance beyond the middle shelf. We infer from these data that no major WAIS advance occurred over many orbital cycles (~5–7 obliquity cycles) in the mid-Pliocene. This interval may have been dominated by interglacial deposition of meltwater plumes and IRD in a possibly temperate glacial setting, where subglacial bedforms can be quickly buried (e.g., Dowdeswell & Vasquez, 2013; Dowdeswell et al., 1998). Alternatively, biogenic-rich material, as recorded in the early to mid-Pliocene diatomites in the AND-1B core from the Ross Sea and in ODP Leg 178 cores of Sites 1095 and 1096 from the western Antarctic Peninsula, indicating warm climatic conditions and high sedimentation rates (Escutia et al., 2009; McKay et al., 2012), may have dominated sedimentation on the ASE shelf. Both scenarios are consistent with the higher abundances of diatoms and IRD in the mid-Pliocene section of Site U1532, especially from 4.2 to 3.2 Ma (Figure 4). This interval may have been an extended period of WAIS retreat with high sedimentation rates on the shelf, causing rapid burial of subglacial bedforms. Reduced downslope transport resulted in the accumulation of hemipelagic, microfossil and IRD-enriched deposits on Resolution Drift under sedimentation rates lower than before 4.2 Ma (Figure 4). Our hypothesis of a long-term WAIS retreat is based on the observation of the bGZW zone, but the determination of the exact time span of this interval can only be tested by future drilling into these shelf sequences.

Our study results for the Amundsen Sea are consistent with the global climate record of generally elevated global temperatures for the mid to late Pliocene (Lisiecki & Raymo, 2005; Westerhold et al., 2020) (Figure S5). The extended warm period inferred from our study falls into a period of Pliocene sea-level highstands from 4.39 to 3.27 Ma (e.g., Dumitru et al., 2019), but occurred earlier than the so-called mid-Pliocene or mid-Piacenzian Warm Period (e.g., Dowsett et al., 2016; Raymo et al., 2018). Our results also compare to an extended period of open-water deposition on the Ross Sea shelf (McKay et al., 2012; Naish et al., 2009). At least when considering processes on broad scales of several orbital cycles, the Amundsen and Ross Sea sectors of the WAIS appear to have behaved in phase. The enormous shelf growth in the ASE, interrupted by long periods of glacial retreat, indicates that this drainage sector of the WAIS was highly dynamic in the Pliocene, likely acting as precursor to phases of partial or complete WAIS collapse since the establishment of a continent-wide ice sheet.

Appendix A

IODP Expedition 379 Scientists: A. Klaus (IODP Texas A&M Univ., USA), D. Kulhanek (IODP Texas A&M Univ., USA), T. Bauersachs (Univ. Kiel, Germany), M. Courtillot (Univ. Perpignan, France), E. Cowan (Appalachian State Univ., USA), M. De Lira Mota (Unisinos Univ., Brazil), M. Esteves (Univ. Tromsø, Norway), J. Fegyveresi (Northern Arizona Univ., USA), L. Gao (China Univ. Geosciences, China), A. Halberstadt (Univ. Massachusetts-Amherst, USA), K. Horikawa (Toyama Univ., Japan), M. Iwai (Kochi Univ., Japan), J. Kim (Korea Inst. Geoscience and Minerals, Rep. Korea), T. King (Univ. South Florida, USA), M. Penkrot (Texas A&M Univ., USA), J. Prebble (GNS Science, New Zealand), W. Rahaman (National Center for Polar and Ocean Res., India), B. Reinardy (Univ. Stockholm, Sweden), J. Renaudie (Museum f. Naturkunde Berlin, Germany), D. Robinson (Univ. Houston, USA), R. Scherer (Northern Illinois Univ., USA), C. Siddoway (Colorado College, USA), L. Wu (Tonji Univ., China), M. Yamane (Nagoya Univ., Japan).

Data Availability Statement

Drill core data from IODP Expedition 379 are available from <http://web.iodp.tamu.edu/LORE/>. Seismic data files of profile TH86003B are available from <https://sdls.ogs.trieste.it>, and those of profiles AWI-20100130, -0131, -0132 and -0134 are available from <https://doi.pangaea.de/10.1594/PANGAEA.931266>, <https://doi.pangaea.de/10.1594/PANGAEA.931265>, <https://doi.pangaea.de/10.1594/PANGAEA.931268>, and <https://doi.pangaea.de/10.1594/PANGAEA.931263>.

Acknowledgments

The authors are grateful to the International Ocean Discovery Program (IODP) organizations, in particular the JOIDES Resolution Science Operator (JRSO) as well as the master and his ship and drill crews of RV JOIDES Resolution for their support in conducting IODP Expedition 379. This research used samples and data provided by the IODP. Thanks to John B. Anderson for providing SCS line NBP9902-11. The TH86 line is from the SCAR Antarctic Seismic Data Library System (SDLS). Our thanks also go to the master, crew and science team of the RV Akademik Alexander Karpinsky for collecting and providing swath bathymetry data of the deep-sea channel. The contribution of R. Lamb was funded by the Deutsche Forschungsgemeinschaft (DFG) in the framework of Priority Program 527 IODP through grant GO724/19-1. Contribution of C.-D. Hillenbrand was funded through Natural Environment Research Council (NERC) UK IODP grant NE/T010975/1. S. Passchier was funded by the U.S. Science Support Program under National Science Foundation (NSF) award #OCE1450528 subcontract 94 (GG009393). The study contributes to the Scientific Research Program “Instabilities and Thresholds in Antarctica” (INSTANT) of the Scientific Committee for Antarctic Research (SCAR). The authors would like to thank Emerson E&P Software, Emerson Automation Solutions, for providing licenses for the seismic software Paradigm in the scope of the Emerson Academic Program. Constructive comments by Laura De Santis and two anonymous reviewers helped improve the manuscript. Open access funding enabled and organized by Projekt DEAL.

References

- Anderson, J. B., & Bartek, L. R. (1992). Cenozoic glacial history of the Ross Sea revealed by intermediate resolution seismic reflection data combined with drill site information. In J. P. Kennett, & D. A. Warnke (Eds.), *The Antarctic Paleoenvironment: A perspective on Global Change, Part One Antarctic Research Series* (Vol. 56, pp. 231–263). Washington, DC: American Geophysical Union.
- Arndt, J. E., Schenke, H.-W., Jakobsson, M., Nitsche, F. O., Buys, G., Goleby, B., et al. (2013). The international bathymetric chart of the southern ocean (IBCSO) version 1.0: A new bathymetric compilation covering circum-Antarctic waters. *Geophysical Research Letters*, *40*, 3111–3117. <https://doi.org/10.1002/grl.50413>
- Barker, P. F., Camerlenghi, A., Acton, G. D., & Ramsay, A. T. S. (Eds.), (2002). Proceedings of the ocean drilling program, scientific results. College Station: Ocean Drilling Program.
- Bart, P. F., & Cone, A. N. (2012). Early stall of West Antarctic Ice Sheet advance on the eastern Ross Sea middle shelf followed by retreat at 27,500 ¹⁴C yr BP. *Palaeogeography, Palaeoclimatology, Palaeoecology*, *335–336*, 52–60. <https://doi.org/10.1016/j.palaeo.2011.08.007>
- Bart, P. J. (2001). Did the Antarctic ice sheets expand during the early Pliocene? *Geology*, *29*, 67–70. [https://doi.org/10.1130/0091-7613\(2001\)029<0067:dtaise>2.0.co;2](https://doi.org/10.1130/0091-7613(2001)029<0067:dtaise>2.0.co;2)
- Bart, P. J., Anderson, J. B., & Nitsche, F. (2017). Post-LGM grounding-line positions of the Bindschadler Paleo ice stream in the Ross Sea Embayment, Antarctica. *Journal of Geophysical Research: Earth Surface*, *122*, 1827–1844. <https://doi.org/10.1002/2017JF004259>
- Bart, P. J., & De Santis, L. (2012). Glacial intensification during the Neogene: A review of seismic stratigraphic evidence from the Ross Sea, Antarctica, continental shelf. *Oceanography*, *25*, 166–183. <https://doi.org/10.5670/oceanog.2012.92>
- Bart, P. J., Hillenbrand, C. D., Ehrmann, W., Iwai, M., Winter, D., & Warny, S. A. (2007). Are Antarctic Peninsula Ice Sheet grounding events manifest in sedimentary cycles on the adjacent continental rise? *Marine Geology*, *236*, 1–13. <https://doi.org/10.1016/j.margeo.2006.09.008>
- Bart, P. J., & Iwai, M. (2012). The overdeepening hypothesis: How erosional modification of the marine-scape during the early Pliocene altered glacial dynamics on the Antarctic Peninsula's Pacific margin. *Palaeogeography, Palaeoclimatology, Palaeoecology*, *335–336*, 42–51. <https://doi.org/10.1016/j.palaeo.2011.06.010>
- Batchelor, C. L., & Dowdeswell, J. A. (2015). Ice-sheet grounding zone wedges (GZWs) on high-latitude continental margins. *Marine Geology*, *363*, 65–92. <https://doi.org/10.1016/j.margeo.2015.02.001>
- Boer, de, B., Dolan, A. M., Bernaldes, J., Gasson, E., Goelzer, H., Gollidge, N. R., et al. (2015). Simulating the Antarctic ice sheet in the late-Pliocene warm period: PLISMIP-ANT, an ice-sheet model intercomparison project. *The Cryosphere*, *9*, 881–903. <https://doi.org/10.5194/tc-9-881-2015>
- DeConto, R. M., & Pollard, D. (2016). Contribution of Antarctica to past and future sea-level rise. *Nature*, *531*, 591–597. <https://doi.org/10.1038/nature17145>
- De Santis, L., Prato, S., Brancolini, G., Lovo, M., & Torelli, L. (1999). The Eastern Ross Sea continental shelf during the Cenozoic: Implications for the West Antarctic ice sheet development. *Global and Planetary Change*, *23*, 173–196. [https://doi.org/10.1016/S0921-8181\(99\)00056-9](https://doi.org/10.1016/S0921-8181(99)00056-9)
- Dowdeswell, J. A., Cofaigh, C. Ó., & Anderson, J. B. (2006). Morphology and sedimentary processes on the continental slope off Pine Island Bay, Amundsen Sea, West Antarctica. *Geological Society of America Bulletin*, *118*, 606–619. <https://doi.org/10.1130/b25791.1>
- Dowdeswell, J. A., Elverhoi, A., & Spielhagen, R. (1998). Glacimarine sedimentary processes and facies on the Polar North Atlantic margins. *Quaternary Science Reviews*, *17*, 243–272. [https://doi.org/10.1016/S0277-3791\(97\)00071-1](https://doi.org/10.1016/S0277-3791(97)00071-1)
- Dowdeswell, J. A., & Fugelli, E. M. G. (2012). The seismic architecture and geometry of grounding-zone wedges formed at the marine margins of past ice sheets. *Bulletin of the Geological Society of America*, *124*, 1750–1761. <https://doi.org/10.1130/B30628.1>
- Dowdeswell, J. A., & Vasquez, M. (2013). Submarine landforms in the fjords of southern Chile: Implications for glacimarine processes and sedimentation in a mild glacier-influenced environment. *Quaternary Science Reviews*, *64*, 1–19. <https://doi.org/10.1016/j.quascirev.2012.12.003>
- Dowsett, H., Dolan, A., Rowley, D., Moucha, R., Forte, A. M., Mitrovica, J. X., et al. (2016). The PRISM4 (mid-Piacenzian) paleoenvironmental reconstruction. *Climate of the Past*, *12*, 1519–1538. <https://doi.org/10.5194/cp-12-1519-2016>
- Dumitru, O. A., Austermann, J., Polyak, V. J., Fornós, J. J., Asmerom, Y., Ginés, J., et al. (2019). Constraints on global mean sea level during Pliocene warmth. *Nature*, *574*, 233–236. <https://doi.org/10.1038/s41586-019-1543-2>
- Dutton, A., Carlson, A. E., Long, A. J., Milne, G. A., Clark, P. U., DeConto, R., et al. (2015). Sea-level rise due to polar ice-sheet mass loss during past warm times. *Science*, *349*. <https://doi.org/10.1126/science.aaa4019>
- Ehrmann, W. U., Hillenbrand, C.-D., Smith, J. A., Graham, A. G. C., Kuhn, G., & Larter, R. D. (2011). Provenance changes between recent and glacial-time sediments in the Amundsen Sea embayment, West Antarctica: Clay mineral assemblage evidence. *Antarctic Science*, *23*, 471–486. <https://doi.org/10.1017/S0954102011000320>
- Escutia, C., Bárcena, M. A., Lucchi, R. G., Ballegeer, A. M., Gonzales, J. J., & Harwood, D. M. (2009). Circum-Antarctic warming events between 4 and 3.5 Ma recorded in marine sediments from the Prydz Bay (ODP Leg 188) and the Antarctic Peninsula (ODP Leg 178) margins. *Global and Planetary Change*, *69*, 170–184. <https://doi.org/10.1016/j.gloplacha.2009.09.003>
- Gasson, E., DeConto, R. M., & Pollard, D. (2016). Modeling the oxygen isotope composition of the Antarctic ice sheet and its significance to Pliocene sea level. *Geology*, *44*(10), 827–830. <https://doi.org/10.1130/G38104.1>
- Gohl, K., Freudenthal, T., Hillenbrand, C.-D., Klages, J., Larter, R., Bickert, T., et al., (2017). MeBo70 seabed drilling on a polar continental shelf: Operational report and lessons from drilling in the Amundsen Sea Embayment of West Antarctica. *Geochemistry, Geophysics, Geosystems*, *18*, 4235–4250. <https://doi.org/10.1002/2017GC007081>
- Gohl, K., Unzelmann-Neben, G., Larter, R. D., Hillenbrand, C.-D., Hochmuth, K., Kalberg, T., et al. (2013). Seismic stratigraphic record of the Amundsen Sea Embayment shelf from pre-glacial to recent times: Evidence for a dynamic West Antarctic ice sheet. *Marine Geology*, *344*, 115–131. <https://doi.org/10.1016/j.margeo.2013.06.011>
- Gohl, K., Wellner, J. S., Klaus, A., & Expedition 379 Scientists. (2021). *Amundsen Sea West Antarctic Ice Sheet History, Proceedings of the International Ocean Discovery Program* (Vol. 379). College Station, TX: International Ocean Discovery Program. <https://doi.org/10.14379/iodp.proc.379.2021>
- Gollidge, N. R., Thomas, Z. A., Levy, R. H., Gasson, E. G. W., Naish, T. R., McKay, R. M., et al. (2017). Antarctic climate and ice-sheet configuration during the early Pliocene interglacial at 4.23 Ma. *Climate of the Past*, *13*, 959–975. <https://doi.org/10.5194/cp-13-959-2017>
- Graham, A. G. C., Jakobsson, M., Nitsche, F. O., Larter, R. D., Anderson, J. B., Hillenbrand, C.-D., et al. (2016). Submarine glacial-landform distribution across the West Antarctic margin, from grounding line to slope: The Pine Island-Thwaites ice-stream system. In J. A. Dowdeswell, M. Canals, M. Jakobsson, B. J. Todd, E. K. Dowdeswell, & K. A. Hogan (Eds.), *Atlas of submarine glacial landforms: Modern, quaternary and ancient* (pp. 493–500). Geological Society London, Memoirs. <https://doi.org/10.1144/M46.173>

- Graham, A. G. C., Larter, R. D., Gohl, K., Dowdeswell, J. A., Hillenbrand, C.-D., Smith, J. A., et al. (2010). Flow and retreat of the Late Quaternary Pine Island-Thwaites palaeo-ice stream, West Antarctica. *Journal of Geophysical Research*, *115*, F03025. <https://doi.org/10.1029/2009JF001482>
- Hepp, D., Mörz, T., & Grütznér, J. (2006). Pliocene glacial cyclicity in a deep-sea sediment drift (Antarctic Peninsula Pacific Margin). *Palaeogeography, Palaeoclimatology, Palaeoecology*, *231*, 181–198. <https://doi.org/10.1016/j.palaeo.2005.07.030>
- Hillenbrand, C.-D., & Ehrmann, W. (2002). Distribution of clay minerals in drift sediments on the continental rise west of the Antarctic Peninsula, ODP Leg 178, Sites 1095 and 1096. In P. F. Barker, A. Camerlenghi, G. D. Acton, & A. T. S. Ramsay (Eds.), *Proceedings of the ocean drilling program, scientific results 178* (pp. 1–29). College Station, TX: Ocean Drilling Program, Texas A&M University.
- Hochmuth, K., & Gohl, K. (2013). Glaciomarine sedimentation dynamics of the Abbot glacial trough of the Amundsen Sea embayment shelf, West Antarctica. In M. J. Hambrey, P. F. Barker, P. J. Barrett, V. Bowman, B. Davies, J. L. Smellie, & M. Tranter (Eds.), *Antarctic palaeoenvironments and earth-surface processes* (Vol. 381, pp. 233–244). Geological Society, London: Special Publications. <https://doi.org/10.1144/SP381.21>
- Hochmuth, K., & Gohl, K. (2019). Seaward growth of Antarctic continental shelves since establishment of a continent-wide ice sheet: Patterns and mechanisms. *Palaeogeography, Palaeoclimatology, Palaeoecology*, *520*, 44–54. <https://doi.org/10.1016/j.palaeo.2019.01.025>
- Jakobsson, M., Anderson, J. B., Nitsche, F. O., Gyllencreutz, R., Kirshner, A. E., Kirchner, N., et al. (2012). Ice sheet retreat dynamics inferred from glacial morphology of the central Pine Island Bay Trough, West Antarctica. *Quaternary Science Reviews*, *38*, 1–10. <https://doi.org/10.1016/j.quascirev.2011.12.017>
- Klages, J. P., Kuhn, G., Graham, A. G. C., Hillenbrand, C.-D., Smith, J. A., Nitsche, F. O., et al. (2015). Palaeo-ice stream pathways and retreat style in the easternmost Amundsen Sea Embayment, West Antarctica, revealed by combined multibeam bathymetric and seismic data. *Geomorphology*, *245*, 207–222. <https://doi.org/10.1016/j.geomorph.2015.05.020>
- Klages, J. P., Kuhn, G., Hillenbrand, C.-D., Smith, J. A., Graham, A. G. C., Nitsche, F. O., et al. (2017). Limited grounding-line advance onto the West Antarctic continental shelf in the easternmost Amundsen Sea Embayment during the last glacial period. *PLoS One*, *12*. <https://doi.org/10.1371/journal.pone.0181593>
- Klages, J. P., Salzmann, U., Bickert, T., Hillenbrand, C.-D., Gohl, K., Kuhn, G., et al. (2020). Temperate rainforests near the South Pole during peak Cretaceous warmth. *Nature*, *580*. <https://doi.org/10.1038/s41586-020-2148-5>
- Lisiecki, L. E., & Raymo, M. E. (2005). A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records. *Paleoceanography*, *20*. <https://doi.org/10.1029/2004PA001071>
- Lowe, A. J., & Anderson, J. B. (2002). Reconstruction of the West Antarctic ice sheet in Pine Island Bay during the Last Glacial Maximum and its subsequent retreat history. *Quaternary Science Reviews*, *21*, 1879–1897. [https://doi.org/10.1016/s0277-3791\(02\)00006-9](https://doi.org/10.1016/s0277-3791(02)00006-9)
- McKay, R. M., De Santis, L., Kulhanek, D. K., Ash, J. L., Beny, F., Browne, I. M., et al. (2019). Site U1522. In R. M. McKay, L. De Santis, D. K. Kulhanek, & the Expedition 374 Scientists (Eds.), *Ross Sea West Antarctic ice sheet history, Proceedings of the international ocean discovery program* (Vol. 374). College Station, TX: International Ocean Discovery Program. <https://doi.org/10.14379/iodp.proc.374.104.2019>
- McKay, R. M., Naish, T., Carter, L., Riesselman, C., Dunbar, R., Sjunneskog, C., et al. (2012). Antarctic and Southern Ocean influences on Late Pliocene global cooling. *Proceedings of the National Academy of Sciences of the USA*, *109*, 6423–6428. <https://doi.org/10.1073/pnas.1112248109>
- Naish, T., Powell, R., Levy, R., Wilson, G., Scherer, R., Talarico, F., et al. (2009). Obliquity-paced Pliocene West Antarctic ice sheet oscillations. *Nature*, *458*, 322–328. <https://doi.org/10.1038/nature07867>
- Nitsche, F. O., Cunningham, A. P., Larter, R. D., & Gohl, K. (2000). Geometry and development of glacial continental margin depositional systems in the Bellingshausen Sea. *Marine Geology*, *162*, 277–302. [https://doi.org/10.1016/s0025-3227\(99\)00074-2](https://doi.org/10.1016/s0025-3227(99)00074-2)
- Nitsche, F. O., Gohl, K., Vanneste, K., & Miller, H. (1997). Seismic expression of glacially deposited sequences in the Bellingshausen and Amundsen Seas, West Antarctica. In P. F. Barker, & A. K. Cooper (Eds.), *Geology and seismic stratigraphy of the Antarctic margin 2 Antarctic research series* (pp. 95–108). Washington, DC: American Geophysical Union.
- O'Brien, P. E., De Santis, L., Harris, P. T., Domack, E., & Quilty, P. G. (1999). Ice shelf grounding zone features of western Prydz Bay, Antarctica: Sedimentary processes from seismic and sidescan images. *Antarctic Science*, *11*, 78–91.
- Pagani, M., Liu, Z., LaRiviere, J., & Ravelo, A. C. (2010). High Earth-system climate sensitivity determined from Pliocene carbon dioxide concentrations. *Nature Geoscience*, *3*(1), 27–30. <https://doi.org/10.1038/ngeo724>
- Pollard, D., & DeConto, R. (2020). Continuous simulations over the last 40 million years with a coupled Antarctic ice sheet-sediment model. *Palaeogeography, Palaeoclimatology, Palaeoecology*, *537*, 109374. <https://doi.org/10.1016/j.palaeo.2019.109374>
- Raymo, M. E., Kozdon, R., Evans, D., Lisiecki, L., & Ford, H. L. (2018). The accuracy of mid-Pliocene $\delta^{18}\text{O}$ -based ice volume and sea level reconstructions. *Earth-Science Reviews*, *177*, 291–302. <https://doi.org/10.1016/j.earscirev.2017.11.022>
- Rebesco, M., Pudsey, C. J., Canals, M., Camerlenghi, A., Barker, P. F., Estrada, F., & Giorgetti, A. (2002). Sediment drifts and deep-sea channel systems, Antarctic Peninsula Pacific margin. In D. A. V. Stow, C. J. Pudsey, J. A. Howe, J.-C. Faugeres, & A. R. Viana (Eds.), *Deep-water contourite systems: Modern drifts and ancient series, seismic and sedimentary characteristics* (pp. 353–372). Geological Society London. <https://doi.org/10.1144/gsl.mem.2002.022.01.25>
- Rignot, E., Mouginot, J., & Scheuchl, B. (2011). Ice flow of the Antarctic Ice Sheet. *Science*, *333*. <https://doi.org/10.1126/science.1208336>
- Rignot, E., Mouginot, J., Scheuchl, B., Broeke, van den, M., Wessm, van, M. J., & Morlighem, M. (2019). Four decades of Antarctic Ice Sheet mass balance from 1979–2017. *Proceedings of the National Academy of Sciences of the USA*, *116*, 1095–1103. <https://doi.org/10.1073/pnas.1812883116>
- Scambos, T. A., Bell, R. E., Alley, R. B., Anandkrishnan, S., Bromwich, D. H., Brunt, K., et al. (2017). How much, how fast?: A science review and outlook for research on the instability of Antarctica's Thwaites Glacier in the 21st century. *Global and Planetary Change*, *153*, 16–34. <https://doi.org/10.1016/j.gloplacha.2017.04.008>
- Scheuer, C., Gohl, K., & Eagles, G. (2006). Gridded isopach maps from the South Pacific and their use in interpreting the sedimentation history of the West Antarctic continental margin. *Geochemistry, Geophysics, Geosystems*, *7*. <https://doi.org/10.1029/2006GC001315>
- Smith, B., Fricker, H. A., Gardner, A. S., Medley, B., Nilsson, J., Paolo, F. S., et al. (2020). Pervasive ice sheet mass loss reflects competing ocean and atmosphere processes. *Science*, *368*, 1239–1242. <https://doi.org/10.1126/science.aaz5845>
- Tucholke, B. E., Hollister, C. D., Weaver, F. M., & Vennum, W. R. (1976). Continental rise and abyssal plain sedimentation in the southeast Pacific Basin - Leg 35 Deep Sea Drilling Project. In C. D. Hollister, & C. Craddock (Eds.), *Deep Sea Drilling Project Initial Reports* (Vol. 35, pp. 359–400). Washington, DC: Deep Sea Drilling Project. <https://doi.org/10.2973/dsdp.proc.35.119.1976>
- Uenzelmann-Neben, G., & Gohl, K. (2012). Amundsen Sea sediment drifts: Archives of modifications in oceanographic and climatic conditions. *Marine Geology*, *299–302*, 51–62. <https://doi.org/10.1016/j.margeo.2011.12.007>

- Uenzelmann-Neben, G., & Gohl, K. (2014). Early glaciation already during the Early Miocene in the Amundsen Sea, Southern Pacific: Indications from the distribution of sedimentary sequences. *Global and Planetary Change*, *120*, 92–104. <https://doi.org/10.1016/j.gloplacha.2014.06.004>
- Wellner, J. S., Gohl, K., Klaus, A., Bauersachs, T., Bohaty, S. M., Courtillot, M., et al. (2021a). Site U1532. In K. Gohl, J. S. Wellner, & A. Klaus (Eds.), *Amundsen Sea West Antarctic Ice Sheet History, Proceedings of the International Ocean Discovery Program* (Vol. 379). College Station, TX: International Ocean Discovery Program. <https://doi.org/10.14379/iodp.proc.379.103.2021>
- Wellner, J. S., Gohl, K., Klaus, A., Bauersachs, T., Bohaty, S. M., Courtillot, M., et al. (2021b). Site U1533. In K. Gohl, J. S. Wellner, & A. Klaus (Eds.), *Amundsen Sea West Antarctic Ice Sheet History, Proceedings of the International Ocean Discovery Program* (Vol. 379), College Station, TX: International Ocean Discovery Program. <https://doi.org/10.14379/iodp.proc.379.104.2021>
- Wellner, J. S., Lowe, A. L., Shipp, S. S., & Anderson, J. B. (2001). Distribution of glacial geomorphic features on the Antarctic continental shelf and correlation with substrate: Implications for ice behavior. *Journal of Glaciology*, *47*, 397–411. <https://doi.org/10.3189/172756501781832043>
- Westerhold, T., Marwan, N., Drury, A. J., Liebrand, D., Agnini, C., Anagnostou, E., et al. (2020). An astronomically dated record of Earth's climate and its predictability over the last 66 million years. *Science*, *369*, 1383–1387. <https://doi.org/10.1126/science.aba6853>
- Yamaguchi, K., Tamura, Y., Mizukoshi, I., & Tsuru, T. (1988). Preliminary report of geophysical and geological surveys in the Amundsen Sea, West Antarctica. *Proceedings of the NIPR Symposium on Antarctic Geosciences*, *2*, 55–67.