


Analysis of Determinants of Stunting and Identifications of Stunting Risk Profiles Among Under 2-Year-Old Children in Ethiopia. A Latent Class Analysis

Anteneh Fikrie¹ , Berhanu Adula¹, Jitu Beka¹, Dejene Hailu², Cheru Atsmegiorgis Kitabo³, and Mark Spigt^{4,5}

Abstract

Background: Childhood stunting has a long-term impact on cognitive development and overall well-being. Understanding varying stunting profiles is crucial for targeted interventions and effective policy-making. Therefore, our study aimed to identify the determinants and stunting risk profiles among 2-year-old children in Ethiopia.

Methods and materials: A cross-sectional study was conducted on 395 mother–child pairs attending selected public health centers for growth monitoring and promotion under 5 outpatient departments and immunization services. The data were collected by face-to-face interviews, with the anthropometric data collected using the procedure stipulated by the World Health Organization. The data were entered using Epi Data version 4.6 and exported to STATA 16 and Jamovi version 2.3.28 for analysis. Bayesian logistic regression analysis was conducted to identify potential factors of stunting. Likewise, lifecycle assessment analysis (LCA) was used to examine the heterogeneity of the magnitude of stunting.

Results: The overall prevalence of stunting in children under 24 months was 47.34% (95% confidence interval (CI): 42.44–52.29%). The LCA identified 3 distinct risk profiles. The first profile is **Class 1**, which is labeled as low-risk, comprised 23.8% of the children, and had the lowest prevalence of stunting (23.4%). This group characterized as having a lower risk to stunting. The second profile is **Class 2**, which is identified as high-risk, comprised 47.1%, and had a high prevalence of stunting (66.7%), indicating a higher susceptibility to stunting compared to Class 1. The third profile is **Class 3**, which is categorized as mixed-risk and had a moderate stunting prevalence of 35.7%, indicating a complex interplay of factors contributing to stunting.

Conclusion: Our study identified 3 distinct risk profiles for stunting in young children. A substantial amount (almost half) is in the high-risk category, where stunting is far more common. The identification of stunting profiles necessitates considering heterogeneity in risk factors in interventions. Healthcare practitioners should screen, provide nutrition counseling, and promote breastfeeding. Policymakers should strengthen social safety nets and support primary education.

Keywords

child stunting, Bayesian logistic regression, latent class analysis, Southern Ethiopia

¹ School of Public Health, Institute of Health, Bule Hora University, Bule Hora, Ethiopia

² School of Public Health, College of Medicine and Health Sciences, Hawassa University, Hawassa, Ethiopia

³ Department of Statistics, College of Natural and Computational Sciences, Hawassa University, Hawassa, Ethiopia

⁴ CAPHRI, School for Public Health and Primary Care, Maastricht University, Maastricht, the Netherlands

⁵ General Practice Research Unit, Department of Community Medicine, UiT the Arctic University of Norway, Tromsø, Norway

Submitted May 7, 2024. Accepted June 22, 2024.

Corresponding Author:

Anteneh Fikrie, School of Public Health, Institute of Health, Bule Hora University, P.O. Box 144, Bule Hora, Ethiopia.
Email: antenehfikrie3@gmail.com

Data Availability Statement included at the end of the article



Introduction

Stunting, defined as low height-for-age, is a significant public health concern in children under 2 years old.¹ It is an indicator of developmental delays, poor child health, and chronic malnutrition, affecting children's recovery, vulnerability to long-term illnesses, missed learning opportunities, and school performance.^{2,3} It is estimated to be a significant contributor to around 45% of children's deaths.⁴ Despite the global prevalence decreasing from 32.4% in 2000 to 22.3% in 2022,⁵ this reduction is insufficient to meet the World Health Organization (WHO)'s Global Nutrition Targets 2025.⁴ In 2022, Africa had the highest number of stunted children, accounting for 43%.⁶ Ethiopia's stunting rate is pervasive, with an overall rate of 35.1% among children under 2.⁷ This highlights the urgency for effective strategies to address childhood stunting.

The Ethiopian government began implementing growth monitoring and promotion (GMP) in the 1990s as part of its attempts to tackle child malnutrition.⁸⁻¹⁰ GMP is a nutritional intervention that measures, plots, and interprets child growth regularly to counsel and take action when abnormalities are detected, empowering mothers to improve their children's nutritional status.¹¹ However, despite this intervention, the prevalence of stunting remains unacceptably high in the country.¹²⁻¹⁴ Several studies have identified inadequate dietary intake, maternal nutrition, breastfeeding practices, and frequent infections as significant contributors to stunting in young children¹⁵⁻¹⁸ and emphasizing the need to consider heterogeneity and the complex interplay between these factors.¹⁹⁻²¹

Despite existing research providing valuable insights into the overall prevalence and risk factors of stunting,¹²⁻¹⁴ there is a need for more nuanced analyses that go beyond broad categorizations to uncover hidden patterns and variations within the stunted population. Studies in India have yielded diverse findings on subgroup analysis.^{22,23} For instance, a study in Gondar City, Northern Ethiopia identified a strong association between stunting and occupational status of a mother,¹⁵ while another in Rwanda revealed a more significant influence of water and sanitation.²⁴ Similarly, a study finding from South Africa revealed socio-demographic and infant and young child feeding (IYCF) factors associated with stunting.²⁰ These contrasting results highlight the need for further research that delves deeper into subgroup variations across Ethiopia to address the knowledge gap. Moreover, subgroup analyses may offer new insights for identifying and targeting specific populations, providing valuable insights into underlying mechanisms and potential interventions.²⁵ Therefore, our study aimed to identify the determinants of stunting and identifications of stunting profiles among two-year-old children in Ethiopia. The findings would provide insights into childhood stunting and offer targeted interventions, contributing to the development of more effective approaches to combat stunting.

Methods and Materials

Study Design

We used a facility-based cross-sectional study to investigate potential subgroups of stunted children under 2 years old.

Study Area and Period

The study was conducted in the West Guji Zone, Southern Oromia Regional State. Its administrative center is Bule Hora town, which is located at a distance of 467 km from Addis Ababa, the capital city of Ethiopia, on the paved Addis Ababa to Moyale highway. The zone has 9 woredas and 2 town administrations. It has an estimated total population of 1,389,821, of which 681,012 are males, and 708,809 are females. The zone has immense potential due to its coffee cultivation, mineral wealth, and cultural richness and largely focuses on pastoralism or subsistence farming. Its situation contrasts significantly with some of the region's more stable and developed zones. There is 1 teaching hospital, 3 primary hospitals, 42 health centers, and 166 health posts. The total number of pregnant women, as estimated from the total population of West Guji Zone in 2021, was 48,234 (West Guji Zone Health Department HMIS Report, 2022). The study was conducted from May 1 to 30, 2023.

Population, Sample Size Determination, Sampling Technique, and Procedure

Children aged below 24 months, accompanied by their mothers, who were attending 3 randomly selected public health centers for GMP, in 5 outpatient departments and immunization services in West Guji Zone, Southern Ethiopia, were included in our study. To ensure the accuracy and reliability of our data, we excluded mothers who were only visiting the health center while not living permanently in the area. The sample size was determined using the single population proportion formula. Considering the proportion of children short for their age or stunted 37.1%,¹³ 95% confidence interval, 5% degree of precision, and 10% non-response rate were also considered. The final calculated sample size was 395. To identify factors associated with stunting, the adequacy of the sample size was examined using Epi-info version 7.1. We employed a simple random sampling technique (using the lottery method) to select 3 health centers (Bule Hora Health Center, Finchawa Health Center, and Guangua Health Centre) based on the availability of functional growth monitoring and promotion (GMP), immunization service, and security issues. Thus, to account for variations in the number of clients visiting each health center daily, we estimated the daily attendance based on the previous month's data. The total sample size needed for the study was then divided proportionally among the 3 centers, ensuring that each center had adequate representation. Within each health center, participants were selected using systematic random sampling. This involved randomly selecting a starting point from the daily client list, and then choosing every 2nd children thereafter until the required sample size for that center was attained. This sampling method helped ensure that the participants were selected randomly.

Study Variables and Operational Definitions

Stunting was the dependent variable of the study. In contrast, sociodemographic and economic factors, maternal obstetric

and reproductive health-related factors, child health and dietary and feeding practices, water, sanitation and hygiene, and maternal food and nutrition knowledge were the independent variables of the study.

Stunting: A child with a height-for-age Z-score (HAZ) less than minus 2 standard deviations (<-2 SD).²⁶ Nutrition and food knowledge refer to an individual's understanding of nutrition, including the intellectual ability to remember and recall food- and nutrition-related terminology, specific pieces of information, and facts.²⁷ A total of 24 items were used to assess the nutrition and food knowledge of mothers, and each correct answer was scored as "1," while each "I don't know" answer was scored as "0." The maximum score that was obtained by correctly answering all the knowledge questions was 24, while the minimum was 0. Thus, women who scored $\geq 70\%$ (out of 24) on the knowledge questions were regarded as having good knowledge, whereas women who scored $<70\%$ (out of 24) on the knowledge questions were regarded as having poor knowledge.²⁸ Food Insecurity Access Scale (HFAS) was used to estimate the prevalence of food insecurity and assess the various ways in which households may experience food insecurity by collecting data on access to food, frequency of food insecurity, and the severity of its effects.²⁹ Therefore, when the average HFAS score is 0 or 1, households are regarded as food secured. When the average HFAS score is greater than or equal to 2, households are regarded as food insecure.²⁹

Dietary diversity score: This is the number of food groups consumed by a child aged 6 to 23 months in the last 24 h out of 7 food groups: (a) grains, roots, and tubers; (b) legumes and nuts; (c) dairy products; (d) flesh foods; (e) eggs; (f) vitamin-A-rich fruits and vegetables; and (g) other fruits and vegetables.³⁰

Minimum dietary diversity¹³: A child aged 6 to 23 months having the minimum diet diversity in the last 24 h consumed at least 4 food groups out of the 7 food groups. Thus, a child with a DDS of less than 4 was classified as having poor dietary diversity, whereas a DDS of greater than 4 was good dietary diversity.¹³

Data Collection Instruments, Methods, and Quality Assurance

The data collection tool was constructed by adapting previous peer-reviewed and published articles^{13,28,31,32} and the United Nations Children's Fund (UNICEF) 2020 framework.³³ Three BSc degree nurses collected the data using a face-to-face interview and were supervised by 2 Master of General Public Health holder health workers. The anthropometric data (height/length) was measured as per the procedure stipulated by the WHO/UNICEF for taking anthropometric using³⁴ a horizontal wooden length measuring board, without shoes, with the child in a recumbent position on a hard surface and recorded to the nearest 0.1 cm.³⁵ Data quality was ensured through proper training of data collectors and supervisors, covering

study objectives, participant approach, instrument approach, data handling, and time management. The questionnaire was pretested before the actual data collection on 5% of the sample size in Gerba. The collected data were checked for consistency, completeness, and relevance daily during data collection by the supervisors and principal investigator.

Data Processing and Analysis

Data were entered into Epi Data version 4.6, and then exported to STATA 16 for analysis. Descriptive statistics, including percentages, frequencies, mean with standard deviation, and median with interquartile range, were run to describe the data. To identify the associations between independent variables and stunting prevalence in children, we used a Bayesian logistic regression. Random-walk Metropolis-Hastings (RWMH) sampling algorithm was used to obtain samples from the posterior distribution. MCMC iterations = 12,500, burn-in = 2,500, 10,000 samples from the posterior distribution at 500 burn-in terms discarded. We computed posterior medians, odds ratios, and 95% credible intervals with equal tails to appraise the statistical significance of each independent variable's influence on stunting. Furthermore, the accuracy of posterior estimates was assessed based on the Markov chain (MC) error. Nutritional indices (HFA) were determined using the Emergency Nutrition Assessment (ENA) for Standardizing Monitoring and Assessment of Relief and Transition (SMART) 2011 software, and nutritional status was determined concerning age- and sex-specific growth based on the WHO Child Growth Standards.²⁶ Diagnostic tools, such as a trace plot, density plot, autocorrelation plot, and histogram, were used to assess the convergence of MCMC (Supplementary file 1). The algorithm's convergence is evident in the plots of significant predictors. We used latent class analysis (LCA) to identify unobservable subgroups within a population based on observed variables and run using Jamovi software version 2.3.28.0 for LCA analysis using the SnowRMM module.³⁶ Key indicators included maternal food and nutrition knowledge, birth intervals, hand washing, maternal education, ANC follow-up, family planning utilization, unsafe drinking water, breast milk, and food insecurity. Meanwhile, sex, age, dietary diversity, early breastfeeding initiation, maternal occupation, hand washing before child feeding, and sanitation facility were taken from previous studies.²³ The optimal number of latent classes was selected using statistical fit indices like the Akaike information criterion, Bayesian information criterion, consistent Akaike information criterion, log-likelihood, entropy, and Elbow plot.^{37,38} AIC and CAIC penalize complexity models, with lower values indicating better fit, while BIC penalizes models with more parameters for parsimony. Entropy measures group assignments' certainty, with higher values indicating clear risk separation and lower values indicating potential overlap and uncertainty, with values above 0.80 considered good classification markers.³⁹ The elbow plot shows the relationship between latent class numbers and fit statistics, with plateauing

points indicating diminishing returns and overfitting indicating model complexity.

In our research, Figure 1 displays that in the elbow plot, the CAIC and BIC exhibit a discernible inflection point at 3 classes, signifying that the inclusion of additional classes does not notably enhance the model's goodness of fit. Similarly, Figure 1 indicates that the initial decline in BIC as the number of classes' increases signifies a substantial improvement in fit, while a marginal upsurge from 3 to 5 implies that a 3-class model is the most easily comprehensible solution. Consequently, as indicated in Table 1, the 3-class model was selected for further examination due to its optimal balance of fit, complexity, and interpretability.

Results

Sociodemographic and Economic Characteristics of the Respondents

Table 2 shows that mothers' median (IQR) age was 30 (27-34) years, with the highest number (64.6%) falling from the 26 to 34 age range. In contrast, the children's median (IQR) age was 12 months, ranging from 9 to 18 months, with the highest percentage of 29.6% falling in the 18 and 23-month age range. In terms of the child's gender, 54.9% of them were girls. The majority of mothers, 61.5%, reported moderate food insecurity, while only 15.7% reported having food security.

Maternal Obstetrics and Reproductive Health and Infant and Young Child Feeding Practices

Table 3 shows that a majority (53.7%) of mothers reported a birth interval of less than 24 months between consecutive

births. A total of 349 (88.4%) and 249 (63%) mothers received ANC follow-up and utilized family planning, respectively. Breastfeeding initiation within the first hour of birth remained relatively low (27.6%). Likewise, 61% of children have inadequate dietary intake. Over half (51.4%) of mothers reported a lack of adequate food and nutrition knowledge.

Environmental Health-Related Characteristics of the Participants

Table 4 shows a significant proportion of (21.3%) of the study participants relying on unimproved water sources, including rivers, ponds, wells, and unprotected springs. This aligns with the result that a substantial majority (53.4%) did not employ water treatment methods before consumption. Moreover, hand washing practices were also inadequate, with less than half of 48.1% adhering to proper procedures at critical times.

Bayesian logistic regression analysis of factors linked to stunting in children aged 24 months

As Table 5 shows, a Bayesian binary logistic regression model found that mothers with good knowledge of food and nutrition significantly reduced the odds of having a stunted child (OR = 0.023, 95% CI: 0.004, 0.061). Longer birth intervals (OR = 0.24, 95% CI: 0.105, 0.480) and having primary education (OR = 0.26, 95% CI: 0.099, 0.560) also reduced stunting odds. Additionally, practicing hand washing after toilet use reduced stunting by 88% in children (OR = 0.119, 95% CI: 0.033, 0.299).

Conversely, children of mothers without ANC history (OR = 8.23, 95% CI: 2.72, 20.18), living in households without safe drinking water (OR = 109.44, 95% CI: 232.67, 303.01),




Figure 1. Elbow plot examining model fit statistics to determine the optimal number of latent classes. Abbreviations: AIC, Akaike information criterion; BIC, Bayesian information criterion.

Table 1. Model Comparison for Selecting Optimal Latent Classes.

Class	AIC	BIC	ARABIC	CAIC	Log-likelihood	χ^2	G ²
1	5614	5662	5624	5674	-2795	8009	1546
2	5248	5348	5269	5373	-2599	6186	1206
3	5126	5278	5157	5316	-2525	4998	1081
4	5061	5264	5102	5315	-2480	4954	1025
5	5016	5271	5068	5335	-2444	4181	941

Table 2. Socio-Demographic and Economic Characteristics of Mothers with Children Less Than 24 Months of Age in West Guji Zone.

Variables	Categories	Frequency	Percent (%)
Age of mothers	≤25 years	74	18.7
	26-34 years	255	64.6
	≥35 years	66	16.7
Residence	Urban	191	48.4
	Rural	204	51.6
Educational status of mothers	No formal education	208	52.7
	Primary	114	28.9
	Secondary and above	73	18.5
Educational status of husband	No formal education	209	52.9
	Primary	92	23.3
	Secondary and above	94	23.8
Maternal occupation	Farmer	172	43.5
	Housewife	161	40.8
	Others ^a	62	15.7
Occupation of husband	Farmer	191	48.4
	Government employee	79	20.0
	Daily laborer	83	21.0
	Others [#]	42	10.6
Marital status	Married	374	94.7
	Others	27	5.3
Family size	<5	166	42.0
	≥5	229	58.0
Household food security status	Food secured	62	15.7
	Mild food insecure	63	15.8
	Moderate food insecure	241	61.5
	Severe food insecure	27	6.8
Sex of the child	Male	178	45.1
	Female	217	54.9
Age of the child	<6 months	51	12.9
	6-8 months	35	8.9
	9-11 months	106	26.8
	12-17 months	86	21.8
	18-23 months	117	29.6

^aStudent, private employee, government employee, merchant; # NGO, merchant, private employee.

children whose mothers never used family planning methods (OR = 2.39, 95% CI: 1.019, 4.837), and children who never

received breast milk (OR = 3.83, 95% CI: 1.28, 8.95) have a higher risk of stunting. Additionally, severe food insecurity in the household (OR = 9.45, 95% CI: 1.367, 36.09) also increases the risk of stunting.

Categorization of Child Stunting Prevalence and Heterogeneity in Risk Factors

The overall prevalence of stunting in children aged 24 months was 47.34% (95% CI: 42.44%, 52.29%). However, our LCA identified 3 distinct subgroups of children under <2 years old with heterogeneity or varying risk profiles for stunting. Following the identification of latent classes, children were categorized into 3 classes based on their posterior probabilities of belonging to each category and subsequently analyzed and outlined using the conditional item-response probabilities. Table 6 shows the conditional probabilities of each latent class and the latent class prevalence. The classes differed in terms of the underlying factors that contributed to their risk of stunting. Each latent class is explained below according to its attributes.

Class 1: Low Risk with High Awareness and Practice of Hygiene and Nutrition

This class represents the smallest proportion of 23.8% of the children and likely faces the lowest risk of stunting. The class collectively indicates a group that is well-informed about and actively practices key hygiene and nutrition-related behaviors, which can greatly reduce the risk of childhood stunting. Nearly all mothers in this class wash their hands before feeding children and after using the toilet. Family planning is also widely used, and most mothers possess good nutritional knowledge. While access to safe water is not universal, a good portion of households has it. Compared to the other classes, two-fifth of the mothers is employed. Most importantly, these mothers prioritize their children's health. The space births effectively, with high rates of antenatal care and early breastfeeding initiation. They also ensure food security and a varied diet within their households. The low prevalence of stunting (only 23.4%) in this class is a testament to the positive impact of these combined factors.

Class 2: High Risk With Socioeconomic Disadvantage and Limited Maternal and Child Healthcare Access

This is the largest group, comprising nearly half the children (47.1%). They are likely at a higher risk of stunting compared

Table 3. Maternal Obstetrics and Reproductive Health and Infant and Young Child Feeding Characteristics in West Guji Zone, Southern Ethiopia.

Variables	Categories	Frequency	Percent (%)
Birth interval in years	<2	212	53.7
	≥2	183	46.3
History of ANC visit	Yes	349	88.4
	No	46	11.6
Ever use of family planning	Yes	249	63.0
	No	46	37.0
Have you received Nutrition Counseling service during ANC	Yes	303	76.7
	No	92	23.3
Do you have PNC	Yes	222	56.2
	No	173	43.8
Current breastfeeding status (n = 395)	Yes	313	79.2
	No	82	20.8
How long after birth, did you first put the child to breastfeed?	Within 1 h	286	72.4
	After 1 h	109	27.6
Is the child exclusively breastfed?	Yes	173	50.3
	No	171	49.7
Has the child started complementary feeding (n = 344)	Yes	282	82.0
	No	62	18.0
Does the child continue breastfeeding after 6 months	Yes	254	73.8
	No	90	26.2
Did your child consume breast milk yesterday day and night? (n = 395)	Yes	254	73.8
	No	90	26.2
Did your child consume Grains, white roots, and tubers (n = 282)	No	112	39.7
	Yes	170	60.3
Legumes and nuts (n = 282)	No	173	61.3
	Yes	109	38.7
Milk and dairy foods (n = 282)	No	131	46.5
	Yes	151	53.5
Flesh foods (n = 282)	No	199	70.6
	Yes	83	29.4
Eggs (n = 282)	No	138	48.9
	Yes	144	51.1
Vitamin A, fruits, and vegetables (n = 282)	No	158	56
	Yes	124	44
Other fruits and vegetables (n = 282)	No	141	50
	Yes	141	50
Minimum meal frequency (MMF) (n = 282)	<3 times a day	83	24.1
	≥3 times a day	199	75.9
Dietary diversity (n = 282)	≥4 food groups	110	39
	<4 food groups	172	61
Overall maternal food and nutrition knowledge status	Poor	203	51.4%
	Good	192	48.6%

to Class 1 due to the presence of various risk factors. The majority of mothers within this group are unemployed, reflecting tight

Table 4. Water, Sanitation, and Hygiene (WASH) Practices Related to the Characteristics of Participants.

Variables	Categories	Frequency	Percent
Drinking water source	Improved sources^a	311	78.7
	Unimproved sources^b	84	21.3
Distance from drinking water source	Less than 30 min	256	90.1
	Greater or equal to 30 min	39	9.9
Do you treat water before drinking	Yes	184	46.6
	No	211	53.4
What type of water storage you are using?	Jerrycan	305	77.2
	Bucket	58	14.7
	Pot	24	6.1
	Others	8	2.0
Hand washing before food preparation	No	127	32.2
	Yes	268	67.8
Wash hands before feeding a child	No	105	26.6
	Yes	290	73.4
Hand washing after cleaning animal dung	No	151	38.2
	Yes	244	61.8
Hand washing after visiting a toilet	No	107	27.1
	Yes	288	72.9
Do you have a latrine	Yes	272	68.9
	No	123	31.1
Type of latrine	Improved ^c	342	86.6
	Unimproved ^d	53	13.4

^aPublic tap, protected spring, pond, well, private tap.

^bUnprotected spring, pond, well, and river.

^cFlush/pour flush toilets to piped sewer systems, septic tanks, and pit latrines; ventilated improved pit (VIP) latrines; pit latrines with slabs.

^dOpen field, shared latrine.

household finances and limited resources to meet basic needs. Typically, these mothers have never utilized family planning methods, leading to a high occurrence of short birth intervals among women, which can result in maternal nutrient depletion and hinder child growth. Despite over 80% of women attending antenatal care, a considerable portion, exceeding 20%, lacks this crucial preventive healthcare service. Poor nutrition and a lack of food knowledge are widespread among these mothers, affecting their ability to provide optimal diets for their children. Moreover, nearly all households experience food insecurity, with approximately 90% of children having insufficient dietary diversity, consequently exacerbating potential nutritional deficiencies. A significant 66.7% of the children in this group suffer from stunted growth.

Class 3: Mixed Risk With a Combination of Positive and Negative Factors

This group falls in between the other two in size (29.1%), and their risk of stunting also likely falls somewhere between Class 1 (lowest) and Class 2 (highest). This class presents a complex picture, defying easy categorization due to a mix of positive “Good Nutrition and Early Breastfeeding Practices” and negative

Table 5. Result of Bayesian Logistic Regression Analysis of Stunting Among <24-Month Children.

Variables	Category	OR	SD	MCSE	Equal 2-tailed (95% credible interval)	
Maternal food and nutrition knowledge	Good	0.0229	0.014	0.0011	0.0049	0.0618
	Poor	1				
Birth interval	≥2 years	0.245	0.098	0.0043	0.105	0.480
	<2 years	1				
Ever use of family planning	Yes	1				
	No	2.39	0.983	0.034	1.019	4.837
Received nutrition counseling service during ANC	No	1.894	0.880	0.034	0.729	4.073
	Yes	1				
Initiation of breastfeeding time	After 1 h	0.627	0.340	0.0152	0.208	1.501
	Within 1 h	1				
Mother educational status	No formal education	1				
	Primary	0.265	0.1221	0.004	0.099	0.563
	≥Secondary	0.448	0.221	0.006	0.151	1.006
Age of child in months	<6	1				
	6-11	0.604	0.524	0.0763	0.118	1.888
	12-23	0.691	1.323	0.189	0.0376	3.293
Source of water	Improved	1				
	Unimproved	1.269	0.644	0.0208	0.427	3.026
Food security status	Secured	1				
	Mildly insecure	1.946	1.664	0.117	0.404	6.170
	Moderately insecure	0.412	0.257	0.022	0.0996	1.062
	Severely insecure	9.45	9.879	0.523	1.362	36.09
Treating water before drinking	Yes	1				
	No	109.44	78.51	6.150	32.67	303.01
Previous history of ANC visit	Yes	1				
	No	8.23	4.53	7.13	2.72	20.18
Hand washing after visiting a toilet	No	1				
	Yes	0.119	0.070	0.004	0.033	0.299
Had the child ever breastfed	Yes	1				
	No	3.83	2.04	0.096	1.28	8.95
The child is currently breastfeeding	Yes	1				
	No	0.461	0.273	0.015	0.132	1.162
Dietary diversity score	<4 food groups	1.814	0.606	0.0312	0.903	3.191
	≥4 food groups	1				
Complementary feeding started	No	2.782	1.724	0.079	0.814	7.300
	Yes	1				
Constant		1.35	1.553	0.159	0.191	4.638

“Economic Deprivation and Resource Insecurity” indicators impacting child stunting. While most mothers in this group practice good hygiene (washing hands before feeding and after using the toilet), many also utilize family planning and prioritize health-care through high rates of antenatal care and early breastfeeding. However, this progress is challenged by widespread unemployment, limited access to safe water, and inadequate dietary diversity in over half of their children’s diets. This combination, despite some positive behaviors, results in a significant proportion of 35.7% of children suffering from stunting, suggesting underlying nutritional deficiencies or chronic health problems.

Discussion

In our study, the overall prevalence of stunting in children under 24 months of age was 47.34%. However, our nuanced subgroup

analysis findings reveal a significant disparity in stunting prevalence among the identified subgroups, emphasizing the importance of understanding the unique characteristics and vulnerabilities of each class. Accordingly, Class 1, labeled as ‘Low-Risk,’ comprised 23.8% of the children and exhibited the lowest prevalence of stunting at 23.4%. Class 2, labeled as ‘High-Risk,’ comprised 47.1% of the children and had a significantly higher prevalence of stunting at 66.7%. Class 3, categorized as ‘Mixed-Risk,’ had a moderate stunting prevalence of 35.7%. Remarkably, more than half, that is, 51.4%, of mothers had poor knowledge of food and nutrition. Likewise, 53.4% of mothers had unsafe drinking water. Similarly, the majority of children (61%) had poor dietary diversity. On top of that, only 15.7% of households had food security. Notably, our findings highlight the importance of early prevention and intervention strategies to mitigate the long-term consequences of stunting, particularly among high-risk subgroups.

Table 6. Characteristics of a 3-Group Latent Class Analysis Based on the Estimated Probabilities of Each Risk Factor Within Each Latent Class.

Indicators	Categories	Latent classes		
		Class 1: Low-Risk	Class 2: High-Risk	Class 3: Mixed-Risk
Prevalence (Y)		0.238	0.471	0.291
Conditional probabilities (p)				
Mothers education	No formal education	0.328	0.555	0.682
	Formal education	0.672	0.445	0.318
Hand washing before child feeding	No	0.0347	0.5262	0.0535
	Yes	0.965	0.474	0.947
Hand washing after toilet visit	No	0.0889	0.5153	0.0377
	Yes	0.911	0.485	0.962
Ever used family planning	No	0.238	0.463	0.341
	Yes	0.762	0.537	0.659
Mothers' nutrition and food knowledge	Poor	0.195	1	0.008
	Good	0.805	0	0.992
Safe drinking water	No	0.399	0.471	0.771
	Yes	0.601	0.529	0.229
Mothers employment	Unemployed	0.589	0.944	0.92
	Employed	0.411	0.056	0.08
Have long birth intervals	≤2 years	0.485	0.618	0.451
	>2 years	0.515	0.382	0.549
Antenatal care follow-up	No	0.206	0.212	0.119
	Yes	0.794	0.788	0.881
Early initiation of Breastfeeding	After 1 h	0	0.468	0.223
	<1 h	1	0.532	0.777
Food security	Insecure	0.461	0.971	1
	Secured	0.539	0.029	0
Adequate dietary diversity	No	0.377	0.87	0.515
	Yes	0.623	0.13	0.485

The overall prevalence of stunting among under 24 months was comparable with a study conducted in Kenya, where 51% of children were also reported to be stunted.⁴⁰ The similarity in these findings may be attributed to shared demographic and socioeconomic characteristics among the communities studied. Contrastingly, several other studies have reported lower prevalence rates of stunting in young children. For instance, research conducted in Egypt found a rate of 20.3%,⁴¹ while in Indonesia, the prevalence was reported to be 33.7%.⁴² Similarly, studies in Southern Ethiopia⁴³ and Mumbai (India)⁴⁴ and a systematic review and meta-analysis pooled estimate in Ethiopia⁷ reported rates of 37%, 38%, and 35.1%, respectively. Despite the geographical disparities in prevalence indicating potential variations in underlying risk factors across populations, it is noteworthy that all studies mentioned above uniformly classify the prevalence levels as 'very high,' with the sole exception of Egypt at 20.3%, which falls within the 'high' category, as defined by the WHO.⁶ Therefore, this highlights the need for a standardized response

to the public health crisis that is tailored to address the specific risk factors in each location.

However, our LCA successfully identified 3 distinct subgroups of children under <2 years old with varying stunting risk profiles. The "Class 1: Lower risk" exhibits the most remarkable combination of positive maternal behaviors, good nutritional practices, and access to healthcare. These factors work synergistically to create an environment that significantly reduces the risk of stunting in children.^{8,45,46} However, despite having a positive indicator, the observed moderate prevalence of food insecurity and limited safe water access suggest areas for potential improvement. Interestingly, this subgroup has a much lower stunting rate (23.4%) compared to other groups. This suggests that their high awareness and practice of hygiene and nutrition may be making a real difference. This indicates that implementing education and behavior changes focusing on hygiene and nutrition can effectively decrease the prevalence of child stunting.^{46,47} More importantly, in this class, mothers can serve as advocates and peer educators, sharing their knowledge and experiences with mothers in higher-risk groups to improve children's eating habits.⁴⁸

Conversely, Class 2 is characterized by socioeconomic disadvantages and a lack of maternal and child healthcare access. This Class of women has less formal education, limited access to family planning and safe water, lower breastfeeding initiation rates, and maternal unemployment, which are leading to higher stunting rates. Unsurprisingly, this class of children experiences a high rate (66.7%) of stunting. Our study result supported the existing research on the intricate interplay of various factors causing stunting in Class 2 children.^{23,40,49,50} The finding implies that interventions aimed at enhancing the health of mothers and children who are struggling with various challenges are crucial.⁵¹ Although LCA does not directly identify individual households within particular latent classes,⁵² it can identify similar groupings based on available data. Thus, stakeholders had better focus on this high-risk group by screening and tracking mothers with similar risk profiles to latent classes. Community health extension workers (CHEWs) play a vital role in rural communities and often have a good understanding of local households,⁵³ can identify mothers with similar risk profiles during home visits or health post visits, and may also administer screening questionnaires during ANC visit for key indicators and can consider for identifying mothers with similar risk profiles.

Class 3 is a group that displayed both higher and lower stunting risk characteristics, indicating a complex interplay of factors influencing their stunting. This complexity can make it difficult to determine the overall risk level for stunting and the most effective interventions to address the situation.⁴ Despite good nutrition knowledge, high antenatal care and early breastfeeding can mitigate stunting risk factors,^{19,54-56} however, economic hardship, food insecurity, low family planning utilization, and limited access to diverse foods contribute to high stunting prevalence.^{23,57,58} Analyzing the distribution of positive and negative factors within Class 3 could provide further insights for intervention design. This suggests the

potential for nutrition-focused behavior change communication intervention to promote hygiene practices, address food insecurity through social safety nets programs, and promote consistent family planning use can be effective strategies for Class 3 in resource-constrained settings.

In our study, we found that a child living in a household without access to clean drinking water has an increased chance of stunting. This finding is consistent with different studies conducted in Ethiopia.^{19,21,59} We also found that maternal hand-washing practice after visiting a toilet is a significant protective factor of child stunting. This finding aligns in diverse settings, including Indonesia,⁶⁰ Ethiopia,^{19,21,61} Bangladesh,⁵⁵ and Kenya.⁶² The similarity suggests a potential mechanism to prevent the transmission of parasites and enteric bacteria, which cause diarrheal diseases and stunting in children.⁴⁶ Therefore, this result implies that, in order to lower childhood stunting, public health interventions should prioritize hand washing and access to clean water through behavioral modification, infrastructural development, and water quality monitor. We also found that children with a long birth interval (≥ 2 years) had a lower likelihood of stunting, which is consistent with numerous studies.^{18,63,64} Moreover, stunting is higher in children born to a woman who had never used family planning and had no antenatal care follow-up history. Consistent with this, numerous studies^{18,63,64} have shown that short birth intervals are associated with an increased risk of stunting, while longer intervals are associated with a decreased risk.^{44,65,66} This might be due to the fact that consecutive pregnancies spaced closely together reduces the maternal recovery time and nutrient replenishment.⁶⁷ This suggests that increasing the demand for family planning use is necessary to ensure the best possible health and development of children throughout crucial times.

Our study reveals a significant link between maternal food and nutritional knowledge and child stunting, with women with better knowledge having a 77% lower risk. This finding is consistent with previous studies that found a significant relationship between stunting and maternal nutritional knowledge.^{41,68-72} This might be because nutrition knowledge is crucial for selecting nutritious foods for children, promoting healthy growth,^{73,74} and equipping mothers to feed on time, prepare healthy meals, and identify early disease signs.⁷⁵ This result implies the need for overarching nutrition education programs for pregnant mothers, particularly those who are undernourished, to improve their knowledge and practices regarding child feeding and nutrition. In our study, mothers with at least primary education have a lower likelihood of having a stunted child, which is in line with previous research.^{7,63,64,74,76} This might be due to improved health-seeking behavior, nutrition knowledge, and child growth.^{73,77} Likewise, maternal education significantly reduces the stunting risk by providing opportunities for healthy practices during pregnancy and lactation,⁷⁷ boosting self-efficacy, financial independence, health-seeking behavior, and enhancing knowledge about healthcare and child nutrition.⁷³ By contrast, children born from households with severe food insecurity were more likely to be stunted, which is consistent with previous

studies conducted in Ethiopia and Southeast Asia.^{57,78,79} The similarities might be due to the fact that food insecurity is linked to child stunting due to limited dietary diversity, increased susceptibility to infections, and inadequate nutrient intake. Thus, addressing this issue and promoting dietary diversity are crucial for optimal the child's growth and development.

Our study employed advanced statistical models (Bayesian logistic regression and LCA) to identify groups of children sharing similar stunting risk factors. This approach provides a deeper understanding of the issue, allows for targeted interventions based on stunting risk profiles (Low-Risk, Mixed-Risk, and High-Risk), and this approach enables subgroup analysis. However, we acknowledge the limitations inherent to LCA. While it effectively identifies at-risk groups, it does not pinpoint the exact risk factors defining each group. Further research is needed to shed light on these specific characteristics. Additionally, the study's design (cross-sectional and institution-based) limits generalizability to other populations. Finally, we used a relatively small sample size, which could potentially impact the reliability of the results, as LCA usually requires larger datasets for accurate risk group identification.

Conclusion

This study identified a concerning high stunting prevalence of stunting among under 2-year-old children, disproportionately impacting vulnerable populations. Notably, factors like good nutritional knowledge, longer birth intervals, maternal education, and consistent hand-washing practices were associated with a significant reduction in stunting odds. Conversely, lack of antenatal care, family planning services, access to safe drinking water, and consistent breastfeeding, coupled with food insecurity, emerged as significant risk factors. On top of these all, our nuanced subgroup analysis revealed previously undetected patterns within the stunted population. Three distinct stunting risk profiles were identified: "Low-Risk Group" exhibiting consistently positive behaviors; "High-Risk Group" facing multiple challenges; and "Mixed-Risk Group" demonstrating a combination of both positive and negative factors. The identification of these distinct latent classes underscores the crucial need to account for heterogeneity in risk factors when designing interventions. Therefore, healthcare practitioners should conduct screenings for stunting, provide nutrition counseling, promote breastfeeding, and educate families on nutrition and healthcare access. Policymakers should enhance food access, strengthen social safety nets, and support primary education. Moreover, future research should focus on elucidating the biological and social mechanisms driving these disparities in stunting prevalence.

Acknowledgments

We would also like to extend our deepest gratitude to Bule Hora University Research and Publication Directorate for the financial support to conduct this research. We would also like to express our great appreciation to the data collectors, supervisors, and all sincere study participants for their voluntarism and willingness to participate in our study.

Author Contributions

AF, BA, and JB: conceived, designed the study, supervised data collection, performed analysis, and data interpretation, drafted the manuscript, and approved the final manuscript. DH, CAK, MS: supervised data collection, performed statistical analysis, interpreted data, and approved the final manuscript.

Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.


Ethics Approval and Consent to Participate

All methods were performed according to the relevant ethical guidelines and principles of the Declaration of Helsinki. Accordingly, an ethical approval letter was obtained from the Bule Hora University, Institute of Health, and Institutional Ethical Review Committee (Ref. No: I/O/H/I/R//B/023/14). Informed written consent was obtained from all respondents after an explanation of the purpose of the study. Confidentiality of the participant's response was maintained by excluding personal information, such as the patient's name or card number throughout the study. All study participants were encouraged to participate in the study, and at the same time, they were also informed that they had the right not to participate.

Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Bule Hora University financially supported the study.

ORCID iD

Anteneh Fikrie  <https://orcid.org/0000-0002-9809-3326>

Data Availability Statement

Data essential for the conclusion are included in this manuscript. Additional data can be obtained from the corresponding author upon reasonable request.

Supplemental Material

Supplemental material for this article is available online.

References

1. Organization WH. *Global nutrition targets 2025: Stunting policy brief*. World Health Organization, 2014.
2. Black RE, Victora CG, Walker SP, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet*. 2013;382(9890):427–451.
3. Dewey KG, Begum K. Long-term consequences of stunting in early life. *Matern Child Nutr*. 2011;7(Suppl. 3):5–18.
4. Organization WH. *Reducing stunting in children: equity considerations for achieving the global nutrition targets 2025*. 2018.
5. Levels and trends in child malnutrition: UNICEF/WHO/World Bank Group Joint Child Malnutrition Estimates: key findings of the 2023 edition. World Health Organization, United Nations Children's Fund (UNICEF) & World Bank, 2023. 978-92-4-007379-1.
6. Unicef, WHO W. *Levels and trends in child malnutrition: key findings of the 2019 Edition of the Joint Child Malnutrition Estimates*. World Health Organization; 2020.
7. Kassaw A, Kassie YT, Kefale D, et al. Pooled prevalence and its determinants of stunting among children during their critical period in Ethiopia: A systematic review and meta-analysis. *PloS One*. 2023;18(11):e0294689.
8. Gyampoh S, Otoo GE, Aryeetey RNO. Child feeding knowledge and practices among women participating in growth monitoring and promotion in Accra, Ghana. *BMC Pregnancy Childbirth*. 2014;14(1):1–7.
9. Kang Y, Kim S, Sinamo S, Christian P. Effectiveness of a community-based nutrition programme to improve child growth in rural Ethiopia: A cluster randomized trial. *Matern Child Nutr*. 2017;13(1):1–15.
10. Bilal SM, Moser A, Blanco R, Spigt M, Dinant GJ. Practices and challenges of growth monitoring and promotion in Ethiopia: A qualitative study. *J Health Popul Nutr*. 2014;32(3):441–451.
11. UNICEF. *Revisiting growth monitoring and its evolution to promoting growth as a strategic program approach: building consensus for future program guidance*. Report of the technical consultation on growth monitoring and promotion Book revisiting growth monitoring and its evolution to promoting growth as a strategic program approach: building consensus for future program guidance Report of the technical consultation on growth monitoring and promotion. 2007.
12. Usman MA, Kornher L, Sakketa TG. Do non-maternal adult female household members influence child nutrition? Empirical evidence from Ethiopia. *Matern Child Nutr*. 2021;17(Suppl 1): e13123. 1–12. doi:10.1111/mcn.13123.
13. ICF; EPHIEE. *Ethiopia Mini Demographic and Health Survey 2019: Final Report*. Rockville, Maryland, USA: EPHIICF, May 2021.
14. (MOH); FMoH, (EPHI); EPHI, UNICEF. *National food and nutrition strategy baseline survey: Key findings preliminary report*. Addis Ababa: Ethiopia Federal Ministry of Health March 2023.
15. Tekeba B, Tarekegn BT, Zegeye AF, Ayele AD. Stunting disparities and its associated factors among preschool children of employed and unemployed mothers in Gondar City: A comparative community-based cross-sectional study. *Front Nutr*. 2023;10:1172501.1-13.
16. Safitri NI, Noor NN, , Wahiduddin, Amiruddin R, Jafar N, Mallongi A. Risk factors of stunting in children aged 0-23 months in Katumbangan Health Center, Indonesia. *Pharmacognosy Journal*. 2023;15(5):851–855.
17. Roesler AL, Smithers LG, Wangpakapattanawong P, Moore V. Stunting, dietary diversity and household food insecurity among children under 5 years in ethnic communities of northern Thailand. *J Public Health*. 2019;41(4):772–780.
18. Lewa AF, Kusika SY, Muliani R, Jannah I. Risk factors of stunting events in child 6–23 months old in Biromaru Public Health Center Sigi. *Enfermeria Clinica*. 2020;30(Suppl 4):131–135.
19. Kwami CS, Godfrey S, Gavilan H, Lakanpaul M. Water, Sanitation, and hygiene: Linkages with stunting in rural Ethiopia. *Int J Environ Res Public Health*. 2019;16(20):1–21.
20. Kubeka Z, Modjadji P. Association of stunting with socio-demographic factors and feeding practices among children under

- two years in informal settlements in Gauteng, South Africa. *Children (Basel)*. 2023;10(8):1–15. <https://doi.org/10.3390/children10081280>.
21. Bekele T, Rahman B, Rawstorne P. The effect of access to water, sanitation and handwashing facilities on child growth indicators: Evidence from the Ethiopia demographic and health survey 2016. *PLoS One*. 2020;15(9):e0239313.
 22. Banerjee K, Dwivedi LK. Disparity in childhood stunting in India: Relative importance of community-level nutrition and sanitary practices. *PLoS One*. 2020;15(9):e0238364.
 23. Green MA, Corsi DJ, Mejía-Guevara I, Subramanian SV. Distinct clusters of stunted children in India: An observational study. *Matern Child Nutr*. 2018;14(3):e12592.
 24. Uwiringiyimana V, Veldkamp A, Amer S. Stunting spatial pattern in Rwanda: an examination of the demographic, socio-economic and environmental determinants. *Geospat Health*. 2019;14(2):329–339.
 25. Weller BE, Bowen NK, Faubert SJ. Latent class analysis: A guide to best practice. *J Black Psychol*. 2020;46(4):287–311.
 26. De Onis M, Onyango A, Borghi E, Siyam A, Blössner M, Lutter C. Worldwide implementation of the WHO child growth standards. *Public Health Nutr*. 2012;15(9):160310.
 27. Marias Y, Glasauer P. *Guidelines for assessing nutrition-related knowledge, attitudes and practices*. Food and Agriculture Organization of the United Nations (FAO). 2014.
 28. Gezimu W, Bekele F, Habte G. Pregnant mothers' knowledge, attitude, practice and its predictors towards nutrition in public hospitals of Southern Ethiopia: A multicenter cross-sectional study. *SAGE Open Med*. 2022;V 10:110. doi:10.1177/20503121221085843.
 29. Coates J, Swindale A, Bilinsky P. Household Food Insecurity Access Scale (HFAS) for measurement of food access: indicator guide: version 3. 2007.
 30. Federal Democratic Republic of Ethiopia Ministry of Health, Ethiopian Public Health Institute. Ethiopia: Food-Based Dietary Guidelines—2022. Addis Ababa, Ethiopia, March 2022.
 31. Kasajja M, Nabiwemba E, Wamani H, Kamukama S. Prevalence and factors associated with stunting among children aged 6–59 months in Kabale District, Uganda. *BMC Nutr*. 2022;8(1):1–7.
 32. Khorasani EC, Peyman N, Esmaily H. Measuring maternal health literacy in pregnant women referred to the healthcare centers of Mashhad, Iran, in 2015. *J Midwifery Reproductive Health*. 2018;6(1): 1157–1162. doi:10.22038/JMRH.2017.9613.
 33. (UNICEF) UNCSF. *UNICEF Conceptual framework on maternal and child nutrition*. New York, NY 10017 UNICEF: Nutrition and Child Development Section, Programme Group; November 2021. p. 4.
 34. WHO Guidelines Approved by the Guidelines Review Committee. *WHO child growth standards and the identification of severe acute malnutrition in infants and children: A joint statement by the World Health Organization and the United Nations Children's Fund*. World Health Organization Copyright © World Health Organization 2009.
 35. Organization WH. Recommendations for data collection, analysis and reporting on anthropometric indicators in children under 5 years old. 2019.
 36. Navarro DJ, Foxcroft DR. Learning statistics with Jamovi: A tutorial for psychology students and other beginners. (Version 0.70), 2019.
 37. Nylund KL, Asparouhov T, Muthén BO. Deciding on the number of classes in latent class analysis and growth mixture modeling: A Monte Carlo simulation study. *Struct Eq Model, Multidiscipl J*. 2007;14(4):535–569.
 38. Aflaki K, Vigod S, Ray JG. Part II: A step-by-step guide to latent class analysis. *J Clin Epidemiol*. 2023;159:348–351.
 39. Muthén LK, Muthén BO. Mplus user's guide: statistical analysis with latent variables: User's guide: Muthén & Muthén; 2004.
 40. Adeladza AT. The influence of socio-economic and nutritional characteristics on child growth in Kwale District of Kenya. *Afr J Food Agric Nutr Dev*. 2010;9(7):1570–1590.
 41. Seedhom AE, Mohamed ES, Mahfouz EM. Determinants of stunting among preschool children, Minia, Egypt. *Int Public Health Forum January*. 2014;1(2):5–9.
 42. Titaley CR, Ariawan I, Hapsari D, Muasyaroh A, Dibley MJ. Determinants of the stunting of children under two years old in Indonesia: A multilevel analysis of the 2013 Indonesia Basic Health Survey. *Nutrients*. 2019;11(5):1–13.
 43. Mengesha A, Hailu S, Birhane M, Belay MM. The prevalence of stunting and associated factors among children under five years of age in southern Ethiopia: Community based cross-sectional study. *Ann Glob Health*. 2021;87(1):111.
 44. Das S, Chanani S, Shah More N, Osrin D, Pantvaidya S, Jayaraman A. Determinants of stunting among children under 2 years in urban informal settlements in Mumbai, India: Evidence from a household census. *J Health Popul Nutr*. 2020;39(1):10.
 45. Headey D, Palloni G. Water, sanitation, and child health: Evidence from subnational panel data in 59 countries. *Demography*. 2019;56(2):729–752.
 46. Grembi JA, Lin A, Karim MA, et al. Effect of water, sanitation, handwashing, and nutrition interventions on enteropathogens in children 14 months old: A cluster-randomized controlled trial in rural Bangladesh. *J Infect Dis*. 2023;227(3):434–447.
 47. Girard AW, Olude O. Nutrition education and counselling provided during pregnancy: Effects on maternal, neonatal and child health outcomes. *Paediatr Perinat Epidemiol*. 2012;26(Suppl 1):191–204. doi:10.1111/j.1365-3016.2012.01278.x.
 48. Vairano M, D'Agnes P, Iavarone F, et al. Mothers as peer educators in a low socio-economic school setting in Southern Italy. 2018.
 49. Kibemo B, Mulugeta A, Hailu D, Gelaw B. The association of socio-demographic and environmental factors with stunting among under-five children in Hawassa City, Sidama National Regional State, Ethiopia. *J Nutr Sci*. 2022;11:e33.
 50. Alemayehu FR, Mariam RG, Loha E, Anato A, Desta DT. Maternal socio demographic characteristics are associated with child stunting in Alamura Subcity of Hawassa, Ethiopia. *Public Health Res*. 2020;10(1):12–20.
 51. Lassi ZS, Padhani ZA, Rabbani A, Rind F, Salam RA, Bhutta ZA. Effects of nutritional interventions during pregnancy on birth, child health and development outcomes: A systematic review of evidence from low- and middle-income countries. *Campbell Syst Rev*. 2021;17(2):e1150.

52. Fox B, Cochran JK, Escue M. Profiles of death penalty attitudes: A latent class approach. *Deviant Behav.* 2024;1–22. doi:10.1080/01639625.2024.2307962.
53. Amare Y, Scheelbeek P, Schellenberg J, Berhanu D, Hill Z. Early postnatal home visits: A qualitative study of barriers and facilitators to achieving high coverage. *BMC Public Health.* 2018;18(1):1–8.
54. Mahmoud NM, Ghaly AS. Dietary knowledge, practices and adequacy among Bedouin pregnant women. *Int J of Nurs.* 2019;6(2):68–83.
55. Nizame FA, Unicomb L, Sanghvi T, et al. Handwashing before food preparation and child feeding: A missed opportunity for hygiene promotion. *Am J Trop Med Hyg.* 2013;89(6):1179.
56. Mishra R, Bera S. Geospatial and environmental determinants of stunting, wasting, and underweight: Empirical evidence from rural South and Southeast Asia. *Nutrition.* 2024;120:112346. <https://doi.org/10.1016/j.nut.2023.112346>
57. Mulu E, Mengistie B. Household food insecurity and its association with nutritional status of under five children in Sekela District, western Ethiopia: A comparative cross-sectional study. *BMC Nutr.* 2017;3(1):35.
58. Gassara G, Lin Q, Deng J, Zhang Y, Wei J, Chen J. Dietary diversity, household food insecurity and stunting among children aged 12 to 59 months in N'Djamena—Chad. *Nutrients.* 2023;15(3):573.
59. Ademas A, Adane M, Keleb A, Berihun G, Tesfaw G. Water, sanitation, and hygiene as a priority intervention for stunting in under-five children in northwest Ethiopia: A community-based cross-sectional study. *Ital J Pediatr.* 2021;47(1):174.
60. Ahmadi LS, Azizah R, Oktarizal H. Association between toilet availability and handwashing habits and the incidence of stunting in young children in Tanjung Pinang City, Indonesia. *Indonesian Malaysian J Med Heal Sci.* 2020;16(2):215–218.
61. Baye K. Prioritizing the scale-up of evidence-based nutrition and health interventions to accelerate stunting reduction in Ethiopia. *Nutrients.* 2019;11(12):3065.
62. Null C, Stewart CP, Pickering AJ, et al. Effects of water quality, sanitation, handwashing, and nutritional interventions on diarrhoea and child growth in rural Kenya: A cluster-randomised controlled trial. *Lancet Glob Health.* 2018;6(3):e316–ee29.
63. Das S, Hossain M, Islam M. Predictors of child chronic malnutrition in Bangladesh. *Cell.* 2008;45(3):137–155.
64. Adekanmbi VT, Kayode GA, Uthman OA. Individual and contextual factors associated with childhood stunting in Nigeria: A multilevel analysis. *Matern Child Nutr.* 2013;9(2):244–259.
65. Chungkham HS, Sahoo H, Marbaniang SP. Birth interval and childhood undernutrition: Evidence from a large scale survey in India. *Clin Epidemiol Glob Health.* 2020;8(4):1189–1194.
66. Ahmed S, Kane TT, Khuda B-E, Levin ANN, Mozumder ABMKA. The effect of birth interval on malnutrition in Bangladeshi infants and young children. *J Biosoc Sci.* 2000;32(3):289–300.
67. Dewey KG, Cohen RJ. Does birth spacing affect maternal or child nutritional status? A systematic literature review. *Matern Child Nutr.* 2007;3(3):151–173.
68. Fadare O, Amare M, Mavrotas G, Akerele D, Ogunniyi A. Mother's nutrition-related knowledge and child nutrition outcomes: Empirical evidence from Nigeria. *PloS One.* 2019;14(2):e0212775.
69. Christian P, Abbi R, Gujral S, Gopaldas T. The role of maternal literacy and nutrition knowledge in determining children's nutritional status. *Food Nutr Bull.* 1988;10(4):16.
70. Maheri M, Bidar M, Farrokh-Eslamlou H, Sadaghianifar A. Evaluation of anthropometric indices and their relationship with maternal nutritional literacy and selected socio-economic and demographic variables among children under 5 years old. *Ital J Pediatr.* 2022;48(1):1–11.
71. Saaka M. Relationship between mothers' nutritional knowledge in childcare practices and the growth of children living in impoverished rural communities. *J Health Popul Nutr.* 2014;32(2):237–248.
72. Oly-Alawuba NMA, Ihedioha S. Nutritional knowledge of mothers/caregivers in relation to the anthropometric indices of children (2–5 years) in Obowu local government area, Imo State, Nigeria. *FASEB J.* 2018;4(1):19–23. <https://ijisset.org/storage/Volume4/Issue1/IJISSET-040102.pdf>
73. Abuya BA, Ciera J, Kimani-Murage E. Effect of mother's education on child's nutritional status in the slums of Nairobi. *BMC Pediatr.* 2012;12(1):80.
74. Vollmer S, Bommer C, Krishna A, Harttgen K, Subramanian S. The association of parental education with childhood undernutrition in low-and middle-income countries: Comparing the role of paternal and maternal education. *Int J Epidemiol.* 2017;46(1):312–323.
75. Mu'awiyyah Babale Sufiyan SS, Bashir AAU. Effect of maternal literacy on nutritional status of children under 5 years of age in the Babban-Dodo community Zaria City, Northwest Nigeria. *Ann Nigerian Med.* 2012;6(2):61.
76. Siddiqi MNA, Haque MN, Goni MA. Malnutrition of under-five children: Evidence from Bangladesh. *Asian J Med Sci.* 2011;2(2):113–119.
77. Chinnakotla B, Susarla SM, Mohan DC, Turton B. Associations between maternal education and child nutrition and oral health in an indigenous population in Ecuador. *Int J Environ Res Public Health.* 2022;20(1):1–14.
78. Paudel A, Bhandari TR, Dangi NB. Household food security and nutritional status of under-five-year children: A case study of Nepal. *J Nutr Food Sec.* 2021;6(2):179–184. <https://doi.org/10.18502/jnfs.v6i2.6073>
79. Nepali S, Simkhada P, Davies IG. Association between wasting and food insecurity among children under five years: Findings from Nepal demographic health survey 2016. *BMC Public Health.* 2020;20(1):1–7. doi:10.1186/s12889-020-09146-x.

Author Biographies

Anteneh Fikrie is an Assistant Professor of Public Health at Bule Hora University in Ethiopia. With over eleven years of experience, Mr. Fikrie has made significant contributions to public health and health-care management through teaching, research, and leadership. His research focuses on women's empowerment, nutrition, maternal and child health, and social epidemiology. Mr. Fikrie has authored over twenty three peer-reviewed publications and possesses expertise in quantitative and qualitative research methodologies. He currently

serves as an editor for PLOS ONE and assistant editor for the BHU Journal of Indigenous Knowledge and Development Studies.

Berhanu Adula is a lecturer at School of Public Health Institute of Health, Bule Hora University. He has completed his BSc degree in Public Health from Ambo University and Masters of Public in Public health Nutrition from Dilla University. Currently, he has offering Nutrition and related course for undergraduate students in Institute of Health, Bule Hora University. He is interested in multidisciplinary research approach using both qualitative and quantitative methods with focus on Maternal and child nutrition, Nutrition related Non-communicable diseases, food security, and other Nutrition Related researches.

Jitu Beka is a lecturer at Bule Hora University, Institute of Health, School of Public Health. She has earned BSc degree in Nursing from Addis Ababa University, and Masters in Human Nutrition from Jimma University. Currently, she has been offering Nutrition related courses for undergraduate students of Institute of Health, Bule Hora University. Her interest of research areas are primarily concerns of malnutrition, food security, Nutritional and metabolic diseases. She has two original research articles on the process of publication.

Dr. Dejene Hailu is an Associate Professor of Public Health at the School of Public Health, Hawassa University, Sidama Regional State, Ethiopia. He has held several leadership positions, including Coordinator of Research and Technology Transfer at the College of Medicine and Health Sciences (2015-2020), Chairperson of the Health Research Ethics Committee (2007-2009), and Dean of the Public Health Faculty (2004-2007). He currently serves as Associate Editor for the Journal of Science and Development.

Dr. Cheru Atsmegiorgis Kitabo an Assistant Professor of Statistics with extensive experience in statistics, data analysis, and survey methodology. With a strong background in leading large-scale research projects and managing complex datasets, Dr. Cheru has a proven track record in delivering high-quality research outputs. Expertise includes advanced statistical modeling, data management, and training. Holds a leadership role in quality assurance at Hawassa University.

Dr. Mark Spigt is an Associate Professor within the Department of Family Medicine at the Faculty of Health, Medicine and Life Sciences, Maastricht University, Netherlands. He holds expertise in implementing and evaluating innovations in primary care and public health. As of 1 January 2021, Dr Mark is a chair of the Research Line Optimising Patient Care (OPC).