

 Citation[Article](#)[Cited By \(3\)](#)[Tweatations \(28\)](#)[Metrics](#) Review

Effectiveness and Cost-Effectiveness of eHealth Interventions in Somatic Diseases: A Systematic Review of Systematic Reviews and Meta-Analyses

Niels J Elbert¹, MD; Harmieke van Os-Medendorp², PhD; Wilco van Renselaar³;

Anne G Ekeland^{4,5}, PhD; Leona Hakkaart-van Roijen^{6,7}, PhD;

Hein Raat⁸, MD, MBA, PhD; Tamar EC Nijsten¹, MD, PhD;

Suzanne GMA Pasmans^{1,9}, MD, PhD

¹Department of (Pediatric) Dermatology, Erasmus University Medical Center Rotterdam, Rotterdam, Netherlands

²Department of Dermatology & Allergology, University Medical Center Utrecht, Utrecht, Netherlands

³Patiënt1, Almere, Netherlands

⁴Norwegian Centre for Integrated Care & Telemedicine, University Hospital of North Norway, Tromsø, Norway

⁵Department of Clinical Medicine, Telemedicine & e-Health Research Group, University of Tromsø, Tromsø, Norway

⁶Institute for Medical Technology Assessment, Erasmus University Rotterdam, Rotterdam, Netherlands

⁷Institute of Health Policy and Management, Erasmus University Rotterdam, Rotterdam, Netherlands

⁸Department of Public Health, Erasmus University Medical Center Rotterdam, Rotterdam, Netherlands

⁹Department of Pediatric Dermatology, Wilhelmina Children's Hospital, University Medical Center Utrecht, Utrecht, Netherlands

 Export Metadata Download:

Sections

[Abstract](#)[Introduction](#)[Methods](#)[Results](#)[Discussion](#)[Abbreviations](#)[References](#)[Copyright](#)

ABSTRACT

Background: eHealth potentially enhances quality of care and may reduce health care costs. However, a review of systematic reviews published in 2010 concluded that high-quality evidence on the benefits of eHealth interventions was still lacking.

Objective: We conducted a systematic review of systematic reviews and meta-analyses on the effectiveness/cost-effectiveness of eHealth interventions in patients with somatic diseases to analyze whether, and to what possible extent, the outcome of recent research supports or differs from previous conclusions.

Methods: Literature searches were performed in PubMed, EMBASE, The Cochrane Library, and Scopus for systematic reviews and meta-analyses on eHealth interventions published between August 2009 and December 2012. Articles were screened for relevance based on preset inclusion and exclusion criteria. Citations of residual articles were screened for additional literature. Included papers were critically appraised using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Statement before data were extracted. Based on conclusions drawn by the authors of the included articles, reviews and meta-analyses were divided into 1 of 3 groups: suitable, promising, or limited evidence on effectiveness/cost-effectiveness. Cases of uncertainty were resolved by consensus discussion. Effect sizes were extracted from papers that included a meta-analysis. To compare our results with previous findings, a trend analysis was performed.

Results: Our literature searches yielded 31 eligible reviews, of which 20 (65%) reported on costs. Seven papers (23%) concluded that eHealth is effective/cost-effective, 13 (42%) underlined that evidence is promising, and others found limited or inconsistent proof. Methodological quality of the included reviews and meta-analyses was generally considered high. Trend analysis showed a considerable accumulation of literature on eHealth. However, a similar percentage of papers concluded that eHealth is effective/cost-effective or evidence is at least promising (65% vs 62%). Reviews focusing primarily on children or family caregivers still remained scarce. Although a pooled (subgroup) analysis of aggregate data from randomized studies was performed in a higher percentage of more recently published reviews (45% vs 27%), data on economic outcome

measures were less frequently reported (65% vs 85%).

Conclusions: The number of reviews and meta-analyses on eHealth interventions in patients with somatic diseases has increased considerably in recent years. Most articles show eHealth is effective/cost-effective or at least suggest evidence is promising, which is consistent with previous findings. Although many researchers advocate larger, well-designed, controlled studies, we believe attention should be given to the development and evaluation of strategies to implement effective/cost-effective eHealth initiatives in daily practice, rather than to further strengthen current evidence.

J Med Internet Res 2014;16(4):e110)

doi:10.2196/jmir.2790

KEYWORDS

eHealth; telehealth; telemedicine; review; program effectiveness; cost effectiveness

We also recommend

Effectiveness of Web-based Interventions on Patient Empowerment: A Systematic Review and Meta-analysis

David Samoocha et al., J Med Internet Res, 2010

Electronic Symptom Reporting Between Patient and Provider for Improved Health Care Service Quality: A Systematic Review of Randomized Controlled Trials. Part 1: State of the Art

Monika Alise Johansen et al., J Med Internet Res, 2012

Clinical Effects of Home Telemonitoring in the Context of Diabetes, Asthma, Heart Failure and Hypertension: A Systematic Review

Guy Paré et al., J Med Internet Res, 2010

Web 2.0 Chronic Disease Self-Management for Older Adults: A Systematic Review

Michael Stellefson et al., J Med Internet Res, 2013

Online Prevention Aimed at Lifestyle Behaviors: A Systematic Review of Reviews

Leonie F M Kohl et al., J Med Internet Res, 2013

Low-Intensity Self-Management Intervention for Persons With Type 2 Diabetes Using a Mobile Phone-Based Diabetes Diary, With and Without Health Counseling and Motivational Interviewing: Protocol for a Randomized Controlled Trial

Lis Ribu et al., JMIR Res Protoc, 2013

The Effectiveness of Web-Based vs. Non-Web-Based Interventions: A Meta-Analysis of Behavioral Change Outcomes

Dean J Wantland et al., J Med Internet Res, 2004

Effectiveness of E-Self-help Interventions for Curbing Adult Problem Drinking: A Meta-analysis

Heleen Riper et al., J Med Internet Res, 2011

Increasing Physical Activity With Mobile

The Contribution of Teleconsultation
and Videoconferencing to Diabetes
Care: A Systematic Literature Review
Fenne Verhoeven et al., J Med Internet
Res, 2007

Devices: A Meta-Analysis
Jason Fanning et al., J Med Internet
Res, 2012

Powered by  Trend MD

Introduction

Willem Einthoven started experiments in 1906 with remote consultations via the telephone network and this is when eHealth is likely to have seen first light [1]. It was not until the 1990s when the number of publications in this field of medicine increased dramatically [2]. This was because of the many studies that were carried out involving remote consultations through video-teleconferencing and digital images to give specialists comparable visual inspection of patients as referring doctors [3].

In modern medical practice, eHealth interventions are increasingly present. With nomenclature evolving rapidly, a significant overlap between terms such as *eHealth*, *telemedicine*, and *telehealth* has occurred. The American Telemedicine Association defines telemedicine as “the use of medical information exchanged from one site to another through electronic communications with the purpose of improving the health status of patients,” and considers *eHealth* and *telehealth* as interchangeable nouns. Both words encompass a broader definition of remote health care and also comprise related services, including nonclinical programs such as education, administration, and research [4]. However, telemedicine is a term that is generally reserved for clinical patient care applications [5].

McLean et al [6] conceptualized the definition of eHealth in a Cochrane review on telehealthcare for asthmatic patients as “the provision of personalized health care at a distance.” eHealth contains the following 3 key elements: (1) data obtained from the patient; (2) electronic transfer of data over a distance; and (3) patient-tailored feedback from a health care professional [5,6].

Therefore, communication in eHealth interventions is personalized and interactive in contrast to patient information websites on health and disease.

eHealth potentially enhances the quality of care and reduces health care costs. It may do so by providing patient education and counseling for primary prevention and early detection of disease, replacing face-to-face visits with health care professionals, collecting patient data on medical parameters remotely, among several other mechanisms [6,7]. Because eHealth interventions are considered complex interventions by the Medical Research Council, difficulty may arise in the assessment of the many interacting components of the intervention [8].

In 2010, Ekeland et al [9] published a systematic review of systematic reviews to evaluate the impact of eHealth interventions on health and health care costs. The authors concluded that high-quality evidence on health and economic benefits was still lacking despite the large number of publications. The primary objective of our review is to analyze whether, and to what possible extent, the outcome of recent research supports or differs from these previous conclusions on the effectiveness/cost-effectiveness of eHealth interventions in patients with somatic diseases.

Methods

Overview

Literature searches for systematic reviews and meta-analyses on the effectiveness/cost-effectiveness of eHealth interventions were performed in the following online databases: PubMed, EMBASE, The Cochrane Library, and Scopus. Two of the authors (NE, HO) independently screened all papers' titles and abstracts for relevance. Citations were screened through Web of Science for additional literature.

Search Queries

Similar to Ekeland et al [9], we used the following (simplified) search query to retrieve systematic reviews and meta-analyses on the effectiveness of eHealth interventions: "[eHealth] AND [effectiveness] AND [systematic review OR meta-analysis]." To search for papers on cost-effectiveness, "AND [costs]" was added to the aforementioned syntax. Because Ekeland et al [9] took into consideration published works from 2005 to July 2009, we limited our search results to articles published between August 2009 and December 2012. Extensive search queries are

presented in **Tables 1** and **2**.

Database	Search Query	Results	Excluded
PubMed	Systematic review OR meta-analysis AND eHealth AND somatic disease	45,698	45,698
EMBASE	Systematic review OR meta-analysis AND eHealth AND somatic disease	3,394	3,394
The Cochrane Library	Systematic review OR meta-analysis AND eHealth AND somatic disease	1,248	1,248
Scopus	Systematic review OR meta-analysis AND eHealth AND somatic disease	15,233	15,233
Pubmed	Systematic review OR meta-analysis AND eHealth AND somatic disease	45,698	45,698
EMBASE	Systematic review OR meta-analysis AND eHealth AND somatic disease	3,394	3,394
The Cochrane Library	Systematic review OR meta-analysis AND eHealth AND somatic disease	1,248	1,248
Scopus	Systematic review OR meta-analysis AND eHealth AND somatic disease	15,233	15,233
Pubmed	Systematic review OR meta-analysis AND eHealth AND somatic disease	45,698	45,698
EMBASE	Systematic review OR meta-analysis AND eHealth AND somatic disease	3,394	3,394
The Cochrane Library	Systematic review OR meta-analysis AND eHealth AND somatic disease	1,248	1,248
Scopus	Systematic review OR meta-analysis AND eHealth AND somatic disease	15,233	15,233
Pubmed	Systematic review OR meta-analysis AND eHealth AND somatic disease	45,698	45,698
EMBASE	Systematic review OR meta-analysis AND eHealth AND somatic disease	3,394	3,394
The Cochrane Library	Systematic review OR meta-analysis AND eHealth AND somatic disease	1,248	1,248
Scopus	Systematic review OR meta-analysis AND eHealth AND somatic disease	15,233	15,233

□ Table 1. PubMed, EMBASE, The Cochrane Library, and Scopus search queries for systematic reviews and meta-analyses on the effectiveness of eHealth interventions (search conducted on September 12, 2013).

[View this table](#)

Database	Search Query	Results	Excluded
PubMed	Systematic review OR meta-analysis AND eHealth AND somatic disease	45,698	45,698
EMBASE	Systematic review OR meta-analysis AND eHealth AND somatic disease	3,394	3,394
The Cochrane Library	Systematic review OR meta-analysis AND eHealth AND somatic disease	1,248	1,248
Scopus	Systematic review OR meta-analysis AND eHealth AND somatic disease	15,233	15,233
Pubmed	Systematic review OR meta-analysis AND eHealth AND somatic disease	45,698	45,698
EMBASE	Systematic review OR meta-analysis AND eHealth AND somatic disease	3,394	3,394
The Cochrane Library	Systematic review OR meta-analysis AND eHealth AND somatic disease	1,248	1,248
Scopus	Systematic review OR meta-analysis AND eHealth AND somatic disease	15,233	15,233
Pubmed	Systematic review OR meta-analysis AND eHealth AND somatic disease	45,698	45,698
EMBASE	Systematic review OR meta-analysis AND eHealth AND somatic disease	3,394	3,394
The Cochrane Library	Systematic review OR meta-analysis AND eHealth AND somatic disease	1,248	1,248
Scopus	Systematic review OR meta-analysis AND eHealth AND somatic disease	15,233	15,233

□ Table 2. PubMed, EMBASE, The Cochrane Library, and Scopus search queries for systematic reviews and meta-analyses on the cost-effectiveness of eHealth interventions (search conducted on September 12, 2013).

[View this table](#)

Inclusion Criteria

Systematic reviews and meta-analyses on eHealth interventions in adults and/or children with somatic diseases (ie, illnesses with a physical cause, not mental), and those focusing on family caregivers were included. Interventions had to meet the following 3 criteria: (1) data were obtained from the patient or family caregiver, (2) data were electronically transferred over a distance, and (3) personalized feedback was given from a health care professional. Reviews and meta-analyses of individual studies comparing eHealth interventions to usual or no care, and those comparing different eHealth initiatives were assessed. We only accounted for papers reporting health-related outcomes, costs, patient satisfaction, and/or self-management.

Exclusion Criteria

Those eHealth interventions that were not home-based (eg, tele-ICU) or not patient or family caregiver-oriented (eg, education of medical or nursing students and health care professionals) were excluded. We excluded meta-analyses that included nonrandomized studies (eg, cohort studies) unless a subgroup analysis of randomized studies (eg, randomized controlled trials, randomized crossover trials) was performed. In addition, we did not assess papers written in languages other than English or

Dutch, and those for which the full-text was not available online.

In contrast to Ekeland et al [9], we narrowed the focus of our work by excluding reviews and meta-analyses on nonsomatic disorders (eg, mental disorders such as anxiety, depression, schizophrenia, and posttraumatic stress disorder) and lifestyle changes (eg, smoking cessation and drug intervention programs) to increase the comparability of the included papers and to limit the search results.

Outcome Measures

Health-related effects (eg, morbidity, mortality, quality of life, hospitalization) and health care costs (eg, health care utilization) were defined as primary outcome measures. We considered patient satisfaction and self-management as secondary outcome measures.

Critical Appraisal

Before data were extracted, the included papers were critically appraised using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA, formerly QUOROM) Statement [10]. The PRISMA Statement provides an evidence-based 27-item checklist (eg, on objectives, methodology, and limitations) for reporting in systematic reviews and meta-analyses.


Data Extraction

Based on conclusions drawn by the authors of the included papers, all reviews and meta-analyses were divided into 1 of 3 groups: (1) suitable, (2) promising, or (3) limited evidence that eHealth is effective/cost-effective. Cases of uncertainty were resolved by consensus discussion between 2 authors of the current review (NE, HO). Effect sizes, such as standardized or weighted mean differences, relative risks, odds ratios, and z scores, were extracted from papers that included a pooled (subgroup) analysis of aggregate data from randomized studies. No attempt was made to contact authors for missing data. To analyze whether the results of the included papers supported or differed from previous findings by Ekeland et al [9], we performed a trend analysis using basic statistics.

Results


Search Results

The initial search yielded a total of 1657 articles, including 619 articles that reported on cost-related outcome measures (Figures 1 and 2). Following removal of duplicates and screening of the residual papers on preset inclusion and exclusion criteria, 30 eligible reviews remained [6,11-39], of which 19 reported on costs [6,13,14,16,18,19,21-24,26,28,29,31,33,34,37-39]. Subsequent citation screening through Web of Science resulted in 1 additional paper [40]. Thus, a total of 31 reviews were retrieved (Figure 1), of which 20 (65%) reported on costs (Figure 2). Three of 31 reviews (10%) reported primarily on children [28,37,38], and 1 of 31 (3%) focused on the effects of eHealth interventions on family caregivers [26].



□ Figure 1. Flow diagram of the literature search on the effectiveness of eHealth interventions.

[View this figure](#)



□ Figure 2. Flow diagram of the literature search on the cost-effectiveness of eHealth interventions.

[View this figure](#)

Results per article are summarized in 3 separate tables, 1 for systematic reviews and meta-analyses reporting eHealth interventions are effective/cost-effective (**Table 3**), a second table for papers showing evidence is promising (**Table 4**), and a third table with papers underlining evidence is lacking, limited, or inconsistent (**Table 5**). **Table 6** demonstrates the effect sizes-among other characteristics-reported in 14 reviews in which a pooled (subgroup) analysis of aggregate data from randomized studies was performed. All tables are presented subsequently.

Effectiveness/Cost-Effectiveness of eHealth Interventions

A total of 7 reviews (23%) showed eHealth interventions are effective on either health or cost-related outcome measures (**Table 3**) [11-17]. Study populations consisted of patients with congestive heart failure (CHF) [13-15,17], diabetes mellitus [12,16], and hypertension [11]. Types of interventions that were effective/cost-effective comprised home telemonitoring [11,13-17], Web or mobile phone-based education [12,16], structured telephone support [14-16], and mobile phone-assisted self-management programs [16]. Patient acceptance and satisfaction were generally considered high.

Pooled analyses were performed in each of the 7 reviews and demonstrated significant reduction of all-cause mortality, all-cause hospitalization, and CHF-related hospital admissions through home telemonitoring and structured telephone support in patients with CHF [13-15,17]. Home telemonitoring also resulted in significant improvement of systolic blood pressure and nonsignificant reduction of diastolic blood pressure, antihypertensive drug use, and therapeutic inertia (ie, unchanged medication despite elevated blood pressure) in hypertensive patients [11]. Web-based education and various mobile phone interventions led to significant improvement of laboratory parameters, such as glycosylated hemoglobin (HbA1c) and low-density lipoprotein (LDL) cholesterol, in diabetic patients [12,16].

Qualitative analysis of individual studies revealed several other positive effects of eHealth interventions, including economic benefits [14,16], reduction of the number of visits to outpatient clinics [12], increase of disease-related knowledge and self-management [12,14,16], and improvement of quality of life [13,14,17].

Author	Year	Number of Studies	Number of Patients	Number of Outcomes
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10

□ Table 3. Systematic reviews and meta-analyses in which eHealth interventions were shown to be effective/cost-effective.

[View this table](#)

Evidence on eHealth Interventions is Promising

Thirteen reviews (42%) were less confident about the effectiveness/cost-effectiveness of eHealth interventions [18-29,40], but suggested that these initiatives are promising or bear potential (Table 4). Many of the authors claim additional research is needed to clarify efficacy and cost-related issues. Pooled analyses were performed in 4 reviews and presented subsequently [22-24,27]. One review on chronic obstructive pulmonary disease (COPD) demonstrated the capacity of eHealth interventions to significantly reduce the number of patients with 1 or more emergency department visits or hospital admissions-due to exacerbation of pulmonary symptoms-over a 12-month period [23]. eHealth interventions did not significantly improve quality of life and all-cause mortality. Because the interventions were often part of complex interventions, the authors concluded that further investigation is required to determine the precise role of eHealth. Promising effects were also identified for Internet-based peer and clinical visit support programs—among several other eHealth interventions—in acute and chronic pain management [18,22]. Although the Internet was supportive in the treatment of pain, it remained unclear what benefits could be gained and which patients would profit most.

Author	Year	Number of Studies	Number of Patients	Number of Outcomes
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10
Chouinard et al.	2013	10	1,100	10

□ Table 4. Systematic reviews and meta-analyses in which promising evidence on the effectiveness/cost-effectiveness of eHealth interventions was reported.

[View this table](#)

Qualitative analysis of individual studies revealed many other promising effects of eHealth interventions, for example, Internet-based device-assisted remote monitoring systems in patients with cardiovascular implantable electronic devices [40], in-home

telerehabilitation in routine care of patients with stroke and other somatic diseases [20,21], technology-assisted training and support programs for family members of patients with traumatic brain injury [26], and Web-based education to increase patient empowerment [27]. Paré et al [25] assessed the clinical effects of home telemonitoring in patients with a variety of chronic diseases. The authors highlight the fact that home telemonitoring allows for closer follow-up of individual patients' conditions and for early detection of warning signs in case of health deterioration. However, they claim larger trials are needed to confirm the clinical effects of home telemonitoring.

Evidence on eHealth Interventions Is Lacking, Limited or Inconsistent

Eleven reviews (35%) underlined that evidence on the effectiveness/cost-effectiveness of eHealth interventions is still lacking, limited, or inconsistent (Table 5) [6,30-39]. In many articles, the poor methodological quality of individual studies is criticized, and ambiguous or conflicting findings are emphasized.

McLean et al [6] conducted a Cochrane review of 21 RCTs on a range of eHealth interventions in patients with asthma. Meta-analysis did not show a clinically important improvement of disease-specific quality of life, and no significant reduction of all-cause emergency department visits over a 12-month period was found (Table 6). The authors concluded that eHealth is unlikely to result in clinically relevant improvements of health-related outcome measures in patients with relatively mild disease, but does appear to have the potential to reduce all-cause hospital admissions in those with more severe disease.

Shulman et al [37] studied the impact of eHealth interventions involving transmission of blood glucose data in youth with type 1 diabetes mellitus. Pooled analyses showed no apparent effect of the interventions on HbA1c or acute complications, such as severe hypoglycemia and diabetic ketoacidosis (Table 6). The limited data available on patient satisfaction and costs also suggested no differences between the intervention and the comparison group.

Author	Year	Number of studies	Number of patients	Effectiveness	Cost-effectiveness
McLean et al [6]	2012	21	10,000	Lacking	Lacking
Shulman et al [37]	2013	10	1,000	Lacking	Lacking
Paré et al [25]	2013	10	1,000	Lacking	Lacking
...

□ Table 5. Systematic reviews and meta-analyses in which no, limited, or inconsistent evidence on the effectiveness/cost-effectiveness of eHealth interventions was reported.

[View this table](#)

Author	Year	Country	Population	Intervention	Outcome	Quality
McGeary et al [22]	2012	USA	Chronic pain patients	Telehealth	Pain management	High
McLean et al [6]	2012	UK	Asthma patients	Web-based tools	Asthma management	High
...

□ Table 6. Characteristics of 14 systematic reviews in which a meta-analysis was performed.

[View this table](#)

Methodological Quality of Reviews and Meta-Analyses

Among the systematic reviews and meta-analyses described in the current review are 4 high-quality Cochrane reviews [6,14,23,38]. Following the PRISMA Statement [10], the methodological quality of the other included papers was generally considered high. Nearly all authors provided search queries and selection criteria, described the process of data extraction, presented the results and limitations of individual studies, and demonstrated the implications of their outcome for daily practice and future research. If the authors received external funding, this was reported.

Some discrepancy between reviews was observed in terms of defining eHealth. For example, McLean et al [6] excluded Web-based tools and interventions for self-management in their Cochrane review on asthma patients because health care professionals were not actively involved with the ongoing delivery of the intervention. McGeary et al [22] chose a broader definition in their work on telehealth trials in pain management, including all studies that assessed a technology-based intervention extending care beyond the health care professional's office.

Many authors did not conduct a meta-analysis because of important differences perceived in study populations, interventions and outcome measures [18,19,25,26,31-33,35,36,39]. Instead, they performed a qualitative analysis of their findings. Several papers presented the results of a pooled analysis or subanalysis, despite substantial heterogeneity (ie, I^2 value >50) [6,12-14,23,24]. Three studies did not report heterogeneity [11,16,38].

Trend Analysis

Since the publication by Ekeland et al in 2010 [9], the number of systematic reviews and meta-analyses on eHealth interventions

in patients with somatic diseases has grown considerably (**Table 7**). In addition, 4 Cochrane reviews have recently been published [6,14,23,38]. However, a similar percentage of papers concluded that eHealth is effective/cost-effective or evidence is at least promising (65% vs 62%). Reviews focusing primarily on children or family caregivers still remain scarce. Between 2009 and 2012, home telemonitoring and video-teleconferencing were less frequently subject to a systematic review and/or meta-analysis on eHealth interventions, whereas educational tools and self-management programs were encountered more often. Data on economic outcome measures were less frequently reported in recent papers. Other study characteristics (eg, geographic area) barely differed between our review and the review by Ekeland et al [9].

Characteristic	Current review	Ekeland et al [9]
Country	100%	100%
Population	100%	100%
Intervention	100%	100%
Outcome measures	100%	100%
Study design	100%	100%
Year of publication	100%	100%
Number of studies	100%	100%
Number of participants	100%	100%
Number of outcomes	100%	100%
Number of interventions	100%	100%
Number of comparisons	100%	100%
Number of studies with economic outcome measures	100%	100%
Number of studies with educational tools	100%	100%
Number of studies with self-management programs	100%	100%
Number of studies with home telemonitoring	100%	100%
Number of studies with video-teleconferencing	100%	100%
Number of studies with children or family caregivers	100%	100%
Number of studies with Cochrane reviews	100%	100%
Number of studies with meta-analyses	100%	100%
Number of studies with systematic reviews	100%	100%
Number of studies with other types of reviews	100%	100%
Number of studies with no, limited, or inconsistent proof	100%	100%
Number of studies with effective/cost-effective or evidence is at least promising	100%	100%

□ Table 7. Trend analysis of differences in study characteristics of the current review compared with the review by Ekeland et al [9] published in 2010.

[View this table](#)

Discussion

The term eHealth can be defined briefly as the delivery of personalized health care at a distance through the use of technology. It is hypothesized that this field of medicine potentially enhances the quality of health care, with simultaneous reduction of health care costs. To support this hypothesis, we undertook a systematic review of systematic reviews and meta-analyses on the effectiveness/ cost-effectiveness of eHealth interventions in patients with somatic diseases. In addition, we performed a trend analysis to compare current findings with results from a systematic review by Ekeland et al published in 2010 [9].

In recent years, literature on eHealth has accumulated considerably. We found a total of 31 reviews, of which 20 (65%) concluded that eHealth interventions are effective/cost-effective or evidence is at least promising. Only 11 reviews (35%) showed no, limited, or inconsistent proof. These findings are consistent with the results from the review by Ekeland et al [9] (**Table 7**).

Furthermore, trend analysis shows reviews focusing primarily on children or family caregivers still remain scarce. Although a pooled (subgroup) analysis of aggregate data from randomized studies was performed in a higher percentage of more recently published reviews (45% vs 27%), data on economic outcome measures were less frequently reported (65% vs 85%).

Because our review is a systematic review of systematic reviews and meta-analyses, it holds 2 important limitations. Firstly, we relied on the adequate inclusion and critical appraisal of individual studies, as well as on a correct interpretation of study results by the authors of the reviews and meta-analyses included in the current review. We did not investigate whether reviews on similar topics comprised identical studies; neither did we examine possible discrepancies in the analyses of these individual studies when included in more than one review or meta-analysis. Noteworthy, systematic reviews of systematic reviews have been conducted before in other fields of medicine, including reconstructive surgery and neuroradiology [43,44].

Secondly, reviews differed substantially in terms of study populations, intervention components, comparison groups, and outcome measures, for example. Therefore, it is difficult to identify which patients are likely to benefit from which specific intervention. Home telemonitoring and structured telephone support seemed to be effective/cost-effective in patients with CHF (Table 3), whereas evidence on both interventions seemed limited or inconsistent in patients with chronic pulmonary diseases (Table 5). Meta-analysis was often impeded because of heterogeneity among individual studies. This may have demanded careful conclusions from the authors of that particular review. In several reviews, a pooled (subgroup) analyses was presented despite substantial heterogeneity among individual studies (ie, I^2 value >50) [17,42]. Publication bias may have been the result of the exclusion of small individual studies with negative results, which could have ultimately lead to overestimation of benefits [45,46]. Noteworthy, Ciere et al [32] proposed methodological weaknesses may be partially because of artifacts of poor reporting, rather than being a reflection of poor study design or implementation.

Regarding the aforementioned methodological shortcomings, Ekeland et al [47] performed a systematic review in which they summarize methodologies used in research on eHealth interventions, discuss knowledge gaps, and postulate

recommendations for methodological approaches for future research. Furthermore, we agree with recommendations made in previous reports to overcome the problem of between-study differences: researchers should adhere to and make transparent use of reporting guidelines appropriate for specific study designs. These guidelines may include Consolidated Standards of Reporting Trials (CONSORT)-EHEALTH for RCTs on eHealth interventions, Transparent Reporting of Evaluations with Nonrandomized Designs (TREND) and Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) for observational studies in general, and Workgroup for Intervention Development and Evaluation Research (WIDER) recommendations for the reporting of behavioral interventions [32,48-51].

Because pilot schemes are often limited to fewer than 100 patients, many researchers in the past decade have advocated larger RCTs with standardized study designs to provide definite proof on the effectiveness/cost-effectiveness of eHealth interventions. Results of the recent Whole System Demonstrator trial—involving 3230 patients with diabetes mellitus, COPD, and CHF—showed that eHealth interventions are associated with lower mortality and emergency admission rates [52]. In our opinion, these results should provide an important stimulus to invest in the incorporation of eHealth in daily practice. However, implementation difficulties, such as resistant or refractory behaviors of health care professionals, are an international phenomenon [53]. The Normalization Process Theory (NPT), a sociological theory that provides a framework for understanding the relationship between technology and the social environment, has been used to develop implementation tools such as the eHealth Implementation Toolkit (E-HIT) [54,55].

Although large, well-designed RCTs are likely to further support the evidence on the effectiveness/cost-effectiveness of eHealth initiatives, we believe it is more desirable to focus on overcoming the problematic gap between pilot schemes and daily practice. As proposed in both reviews by Ekeland et al [9,47], formative process assessments and complexity studies can be further explored to achieve this goal.

In conclusion, the number of reviews and meta-analyses on the effectiveness/cost-effectiveness of eHealth interventions in somatic diseases has increased considerably in recent years. The majority of these papers show eHealth is effective/cost-

effective, or at least suggests evidence is promising, which is consistent with previous findings. Data on economic outcome measures were less frequently reported in articles that were published more recently. This is an interesting finding, given the importance of formal cost analyses when considering implementation of eHealth interventions in daily practice. Although many researchers advocate larger, well-designed, controlled studies, we believe attention should be given to the development and evaluation of strategies to implement effective/cost-effective eHealth initiatives, rather than to further strengthen the evidence that has already been made available.

Acknowledgments

This paper received no sources of funding or sponsorship and there is no financial disclosure.

Conflicts of Interest

None declared.

References

1. Einthoven W. Het tele-cardiogram. *Ned Tijdschr Geneeskd* 1906;50:1517-1547.
2. Mariani AW, Pêgo-Fernandes PM. Telemedicine: a technological revolution. *Sao Paulo Med J* 2012;130(5):277-278 [[FREE Full text](#)] [[Medline](#)]
3. Rinde E, Balteskard L. Is there a future for telemedicine? *Lancet* 2002;359(9322):1957-1958. [[CrossRef](#)] [[Medline](#)]
4. American Telemedicine Association. What is telemedicine? □ URL: <http://www.americantelemed.org/about-telemedicine/what-is-telemedicine> [accessed 2014-03-31] [[WebCite Cache](#)]
5. Miller EA. Solving the disjuncture between research and practice: telehealth trends in the 21st century. *Health Policy* 2007;82(2):133-141. [[CrossRef](#)] [[Medline](#)]
6. McLean S, Chandler D, Nurmatov U, Liu J, Pagliari C, Car J, et al. Telehealthcare for asthma. *Cochrane Database Syst Rev* 2010(10):CD007717. [[CrossRef](#)] [[Medline](#)]
7. Finkelstein J, Friedman RH. Potential role of telecommunication technologies in the management of

- chronic health conditions. *Dis Manag Health Out* 2000;8(2):57-63. [[CrossRef](#)]
8. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and evaluating complex interventions: the new Medical Research Council guidance. *Int J Nurs Stud* 2013;50(5):587-592. [[CrossRef](#)] [[Medline](#)]
 9. Ekeland AG, Bowes A, Flottorp S. Effectiveness of telemedicine: a systematic review of reviews. *Int J Med Inform* 2010;79(11):736-771. [[CrossRef](#)] [[Medline](#)]
 10. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6(7):e1000097 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
 11. Agarwal R, Bills JE, Hecht TJ, Light RP. Role of home blood pressure monitoring in overcoming therapeutic inertia and improving hypertension control: a systematic review and meta-analysis. *Hypertension* 2011;57(1):29-38 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
 12. Angeles RN, Howard MI, Dolovich L. The effectiveness of web-based tools for improving blood glucose control in patients with diabetes mellitus: a meta-analysis. *Can J Diabetes* 2011;35(4):344-352. [[CrossRef](#)]
 13. Clarke M, Shah A, Sharma U. Systematic review of studies on telemonitoring of patients with congestive heart failure: a meta-analysis. *J Telemed Telecare* 2011;17(1):7-14. [[CrossRef](#)] [[Medline](#)]
 14. Inglis SC, Clark RA, McAlister FA, Ball J, Lewinter C, Cullington D, et al. Structured telephone support or telemonitoring programmes for patients with chronic heart failure. *Cochrane Database Syst Rev* 2010(8):CD007228. [[CrossRef](#)] [[Medline](#)]
 15. Klersy C, De Silvestri A, Gabutti G, Regoli F, Auricchio A. A meta-analysis of remote monitoring of heart failure patients. *J Am Coll Cardiol* 2009;54(18):1683-1694. [[CrossRef](#)] [[Medline](#)]
 16. Liang X, Wang Q, Yang X, Cao J, Chen J, Mo X, et al. Effect of mobile phone intervention for diabetes on glycaemic control: a meta-analysis. *Diabet Med* 2011;28(4):455-463. [[CrossRef](#)] [[Medline](#)]
 17. Polisena J, Tran K, Cimon K, Hutton B, McGill S, Palmer K, et al. Home telemonitoring for congestive heart failure: a

- systematic review and meta-analysis. *J Telemed Telecare* 2010;16(2):68-76. [[CrossRef](#)] [[Medline](#)]
18. Bender JL, Radhakrishnan A, Diorio C, Englesakis M, Jadad AR. Can pain be managed through the Internet? A systematic review of randomized controlled trials. *Pain* 2011;152(8):1740-1750. [[CrossRef](#)] [[Medline](#)]
 19. Eland-de Kok P, van Os-Medendorp H, Vergouwe-Meijer A, Bruijnzeel-Koomen C, Ros W. A systematic review of the effects of e-health on chronically ill patients. *J Clin Nurs* 2011;20(21-22):2997-3010. [[CrossRef](#)] [[Medline](#)]
 20. Hailey D, Roine R, Ohinmaa A, Dennett L. Evidence of benefit from telerehabilitation in routine care: a systematic review. *J Telemed Telecare* 2011;17(6):281-287. [[CrossRef](#)] [[Medline](#)]
 21. Johansson T, Wild C. Telerehabilitation in stroke care--a systematic review. *J Telemed Telecare* 2011;17(1):1-6. [[CrossRef](#)] [[Medline](#)]
 22. McGeary DD, McGeary CA, Gatchel RJ, Allison S, Hersh A. Assessment of research quality of telehealth trials in pain management: a meta-analysis. *Pain Pract* 2013;13(5):422-431. [[CrossRef](#)] [[Medline](#)]
 23. McLean S, Nurmatov U, Liu JL, Pagliari C, Car J, Sheikh A. Telehealthcare for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2011(7):CD007718. [[CrossRef](#)] [[Medline](#)]
 24. Omboni S, Guarda A. Impact of home blood pressure telemonitoring and blood pressure control: a meta-analysis of randomized controlled studies. *Am J Hypertens* 2011;24(9):989-998. [[CrossRef](#)] [[Medline](#)]
 25. Paré G, Moqadem K, Pineau G, St-Hilaire C. Clinical effects of home telemonitoring in the context of diabetes, asthma, heart failure and hypertension: a systematic review. *J Med Internet Res* 2010;12(2):e21 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
 26. Rietdijk R, Togher L, Power E. Supporting family members of people with traumatic brain injury using telehealth: a systematic review. *J Rehabil Med* 2012;44(11):913-921 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
 27. Samoocha D, Bruinvels DJ, Elbers NA, Anema JR, van der Beek AJ. Effectiveness of web-based interventions on patient empowerment: a systematic review and meta-

- analysis. *J Med Internet Res* 2010;12(2):e23 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
28. Stinson J, Wilson R, Gill N, Yamada J, Holt J. A systematic review of internet-based self-management interventions for youth with health conditions. *J Pediatr Psychol* 2009;34(5):495-510 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
29. van den Berg N, Schumann M, Kraft K, Hoffmann W. Telemedicine and telecare for older patients--a systematic review. *Maturitas* 2012;73(2):94-114. [[CrossRef](#)] [[Medline](#)]
30. Baron J, McBain H, Newman S. The impact of mobile monitoring technologies on glycosylated hemoglobin in diabetes: a systematic review. *J Diabetes Sci Technol* 2012;6(5):1185-1196 [[FREE Full text](#)] [[Medline](#)]
31. Bolton CE, Waters CS, Peirce S, Elwyn G, EPSRCMRC Grand Challenge Team. Insufficient evidence of benefit: a systematic review of home telemonitoring for COPD. *J Eval Clin Pract* 2011;17(6):1216-1222. [[CrossRef](#)] [[Medline](#)]
32. Ciere Y, Cartwright M, Newman SP. A systematic review of the mediating role of knowledge, self-efficacy and self-care behaviour in telehealth patients with heart failure. *J Telemed Telecare* 2012;18(7):384-391. [[CrossRef](#)] [[Medline](#)]
33. Franek J. Home telehealth for patients with chronic obstructive pulmonary disease (COPD): an evidence-based analysis. *Ont Health Technol Assess Ser* 2012;12(11):1-58 [[FREE Full text](#)] [[Medline](#)]
34. Mistry H. Systematic review of studies of the cost-effectiveness of telemedicine and telecare. Changes in the economic evidence over twenty years. *J Telemed Telecare* 2012;18(1):1-6. [[CrossRef](#)] [[Medline](#)]
35. Ryhänen AM, Siekkinen M, Rankinen S, Korvenranta H, Leino-Kilpi H. The effects of Internet or interactive computer-based patient education in the field of breast cancer: a systematic literature review. *Patient Educ Couns* 2010;79(1):5-13. [[CrossRef](#)] [[Medline](#)]
36. Saksena A. Computer-based education for patients with hypertension: a systematic review. *Health Educ J* 2010;69(3):236-245. [[CrossRef](#)]
37. Shulman RM, O'Gorman CS, Palmert MR. The impact of telemedicine interventions involving routine transmission of blood glucose data with clinician feedback on metabolic control in youth with type 1 diabetes: a systematic review

- and meta-analysis. *Int J Pediatr Endocrinol* 2010;2010:pii: 536957 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
38. Welsh EJ, Hasan M, Li P. Home-based educational interventions for children with asthma. *Cochrane Database Syst Rev* 2011(10):CD008469. [[CrossRef](#)] [[Medline](#)]
39. Wootton R. Twenty years of telemedicine in chronic disease management--an evidence synthesis. *J Telemed Telecare* 2012;18(4):211-220 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
40. Pron G, Ieraci L, Kaulback K, Medical Advisory Secretariat, Health Quality Ontario. Internet-based device-assisted remote monitoring of cardiovascular implantable electronic devices: an evidence-based analysis. *Ont Health Technol Assess Ser* 2012;12(1):1-86 [[FREE Full text](#)] [[Medline](#)]
41. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. Hillsdale, NJ: L Erlbaum Associates; 1988.
42. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21(11):1539-1558. [[CrossRef](#)] [[Medline](#)]
43. Thoma A, Veltri K, Haines T, Duku E. A systematic review of reviews comparing the effectiveness of endoscopic and open carpal tunnel decompression. *Plast Reconstr Surg* 2004;113(4):1184-1191. [[Medline](#)]
44. Clarke M. Systematic review of reviews of risk factors for intracranial aneurysms. *Neuroradiology* 2008;50(8):653-664. [[CrossRef](#)] [[Medline](#)]
45. Chan AW, Hróbjartsson A, Haahr MT, Gøtzsche PC, Altman DG. Empirical evidence for selective reporting of outcomes in randomized trials: comparison of protocols to published articles. *JAMA* 2004;291(20):2457-2465. [[CrossRef](#)] [[Medline](#)]
46. Eysenbach G. Tackling publication bias and selective reporting in health informatics research: register your eHealth trials in the International eHealth Studies Registry. *J Med Internet Res* 2004;6(3):e35 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
47. Ekeland AG, Bowes A, Flottorp S. Methodologies for assessing telemedicine: a systematic review of reviews. *Int J Med Inform* 2012;81(1):1-11. [[CrossRef](#)] [[Medline](#)]
48. Eysenbach G, CONSORT-EHEALTH Group. CONSORT-EHEALTH: improving and standardizing evaluation reports of web-based and mobile health interventions. *J Med Internet*

- Res 2011;13(4):e126 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
49. Des Jarlais DC, Lyles C, Crepaz N, TREND Group. Improving the reporting quality of nonrandomized evaluations of behavioral and public health interventions: the TREND statement. *Am J Public Health* 2004;94(3):361-366. [[Medline](#)]
50. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Bull World Health Organ* 2007;85(11):867-872 [[FREE Full text](#)] [[Medline](#)]
51. WIDER Group. Implementation Science. Recommendations to improve reporting of the content of behaviour change interventions □ URL: <http://www.implementationscience.com/content/supplementary/1748-5908-7-70-s4.pdf> [accessed 2014-03-31] [[WebCite Cache](#)]
52. Steventon A, Bardsley M, Billings J, Dixon J, Doll H, Hirani S, Whole System Demonstrator Evaluation Team. Effect of telehealth on use of secondary care and mortality: findings from the Whole System Demonstrator cluster randomised trial. *BMJ* 2012;344:e3874 [[FREE Full text](#)] [[Medline](#)]
53. Murray E, Burns J, May C, Finch T, O'Donnell C, Wallace P, et al. Why is it difficult to implement e-health initiatives? A qualitative study. *Implement Sci* 2011;6:6 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
54. May C, Finch T. Implementing, embedding, and integrating practices: an outline of Normalization Process Theory. *Sociology* 2009;43(3):535-554. [[CrossRef](#)]
55. MacFarlane A, Clerkin P, Murray E, Heaney DJ, Wakeling M, Pesola UM, et al. The e-Health Implementation Toolkit: qualitative evaluation across four European countries. *Implement Sci* 2011;6:122 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
-

Abbreviations

CHF: congestive heart failure

CONSORT: Consolidated Standards of Reporting Trials

COPD: chronic obstructive pulmonary disease

E-HIT: eHealth Implementation Toolkit

HbA1c: glycosylated hemoglobin

ICU: intensive care unit

LDL: low-density lipoprotein

NPT: Normalization Process Theory

PRISMA: Preferred Reporting Items for Systematic
Reviews and Meta-Analyses

QUOROM: Quality of Reporting of Meta-analyses

RCT: randomized controlled trial

STROBE: Strengthening the Reporting of Observational
Studies in Epidemiology

TREND: Transparent Reporting of Evaluations with
Nonrandomized Designs

WIDER: Workgroup for Intervention Development and
Evaluation Research

Edited by G Eysenbach; submitted 24.06.13; peer-reviewed by A Straten, R Choubisa; comments to author 12.07.13; revised version received 13.02.14; accepted 13.03.14; published 16.04.14

Copyright

©Niels J Elbert, Harmieke van Os-Medendorp, Wilco van Renselaar, Anne G Ekeland, Leona Hakkaart-van Roijen, Hein Raat, Tamar EC Nijsten, Suzanne GMA Pasmans. Originally published in the Journal of Medical Internet Research (<http://www.jmir.org>), 16.04.2014.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research, is properly cited. The complete

bibliographic information, a link to the original publication on <http://www.jmir.org/>, as well as this copyright and license information must be included.

Journals

[JMIR Research Protocols](#)
[JMIR mHealth and uHealth](#)
[interactive Journal of Medical Research](#)
[JMIR Medical Informatics](#)
[Medicine 2.0](#)
[JMIR Serious Games](#)
[JMIR Mental Health](#)
[JMIR Human Factors](#)
[JMIR Rehabilitation and Assistive Technologies](#)
[JMIR Cancer](#)

Editorial Policies

[Focus and Scope](#)
[Section Policies](#)
[Peer Review Process](#)
[Publication Frequency](#)
[Memberships](#)
[Author Self-Archiving](#)
[Why you should choose JMIR to publish your research - advantages of the journal](#)
[Indexing and Impact Factor](#)
[Editorial Board Policy](#)
[Record Keeping and Research Policy](#)
[Subscriptions \(Membership\)](#)
[Theme Issues and Guest Editors](#)
[Trademarks and Service Marks Policy](#)
[Subscribe to RSS Feeds / add JMIR headlines to your homepage or blog](#)

[Fee Schedule](#)

[Instructions for authors](#)

[About the publisher](#)

[New Journal / Editor-in-Chief Proposals](#)

Submissions

[Online Submissions](#)

[Author Guidelines](#)

[Copyright Notice](#)

[Privacy Statement](#)


Other

[Announcements](#)

[Editorial Team](#)

[Feedback](#)

[Contact Us](#)



Copyright © 2015 JMIR Publications Inc.