

Results from the Tromsø Intervention Study on Preterms until children's age of nine.

The influence of structured early parental guidance on behavior-emotional development and well-being among children born preterm and parenting stress in their families.

Inger Pauline Landsem

A dissertation for the degree of Philosophiae Doctor – Month 20xx

Table of Contents

Acknowledgements	3
List of Papers	5
Summary in English	6
Summary in Norwegian	9
Definitions and abbreviations	11
1. Introduction	13
1.1. Neonatal care in health care systems	13
1.2. Health and developmental risks in children born prematurely	14
1.3. Development of behavioral problems among children born preterm.....	15
1.3.1. Internalizing problems	17
1.3.2. Externalizing problems	18
1.3.3. Attention problems	19
1.3.4. Social problems	21
1.3.5. Thought problems	23
1.3.6. Mental health among children born preterm.....	23
1.4. Quality of life (QoL) of children born preterm	24
1.5. Development of competencies until middle childhood	25
1.6. Parental adaptation to the caretaking of a child born preterm	27
1.6.1. Enjoyment of and becoming attached to the infant.....	27
1.6.2. Parenting stress and impact on parenting behavior	28
1.6.3. Contribution of the child to the parent-child interaction	30
1.7. Early interventions addressing developmental difficulties	31
1.8. A transactional model of development	33
1.9. Mechanisms involved in developmental change	34
2. Aims and questions	37

3. Materials and Methods	38
3.1. Participants	38
3.2. Study implementation	39
3.3. The intervention program.....	40
3.4. Measurements	41
3.5. Follow-up procedures	45
3.6. Statistical analysis	46
4. Summary of results	48
5. Discussion	53
5.1. Children’s emotional-behavioral development	54
5.2. Family climate and parental agreement	61
5.3. Socio-emotional problems, social competence and mental health	64
5.4. Attention problems and perceived academic performance	66
5.5. Parenting stress and possible influences on children’s behavioral problems	69
5.6. Influences on children’s quality of life	74
6. Strengths and limitations	76
7. Clinical Implications /implications for later research	77
8. Conclusions	82
References	83
Paper I	
Paper II	
Paper III	
Appendix	

Acknowledgements

I have been following the planning, implementation and progression of the Tromsø Intervention Study on Preterms (TISP) for 17 years. The study, and foremost the knowledge that motivated the study has influenced both my professional and personal life significantly.

First of all, I want to thank my supervisors Professor John Andreas Rønning and Senior Lecturer Bjørn Helge Handegård at the UiT, The Arctic University of Norway.

John was the one that made it possible for me to apply for a research education related to this study. As the director of the study, and a very well informed first supervisor in the field of early child development his knowledge, guidance and availability has been of vital importance. Similarly, without the kind, patient and matter-of-factly supervision and corrections from Bjørn Helge especially related to statistical analysis and the presentation of results, this work would not been completed. I am deeply thankful to you both.

I will also thank Stein Erik Ulvund and Per Ivar Kaaresen who, together with Rønning and Dr. Lauritz B. Dahl designed the study, and contributed with well-informed suggestions to the research articles. Elin Gullhav has, as the leader of the Child and Adolescent Department in our hospital, been enthusiastic and helpful. She obtained a position and a nice work-place for me in the department. In the “pink-room” we have had a warm atmosphere and I have got to know nice persons as Marit, Christian and Catharina among others. Thanks a lot!

A special thanks goes to Jorunn Tunby, the study coordinator in TISP since the beginning and the one I have shared talks, discussions and lunch-breaks with across these three years. I will also mention my dear colleagues who, together with Jorunn and me implemented the interventions; Randi Berg Høier, Nina B. Cheetham, Kirsti Vikhammer, Unni Byrknes, Anne Helene Uhre and Hilde Sletvold. We shared an exciting awakening confronted with new knowledge about early child development and you all taught me a lot. Thank you.

The most important insights were however learned through the meetings with the participating children and their parents during the intervention sessions, and through all the other families I have met since I entered the neonatal intensive care unit for the first time. Studies may have made me

more skilled and well-informed but the meetings with you have nurtured the wisdom I hope to have obtained.

Last, but not least, I would like to thank family and friends who have believed in and supported me - even some of you may have thought "when will she stop studying?...".

My very good-hearted husband has been with me every day and our sons have grown up as two clever adults. I've been introduced to lovely girl-friends and even the most wonderful of all – become a grateful and proud grandmother to Mynthe Annie, the new star in my life.

Tromsø, 12.10.15

Inger Pauline Landsem

List of Papers

Paper I

Landsem IP, Handegård BH, Ulvund SE, Tunby J, Kaaresen PI, Rønning JA. Does an early intervention influence behavioral development until age nine in children born prematurely? *Child Development* 2015; 86(4):1063-1079.

Paper II

Landsem IP, Handegård BH, Tunby J, Ulvund SE, Rønning JA. Early intervention program reduces stress in parents of preterms during childhood, a randomized controlled trial. *Trials* 2014; 15: 387.

Paper III

Landsem IP, Handegård BH, Ulvund SE, Kaaresen PI, Rønning PI. Early intervention influences positively Quality of Life as reported by prematurely born children at age nine and their parents; a randomized clinical trial. *Health and Quality of Life Outcomes* 2015; 13:25.

Summary in English

Three studies emanating from “The Tromsø Intervention Study on Preterms” (TISP) are summarized in the thesis. The main focus is whether an early structured intervention program, given to families with preterm infants in the newborn period may influence; children's socio-emotional behavioral development up to 9 years of age, mothers and fathers reports of parenting stress in the same period of time and children's quality of life by nine years.

TISP started in 1999 and the inclusion of children with families lasted until 2002. Infants born preterm from Troms or Finnmark (birth weight below 2000 gram) were recruited and randomized to a premature intervention group (PI = 72 children) and a premature control group (PC = 74 children). 75 healthy term newborns from the city of Tromsø were recruited to form a term control group (TR). Children of mothers who did not speak Norwegian, which were triplets or was proven with severe neuro-developmental disorders were excluded from the study.

The intervention, a modified version of The Mother-Infant Transaction Program (MITP-M), included 8 hours of supervision for the mother (and father if present) last week of the child's stay in hospital and four home visits approximately 7, 14, 30 and 90 days after discharge. This was given to the PI group while parents in the PC- and the TR group received information according to the hospitals guidelines. The aim of MITP-M was to promote parents' enthusiasm for their children and help them recognize signs of children's organization versus need for regulatory support. Furthermore, to show parents appropriate interactions with the child that would minimize child disturbance and increase interactional satisfaction for all partners.

All participating families are followed up through developmental tests of the children and data collections using questionnaires when the children were 6 months, 1, 2, 3, 5, 7, 9 years.

Dropout rates have been low. 129 preterm infants (88%) were still participating at children's age of nine. The main focus was to analyze whether differences between the PI and the PC group emerged as a possible consequence of the intervention. The PI and the PC group were in addition compared with the results of the TR group.

Children's behavioral development is analyzed using mothers, fathers and teachers reports of behavioral problems at 2, 3, 5, 7 and 9 years of age. Teachers reported at 7 and 9 years. They had no information about children's group belonging. Longitudinal analyzes showed that the average prevalence of behavior problems varied according to the same pattern in the three groups from 2 to 9 years. PC mothers and PC fathers reported more behavior problems than PI parents at all follow-ups, but this constituted non-significant differences between PI and PC group with regard to internalizing and externalizing behavior from 2 to 9 years. No longitudinal differences were detected in longitudinal behavior development between the PI and the TR group but one was detected between the PC and the TR group. Fathers in PC group reported a steeper increase in symptoms of anxiety in children from their age of 5 until 9 years compared with fathers in the TR group.

Significant differences between the PI and the PC group became visible at children's age of 7 and 9 years. Both parents and teachers reported significantly less attentional problems and better social competence, adaptability and school-related performances in the PI group. PI children were in average not reported as different from the TR group at their age of nine, both what came to school adjustment, achievements and the occurrence of problems. Significant differences between the PC and the TR group persisted concerning the extent of behavior problems and competencies.

The second study reported mothers and fathers reports of child- and parent-related stress at all follow-ups from children's age of 6 months until 9 years. PI mothers reported significantly less stress than PC mothers at all times and the same was evident for fathers at children's age of 2, 3 and 5 years. PC mothers reported high and stable levels of child-related stress across pre-school years, especially related to statements concerning children's adaptability and mood. In contrast, PI- and TR-mothers reported decreasing levels of parenting stress from children's age of one. Differences between PI and PC group in terms of parents-related stress referred to less perception of parental attachment and competence in the PC group. The last study in this thesis analyzed children's and parent's reports of children's quality of life at their age of nine. Children in PI group reported higher physical wellbeing than the PC children while PI parents report higher emotional and school-related well-being than parents in PC group.

The research presented in this thesis indicate that the structured guidance given parents of preterms in the newborn period has promoted positive and long-lasting effects on the PI group as these children perform at similar level as their term born peers at 9 years of age.

Summary in Norwegian

Avhandlingen er en sammenfatning av tre studier som utgår fra The Tromsø Intervention Study on Preterms (TISP), tidligere omtalt som "Tidlig Intervensjon 2000". Hovedfokus i studiene er hvorvidt et tidlig strukturert veiledningsprogram, gitt til familier med prematurt fødte barn i nyfødtp perioden kunne påvirke; barnas sosio-emosjonelle adferdsutvikling opp til 9 års alder, mødre og fedres rapport av foreldrestress i samme periode og barnas livskvalitet ved 9 år.

TISP startet i 1999 og inkluderingen av barn pågikk fram til 2002. Prematurt født barn fra Troms eller Finnmark, med fødselsvekt under 2000 gram ble rekruttert og randomisert til en prematur intervensjons gruppe (PI =72 barn) og en prematur kontroll gruppe (PC = 74 barn). I tillegg ble 75 friske, fullbårne nyfødte hjemmehørende i Tromsø rekruttert til en termin kontroll gruppe (TR). Barn av mødre som ikke snakket norsk, som var trillinger eller ble påvist med alvorlige nevro-utviklingsmessige tilstander ble utelukket fra studien.

Veiledningsprogrammet, en modifisert versjon av The Mother-Infant Transaction-Program (MITP-M), omfattet 8 veiledningstimer for mor (og far hvis til stede) siste uka av barnets opphold i sykehuset og fire hjemmebesøk ved cirka 7, 14, 30 og 90 dager etter utskrivelse. Dette ble gitt til PI gruppen mens barn og foreldre i PC og TR gruppene mottok informasjon i henhold til sykehusets retningslinjer. Siktemålet med MITP-M var å fremme foreldrenes begeistring for sitt barn og hjelpe dem å gjenkjenne barnets tegn på likevekt versus behov for reguleringsstøtte. Videre å lære dem hensiktsmessige måter å samhandle med barnet slik at barnet ble minst mulig forstyrret og samhandlingen mest mulig tilfredsstillende både for barn og foreldre.

Alle deltagende familier er fulgt opp med utviklingsmessige tester av barna og data innsamling ved hjelp av spørreskjema når barna var 6 mnd, 1, 2, 3, 5, 7, 9 år. Studien har hatt lite frafall av deltakere. Ved 9 år møtte 129 prematurt fødte barn (88 %). Hovedfokus var å analysere hvorvidt det framkom forskjeller mellom gruppene med prematurt fødte barn (PI & PC). I tillegg er PI og PC gruppene hver for seg sammenlignet med resultatene i termin kontrollgruppen.

Barnas adferdsutvikling er analysert ved hjelp av mødre-, fedre- og lærer rapporterte adferdsproblemer ved 2, 3, 5, 7 og 9 års alder. Lærere rapporterte ved 7 og 9 år. Disse hadde ikke informasjon om hvilken studiegruppe barnet tilhørte. Longitudinelle analyser viste at

gjennomsnittlig forekomst av adferdsproblemer varierte etter samme mønster i de tre gruppene fra 2 til 9 år. PC mødre og PC fedre rapporterte mer problemer enn PI foreldre på alle tidspunkt, men dette utgjorde ikke-signifikante forskjeller mellom PI og PC gruppen med hensyn til utagerende- og tilbaketrekkings adferd fra 2 til 9 år når. I sammenligninger med TR gruppen framkom ingen forskjeller mellom PI- og TR gruppen, mens en forskjell mellom PC- og TR gruppen var signifikant. Fedre i PC gruppen rapporterte en brattere økning av symptomer på engstelighet hos barna fra 5 år og opp til 9 års alder enn fedrene i TR gruppen.

Signifikante forskjeller mellom PI og PC gruppen ble synlig ved 7 og 9 år. Både PI foreldre og PI lærere rapporterte signifikant mindre oppmerksomhetsproblemer samt noe bedre sosiale ferdigheter, tilpasningsevne og skoleferdigheter blant barna enn rapportert i PC gruppen. Ved 9 år ble PI gruppen rapportert på samme nivå som TR gruppen både hva gjaldt skoletilpasning, prestasjoner og forekomst av problemer, mens signifikante forskjeller mellom PC- og TR gruppen vedvarte.

I den andre studien analyserte vi mødres og fedres rapport av barne- og foreldre relatert stress på alle oppfølgingstidspunkt fra barna var 6 måneder og opp til 9 år. PI-mødre rapporterte signifikant mindre stress enn PC-mødre på alle tidspunkt og det samme var tilfelle for PI- versus PC fedre ved 2, 3 og 5 år. PC- mødre rapporterte stabilt høyt barne-relatert stress opp til 5 års alder, spesielt relatert til utsagn som fokuserte på tilpasningsvansker og lunefullt humør hos barnet. PI- og TR-mødre rapporterte på sin side avtagende barne-relatert stress fra 1 års alder. Forskjellene mellom PI og PC gruppen hva gjaldt foreldre-relatert stress omhandlet spesielt rapport av mindre positive tilknytningsfølelser og foreldre kompetanse i PC gruppen.

Den siste studien i avhandlingen rapporterer noen forskjeller mellom prematurgruppene som berører barnas opplevde livskvalitet. Barn i PI gruppen rapporterer høyere kroppslig velvære ved 9 år enn PC barna mens PI foreldre rapporterer høyere følelsesmessig og skolerelatert velvære enn foreldre i PC gruppen.

Studiene i denne avhandlingen indikerer at veiledningsprogrammet som ble gitt PI gruppen har gitt langvarige, positive effekter i familiene og at PI barna fungerer på nivå med sine jevnaldrende født til termin ved 9 års alder.

Definitions and abbreviations

MITP	Mother Infant Transaction Program
MITP-M	Mother Infant Transaction Program, modified version in TISP
NBAS	Neonatal Behavior Assessment Scale
NICU	Neonatal Intensive Care Unit
PSE	Parenting Self-Efficacy
SES	Socio-Economic Status (parental years of education, employment, marital status, income, living conditions, among others)
TISP	Tromsø Intervention Study on Preterms
TISP, study groups:	
PC	Preterm Control group
PI	Preterm Intervention group
TR	Term Control group

Birth & Medical terms

BW	Birth Weight
BW groups:	
LBW	Low Birth Weight (BW < 2500 g)
VLBW	Very Low Birth Weight (BW < 1500 g)
ELBW	Extremely Low Birth Weight (BW < 1000 g)
BPD	Bronchopulmonary Dysplasia
CRIB	Clinical Risk Index for Babies (a tool for assessing initial neonatal medical risk)
GA	Gestation Age (the number of weeks that a baby has been in the uterus. Newborns delivered before 37 GA are considered premature)
KC	Kangaroo Care (the newborn is held with skin-to-skin contact with an adult)

Measurement & outcome subscales

ASEBA	The Achenbach System of Empirically Based Assessment
ASEBA Questionnaires:	
CBCL	Child Behavior Checklist (questionnaire)
TRF	Teachers Report Form (questionnaire)
KINDL	The Kinder Lebensqualität Fragebogen (questionnaire)

PSI Parenting Stress Index, full form:

PSI domains:

CD Child Domain

PD Parent Domain

TS Total Stress

PSI-sf Parenting Stress Index, short form;

PSI-SF domains:

DC Difficult Child

PS Parenting Stress

P-CDI Parent-Child Difficult Interaction

QoL Quality of life

SDQ Strength and Difficulties Questionnaire

Statistical terms

ES Effect Size

GLMM Generalized Linear Mixed Models

ICC Intraclass Correlation

LMM Linear Mixed Models

OR Odds Ratio

Introduction

1.1 Neonatal care in health care systems

The development of neonatal care over the last 60 years has been described as a movement from professional- and institution-based autocracy to a parent-professional partnership for the care of hospitalized children [1,2]. In the earliest years, preterm born children were still cared for as individuals and institutionalized independently of their parent's wishes. Bowlby had questioned the prevailing assumption that a child develops independently of the environment in 1952 [3]. Maternal roles were emphasized as important, and research documented the negative effects on child development of the practice of parent-child separation. [4-7]. The impact of parental involvement on the health and development of children has been noted [8]. This knowledge was gradually incorporated in the care of hospitalized children in general and for children born prematurely in particular [3,8-11].

The development of parental involvement has been described as a hierarchy, beginning with the access of parents to the hospital environment (1960s) and proceeding to participation (1980s), partnership and later family-centered care [12]. These processes involved considerable changes in parental roles and were requested by parents and dedicated health care providers and were also supported by changes in juridical laws and regulations [7,13,14].

Family-centered care aims to empower parents. In terms of care of families with prematurely born children, this implicates the transmission of interdisciplinary knowledge, confidence and self-efficacy [2,15-18]. Parents typically provide the first and most proximal environment in their children's life. For prematurely born infants, this natural protection is disturbed; both because parent-child skin-to-skin contact may be hindered by medical equipment/treatment, parents may have limited access to the neonatal intensive care unit (NICU) and the child may be cared for by many different people.

The multidisciplinary care of children born preterm has experienced significant improvements over the last few decades (antenatal steroids, postnatal surfactant therapy, specialized technology and equipment, specialized personnel and increased parental involvement)[1,19-21]. However, although

the survival rates for preterms have increased, morbidity has also risen [22,23]. In particular, the long-term outcomes related to socio-emotional, attentional and academic competencies among preterm born children continue to lag behind term-born peers [19,24,25].

1.2 Health and developmental risks of children born prematurely

The level of risk associated with preterm birth is inversely related to the length of the pregnancy [19,25-27]. In addition, the developmental impact of preterm birth cannot be described without relating outcomes in different countries and areas with more or less developed health systems. Globally, the rates of preterm birth vary between 5 and 18%. Norway is positioned among the countries with the best outcomes because of the relatively low prevalence of preterm birth (4.9% in 2012) [28], high surveillance rates and relatively low morbidity in comparison to less developed countries. The differences between countries may be exemplified by reports from the World Health Organization (WHO) describing a 50% chance of surveillance at 24 gestational weeks (w of GA) with access to a NICU in a high income country, while a surveillance rate of 50% at 34 w of GA has been reported in several low and middle income countries [29]. Internationally, the prevalence of preterm birth has been reported to be increasing. A low birth weight (LBW) is reported in approximately 15.5% of newborns in developing countries, and in the USA, the occurrence of preterm birth has increased with 31% over the last 35 years [30]. However, the prevalence of LBW newborns (BW < 2500 grams) in Norway has stabilized over the last decade. Approximately 3000 newborns (4.9%) were born before 37 w of GA in 2012, 2% had a BW < 2000 grams and 287 had a BW < 1000 grams. [28].

Children born with the lowest birth weights (500 – 1500 grams) contribute greatly to the rates of perinatal mortality and morbidity despite accounting for less than 2% of all deliveries [19]. Several biological and environmental factors contribute to individual differences in the risk profiles of children. First, the severity of the neonatal course (need for treatment interventions, complications, experiences of pain, length of stay in hospital), followed by the quality of resources in the caring environment (e.g., socio-economic status, support, parental mental health) and, finally, the potential sequelae they experience after preterm birth (e.g., lung disease, brain injury, neurological complications, hearing loss and blindness). The major handicaps mentioned above are reported in

5% of the general population and in 6-8% of low birth weight (LBW) children, 14-17% of very low birth weight (VLBW) children and 20-25% of extremely low birth weight (ELBW) children [19,25]. Moderate to late preterm children (32-36^{±5} w of GA) have been reported to have fewer major handicaps, but they may experience several problems during childhood. After controlling for confounders such as maternal race, prenatal steroids, infant gender and chorioamnionitis, a 23% decrease in adverse outcomes per week longer of pregnancy was reported for children who were born between 32 and 39 w of GA [30]. A high prevalence (50 – 70%) of dysfunction but with a low severity has been reported to occur in very low birth weight (VLBW) children. These problems are often identified as attention-related and/or social problems and, to some degree, to more internalizing behavior [24,26]. Children born preterm are often reported to exhibit under-achievement after entering primary school and more frequently require specialized school assistance [19,31]. Different features of these problems will be reviewed later in this thesis. A recurring problem is the absence of good predictors of the frequently reported, subtle problems that occur among children born preterm [15,19].

1.3 Development of behavioral problems in children born preterm

Unusual or abnormal behavior among children born preterm was reported long before neonatal intensive care medicine was defined. Prior to the Second World War, a syndrome characterized by behavioral difficulties (hyperactivity, susceptibility to distraction, unpredictability and shyness) was described [33]. In the 1970s, a description of transient dystonia was reported, and parents described problems such as irritability, crying and feeding difficulties [34]. The developmental outcomes of preterm children were subsequently extensively studied, but the behavior development of these children has been investigated very little compared with that of short and long-term cognitive growth [35]. Nevertheless, a greater number of behavioral problems and a higher incidence of psychological disorders have been reported in children born preterm compared with those born full-term [35]. A behavioral problem prevalence of 20% (twice the value determined for children with typical development) was reported in the Infant Health and Developmental Program (IHDP) for children aged 3, 5 and 8 years [35]. Similar results have been reported in other studies [26,27], while a Norwegian follow-up study reported that 40% of preterms (BW < 2000 grams) exhibited greater behavioral difficulties [36]. The IHDP study identified predictors of an increased risk of problems

such as maternal psychological stress at 40 w of GA, younger maternal age and maternal smoking. In addition, child- and birth related-factors such as gender, birth weight, gestational age and neonatal health were marginally predictive of behavioral problems [35]. Before specific areas of behavioral problems are described, some features of the children's general behavioral development, as expected across early and middle childhood, are mentioned.

The development of children comprises several dimensions and developmental milestones [37]. Developmental changes are described as normative, nonreversible, relatively stable, frequently sequential and, especially in children born preterm, associated with maturation [38]. The maturation and growth of children is especially apparent in the acquisition of new skills and the understanding and expression of language. Some milestones are included in regular health care assessments throughout childhood, e.g., motor and language achievements. Other milestones exist as more or less defined norms of the society impacting the developing child. To some degree, every child will be viewed in a mirror that is shaped by the expectations and pre-understanding of the norms that are integrated in others and in society. Five dimensions with developmental milestones concerning mental maturation and development have been described as important [37].

Social competence is mentioned first, and it may be defined as the effectiveness in developmentally appropriate social interactions. The main skills identifying such competence are cooperation, helpfulness and the ability to resolve conflicts [39]. During the newborn period, a type of social competence may be viewed as the ability of the child to elicit responses and positive responses from their caregivers [40,41]. The second dimension is attachment, which may be described as the deep, selective and enduring connection between a child and the caregiver that enables the child to form positive, close relationships with parents, peers and, later, partners. The third dimension, emotional competence may be defined as the multifaceted ability to be aware of one's own and other's emotions and to utilize this awareness in interpersonal interactions and in the regulation of emotional experiences. The fourth dimension is the multidimensional construct of self-perceived competence. This dimension considers the child's evaluation of his or her own abilities (cognitive, physical, social), especially in comparison to others. Finally, Denham [37] refers to a dimension called temperament and personality. These features are considered to be fundamental to how children function in social and familial relationships. Temperament is defined as an individual style

concerning reactivity and self-regulation in which emotional reactivity refers to the speed and intensity with which individuals respond to events.

Throughout childhood, all of these features that were briefly mentioned above develop within each individual child and are influenced by biological and environmental conditions [1]. The ease or success of these developments may be reflected in the behavior of the child, which is characterized either by successful adaptation and health or maladaptation and disorder [1]. The socio-emotional functioning of children during childhood is very important because it serves as a predictor and is associated with later behavioral problems, mental health and successful functioning in school [37,43].

Behavioral problems refer to dimensionally measured behavior, including both normal and atypical ranges of behavior [44]. Children born preterm display behavior that is similar to other developing children. Normative descriptions of expected problems in populations at different ages are relatively new, especially those described for children who have not yet reached middle childhood [43,45]. There is suggestive evidence concerning age-related changes in problem behavior, and differences between genders have been described [46]. Behavioral problems are frequently categorized according to a two-dimensional taxonomy that separates problems related to mood, anxiousness or depression as internalizing problems and out-acting/intrusive behaviors as externalizing problems [46-50]. In addition, problems related to the adaptation of children to their social surroundings have frequently been described as either attention-related or social-emotional problems [45,47]. Different types of problems are detailed in the following sections.

1.3.1 Internalizing problems

Internalizing problems are often described as mood and anxiety problems, including symptoms of depression, somatic complaints and withdrawal behavior [45-50]. Each feature may be related to different developmental difficulties at different ages and expressed in different ways across childhood [43], and a greater number of problems have a large influence on the social development of children [46]. In a large, longitudinal study of internalizing behavior in children aged two to eleven years old, different trajectories were described across childhood. Two-thirds of the children were reported to have few problems during childhood, while one-third exhibited problems with

decreasing, increasing or sustained elevated trajectories [43]. Interestingly, a greater number of maternal psychological symptoms during infancy predicted increasing, decreasing or elevated trajectories, while good maternal mental health predicted a stable, low trajectory. Toddler twin studies suggest that genetic factors account for 50% of the variance and a shared environment for 30% of the variance in internalizing symptoms [46]. Internalizing behavior has been less well studied and reported compared with externalizing behavior [46], and it is not as easy to observe in a child's behavior [45]. The detection and interpretation of internalizing problems are more dependent on the person reporting the problems (e.g., mother, father or teacher) and the age of the child [51]. In infancy, distress and fearful behavior related to separation from caregivers may be viewed as developmentally appropriate behavior, whereas it may be seen as separation anxiety in later childhood or a type of social phobia among adolescences [43,46]. Children born preterm have frequently been reported to have increased levels of internalizing behavior [24,51-56]. The etiology of this behavior pattern has been unclear and will be discussed later. In questionnaires that were answered by parents or teachers, this type of problem is typically expressed as the child as follows: acting too young, refusing to participate in activities, seems shy and with little expression in response to positive stimuli [48-50].

Agreement between different observers regarding this type of behavior is generally lower than that in reports on externalizing behavior [51,58-60]. In addition, internalizing behavior varies because many types of behavior differ between countries, especially with respect to socio-economic factors (SES) [61] and cultural differences [62]. A Norwegian population survey reported differences in parental reports of internalizing problems among 8-10-year-old children in Norway and Britain [62]. Norwegian parents seemed to under-report internalizing problems compared with British parents. This observation could depend on a higher "normalizing" view of emotional difficulties among Norwegian adults or, possibly, an under-detection of emotional difficulties in young children.

1.3.2 Externalizing problems

Externalizing behaviors are actions that are directed out towards others. According to the Achenbach System of Empirically Based Assessments (ASEBA), externalizing behavior consists of aggressive and delinquent behavior [47]. In questionnaires, aggressive or destructive behavior

among toddlers are described as, e.g., being jealous, screaming, destroying other things or eating non-food. Later in childhood, this behavior may be described as, e.g., screaming, fighting, teasing, bragging, talking too much or appearing irritable [49,50]. Signs of this type of behavior vary from typically non-appropriate and immature behavior to serious dysfunctional behavior. Thus, most children will to some degree display aggressive or oppositional behavior, especially during early childhood. Delinquent behavior is described as more serious, dysregulated and aversive behavior such as, e.g., stealing, running away from home, skipping school, and swearing [49,50]. Children who display elevated levels of externalizing behavior may be described as possessing underdeveloped self-regulatory abilities that may lead to uninhibited behavior and the expression of poor self-control [63]. Factors such as ethnicity [64,65], SES [64,66] and gender [64,66] are also associated with the amount of externalizing behavior [65]. A greater amount of externalizing behavior in toddlerhood has been reported to be a strong predictor of subsequent adaptation difficulties in early school years [67]; however, most children have decreasing trajectories of externalizing behavior after toddlerhood [67,68].

Normative trajectories of externalizing behavior have been described to decrease during childhood [45], and in contrast to internalizing behaviors, they are not described as elevated among children born preterm compared with full-terms [24]. An even earlier meta-analysis has described elevated levels of externalizing behavior in very preterm compared with full-term children [26]. The results of subsequent reports did not support this finding [24,56] even though a lower BW was found to be associated with more externalizing behavioral problems.

1.3.3. Attention problems

Attention is the ability of a child to orient to, shift between and focus on something in the external world [69]. Attention problems are often described as impulsivity, hyperactivity and inattention [48-50]. Questions concerning attention problems concern the ability of the child to act at an age-appropriate level, concentrate, and sit still and whether behavior as clumsiness and staring are observed [ibid]. While problems such as impulsivity and hyperactivity have been reported to decline during middle childhood, inattention appears to be more stable across the age groups [70]. Boys have been described to have more attention problems than girls [45,71]. Attentional

competence is a basic premise for interactions and contact with others, and it may be considered a potential mechanism for the development of later and/or other socio-emotional problems [69].

Children born preterm are more often identified with early attention problems than those born at term [72]. Teacher and parent ratings of problems have been reported with standard deviations (SDs) that are 0.43 and 0.59 higher than those reported for full-term peers [24]. A neuro-psychological model has been useful to briefly describe some of the elements of the highly complex processes involved in the development of attention [69,73]. Three networks in the brain have been described to be involved and interconnected:¹⁾ the orienting system, ²⁾ the arousal system and an ³⁾ executive attention system [73]. First, an orienting system is visible already in newborns and is fully functional during the first 6 months of life [74]. Newborns gradually become capable of orienting themselves to movements or objects and of disengaging or shifting their direction. The orientation qualities have been explored by assessing the duration of the gaze, changes in attentional focus and the ability of the infant to successfully disengage from stimuli. Second, the alerting or arousal system is related to the capacity of the child to maintain a state of alert arousal that enables him or her to successfully process information [69]. During infancy and early childhood, this development has been assessed by observing sequences of sustained (focused) attention in children during free play. Third, when infants reach toddlerhood, the executive control system matures. In this phase, attention becomes more related to planned and child-generated activities with objects. During further development, the attentional competence of children becomes more directly assessable by caretakers because the caretakers may observe the degree to which the child can pay attention to a task until it is successfully solved. A link between early focused attention and later cognitive outcomes has been suggested for children born preterm; attention at 7 months has been shown to be predictive of reported behavioral problems and cognitive abilities during the preschool years [75]. In their longitudinal study of normative behavior, Bongers et al. claimed that observable attention problems are especially apparent when children attend school [45].

Children born preterm have previously been reported to have greater problems in all attentional systems.¹⁾ They show less efficient orientating attention during the first 6 months of life [69,74],²⁾ and they shift more frequently [76,77]³⁾ and show a reduced ability to disengage from stimuli [74]. From the second half of the first year, some researchers have reported that preterm children exhibit shorter periods of focused attention, while others have not observed this difference [77]. At

the beginning of toddlerhood, when the executive system gains greater control of the arousal system, sustained attention becomes more apparent in preterm children.

Risk factors concerning the development of attention have been reviewed. A low BW negatively influences bio-neurological development, and there is a strong association between a lower BW and a greater number of attention problems [78]. A shorter gestational age predicts poorer attentional skills [31], and a higher medical risk seems to negatively influence attention. However, these observations have been difficult to assess due to the interference of other variables such as birth weight (BW) and GA [79]. In addition, male gender [71], decreased maternal psychological well-being and less successful parent-child interactions throughout childhood [80] have been identified as risk factors. Attention problems have been further reported to partially mediate the relationship between the birth condition (prematurity) and subsequent behavioral problems [81].

Some authors have concluded that children born preterm gradually develop more severe attention problems [ibid], which is consistent with the results of a large French study [82] reporting a greater number of attention problems at age five and of a Norwegian population-based study reporting a slower reaction time, reduced awareness and greater attention problems at age 11 in preterm children compared to term-born peers [83,84].

1.3.4 Social problems

Social problems are related to the adaptation of a child to his or her social surroundings. A tri-component model has been suggested as a conceptual framework to describe the features of social competencies and thereby of potential problems [85,86]. These interrelated components are described as follows: ¹⁾ social skills (cognitive features such as theory of mind, compliance, attention and behavioral skills), ²⁾ social performance (the ability to use skills in appropriate ways in different contexts) and ³⁾ social adjustments, which include, e.g., the ability to form high-quality friendships, several friends, a high level of sociability and low levels of social withdrawal [86]. Importantly, each component level builds on the ability of the child to function at lower levels.

During the first years, it may be difficult to separate socio-emotional from attention problems and vice versa [69]. Parent-reported socio-emotional problems among two to three-year-old children have been estimated to occur 10 to 15% of children [87,88], and several reports have focused on the

problem of the under-detection of such problems in toddlerhood [43,87,89]. Social problems seem to be more easily perceived by caretakers when the child begins to function more independently. An immature or unsuccessful social adaptation may be highly influenced by less successful parent-child interactions [90]. The statements used to identify social problems in pre-school years may concern, e.g., clinging behavior, not going along with other kids, being teased, preferring younger playmates [48-50]. Social competence may be viewed as culturally appropriate manifestations of behavior at the opposite end of the problem behavior spectrum [87,91].

After the first year of life, poorly regulated behavior (frequently referred to as the “terrible twos”) can be identified as a precursor of social problems [44]. Social difficulties may first become visible as a lack of interest in social interactions during infancy, while an inability to manage peer interactions may characterize toddlerhood [37]. During the preschool years, peer interactions become more complex, and the ability of children to regulate arousal, cooperate and form friendships may be observed as more or less successful behaviors. If socio-emotional development decelerates, it will interfere with other features of development such as perception, learning, and achievement, and it will frequently become evident as dysregulated behavior [44].

Socio-emotional problems have been documented as highly stable [86,87]. Approximately 35% of children who were rated by their teachers as having elevated socio-emotional problems during the first year in elementary school had been identified with worrisome test-scores based on standardized measurements before age of three. 68% of those with parent-reported psychiatric disorders had been identified as having problems as early as during toddlerhood [44,88].

Very preterm children, especially those that are extremely preterm, have been reported to exhibit elevated social problems throughout childhood [92-94], and this tendency was confirmed in a recently published review [86]. Features of social adjustment in particular create a gap between VLBW children and full-term peers, while there is less consensus regarding features of social performance. However, VLBW children seem to catch up with their peers born full term in terms of the levels of prosocial behavior [ibid]. Those children who were reported to exhibit the greatest social difficulties across the studies had the lowest birth weights, were males and had reduced intellectual functional ability [66,86].

Children born preterm are at a high biological risk and have been described as being more dependent on maternal sensitivity to facilitate their social engagement [95,96]. Strong associations have been described between the biological regulation concerning the sleep/awake cycle and heart rate variability during the last trimester of pregnancy and later social rhythms/parent-infant interactions during infancy [41]. Parent-infant synchrony [40] and low levels of parental negativity [97] are also known to predict better infant self-regulation and socio-emotional adaptation.

1.3.5 Thought problems

Thought problems have a low prevalence [49,50]. These problems have been reported by parents and teachers as, e.g., fixed thinking about something, hearing things, repeating acts, seeing things, staring blankly, and expressing strange behavior or ideas. This subscale in the ASEBA's questionnaire is meant to intercept behavior associated with early psychopathology, e.g., schizophrenia [ibid].

1.3.6. Mental health among children born preterm

Two Norwegian population-based studies of low birth weight (LBW) children have reported impaired mental health outcomes during late childhood and as young adults compared to the sample born full-term [98,99]. Elgen et al. reported a three-fold increased risk of psychiatric disorders, and only 50% of the LBW young adults (age 19) reported good mental health throughout adolescence [ibid]. Affective, anxiety, and attention deficit hyperactivity (ADHD) disorders and antisocial personality disorders were the most common mental health problems, and 20% of the population had more than one diagnosis. The populations reported in the above-mentioned studies were born in the 1980s, and the results are consistent with previous reports from other countries [25,79,100,101].

1.4 Quality of life (QoL) of children born preterm

The QoL of former preterm children has been investigated extensively the last few decades, mainly by parental proxy reports on their children but also based on the self-reports of adolescents and young adults [102-104]. QoL is defined by the WHO as “an individual’s perception of their position in life in the context of the culture and value systems in which they live and in relation to their goals, expectations, standards and concerns” [105]. The QoL concept is a highly multidimensional construct that attempts to measure subjective experiences. Decades ago, a Danish psychiatrist described QoL based on four classes of needs: biological needs, the need for warm human relationships, meaningful occupations and the need for diverse and exciting experiences [106,107]. Quality of life has also been separated into subjective and objective features in which the final description is based on societal standards, whereas the subjective part of QoL is based on individual life experiences and preferences [ibid]. An overview of the basic features of QoL in children and adolescents has also been described by Mattejat [108] and by Jozefiak [109] (Figure 1.).

Researchers investigating the QoL of children have used different conceptions or definitions of the concept (QoL, health-related QoL, well-being, among others) that are related in part to the type of study or measurement used [104]. Zwicker and Harris identified 6 studies that examined QoL in preschool-aged children who were born preterm [ibid]. Significantly poorer physical functioning was reported in four of these studies compared with the reports of children who were born at full term. Preterm children have also been reported to exhibit less competent social functioning and, in some studies, greater anxiety compared with children born at term [110]. ELBW children in particular have been reported to have a generally lower QoL than their peers born at full term [102,104,111]. Health-related QoL has also been reported by Norwegian parents for children aged 10 years who were born extremely preterm [112]. This Norwegian study indicated that learning and attention problems negatively influence the daily life of children and that boys in particular struggled more than their full-term counterparts.

Self-reporting has been used among preterm children who have reached adolescence, being approximately at 14 years of age [113]. Teenagers born preterm did not report such a large reduced QoL as their parents report compared with their full-term peers [104,114,115]. In a Norwegian study, VLBW teenagers with cognitive deficits had significantly lower global health and behavior, general health perception, self-esteem and family activities [99]. A review by Zwicker and Harris concluded

that the impact of prematurity is greatest during the younger years, and this condition to some degree even negatively influences life during adolescence and adulthood [104].




Figure 1. Basic features of Quality of Life

Figure originally published by T. Jozefiak [109], after Matzejat [108], and reprinted with permission.

1.5 Development of competencies up to middle childhood

The development of competencies is highly influenced by the ability of children to adapt to contexts and challenges [1,116]. Competencies may be difficult to discuss separately from behavior problems because they largely describe similar phenomena, which are described as resources or problems. Children born preterm have frequently been reported to possess impaired physical performance, reduced intellectual and social functioning and poorer school performance [19, 57,117,118]. Outcomes are mostly evaluated by comparing means at group levels. The complexity and nuances that could become visible at an individual level are not a focus.

While behavioral problems are evaluated by pre-defined statements, behavior as competencies in the ASEBA-system [47] are assessed by parent or teacher evaluations of the child *in comparison to children of the same age*. In terms of sports and activities, parents are asked to evaluate how much time the child participates in and to what degree he or she has been successful with the preferred activities. In relation to school performance, parents and teachers evaluate the competencies of children in subjects such as first language, math, history and other subjects [49,50].

A recent meta-analysis reported that VLBW children perform at the same level as full-terms in terms of language achievements across the pre-school years, while their performance in arithmetic was poorer [117]. It has been frequently stated that the burden of prematurity becomes apparent when these children are faced with greater demands from society (pre-school, school, among others). Similarly to the ability of attention and social problems to moderate QoL, it also affects the ability of children to manage new demands and expectations in a classroom setting. Children born preterm are often described as less ready for school and frequently have a delayed start in primary school [119]. LBW children have been reported to receive special educational services in primary school twice as frequently as the general population [83]. Similar results were reported in a sample of children who were born moderately to late preterm; 7.7% of the children received extra services in school versus 2.8% of the general population [57].

The behavior problems and development of competencies of children are associated with biological conditions as well as psycho-social factors such as parent-infant relationships and parental well-being [24, 117, 120-122]. Some authors have suggested that there is a differential impact of biological and psychosocial risk factors at different ages as the psychosocial risks increase in importance with age [123].

1.6 Parental adaptation to the caretaking of a child born preterm

1.6.1 *Enjoyment of and attachment to the infant*

Bromwich suggested that the behavioral establishment of an affective base can lay the foundation for parent-child attachment [124]. This attachment is frequently typed as *bonding*. Different interactional patterns have been described between mothers with children born preterm compared with mothers with full-term children as early as the 1980s [125]. At four months, preterm children were less responsive than their full-term peers despite the heightened levels of involvement of their mothers. At the age of two, mothers of preterm children performed less positive scaffolding and were less involved in interactions with their children compared with the mothers of full-term children. This finding may be considered to be an early observation of the *dependency of parents on children's responses* to their parenting efforts to enable them to continue good work. Parenting of children born preterm has been described to be associated with increased challenges because the children show fewer positive affects and smiles [126], more fuzziness and inattentance [6] and less stimulation of their mothers to perform spontaneous, intuitive maternal behaviors (so-called "motherese" characterized by kissing, snuggling, touching, child-directed talk and physical contact) [127]. Others have studied interactional patterns between mothers and 6-month-old infants and compared premature and full-term at children's age of 18 months on later outcomes [128]. A maternal "controlling pattern" was observed in almost 28% of the mother-preterm infant dyads compared with 12% of the full-term infant dyads. At 18 months, preterm infants in dyads with a controlling pattern had significantly fewer positive outcomes compared with preterms in cooperative pattern dyads and with full-term infants [128].

The impact of early impaired contact between children born preterm and their parents has been studied extensively during recent decades. Schmid et al. reported a long-term impact of maternal responsiveness and early interactions, and their results indicated that a less successful early interaction was associated with a higher risk of depression in offspring up to the age of 19 years [129]. The importance of responsiveness, sensitivity and synchrony will be described later in this thesis as possible mechanisms that play a role in the intervention.

1.6.2. Parenting stress and impact on parenting behavior

Parenting a child that is born preterm is associated with more long-term stress, even in families with a high level of socio-economic resources [130]. Consistent with the early theory of psychological stress is the concept of *parenting stress*, which has been described as a complex process consisting of four components: an *external event* met by a *cognitive appraisal* and followed by a *mobilization of individual coping mechanisms* and finally followed by a *stress reaction* in body and mind that is recognized as a parental behavior or affect [131,132]. Parenting stress is distinct from stress related to other life events and is described as a complex response to the demands of parenthood [ibid]. Parents with reported high levels of stress are more likely to have an authoritarian, harsh and negative parenting style, in which the parent is less involved with his or her children and foster children with a more unsecure attachment quality [132].

Abidin defined two dimensions of parenting stress [133]. They distinguish between features related to parental attributes (aspects of depression, attachment, competence, role restriction, isolation, spouse and health) and child attributes (aspects of adaptability, acceptability, demandingness, mood, distractibility/hyperactivity and ability to reinforce the parent [133,134]. The association between the main dimensions and sub-aspects are described in Abidin’s model of parenting stress shown in Figure 2.




Figure 2. Theoretical Model for PSI

Reprinted with permission from the author (Abidin) [133].

Individual differences in parenting stress are stable over time, and they are associated with the quality of the parent-infant relationship and are essential to address its bidirectionality; parenting stress appears both as actions and as reactions [132].

Reducing parenting stress is considered to be important because it strengthens the mental health of the parent [132], may decrease the impact of maternal depression on parenting behavior [134] and improves the efficacy of interventions that target the sensitivity and responsiveness of the parents [135] and behavioral problems of the children [136]. This finding has been reported to be especially important among the parents of preterms because greater parenting stress has been reported repeatedly in such families [130,137], and parenting stress may have greater negative consequences on children born preterm than on children born full-term [138-140].

Maternal depression has been reported to have a universal negative effect on mother-child interactions across cultural and socio-economic differences [141], and maternal mental health problems have been reported to occur more frequently in families that have reared children born preterm [121] and also to have a significant impact on the outcomes of children born preterm [142,143]. However, paternal depression has also been reported to have a significant and deleterious effect on the parenting behavior of fathers [144]. Essex et al. reported that maternal depression and stress beginning in infancy are the most potent predictors of subsequent stress regulation and cortisol levels [145]. Preschoolers with the highest cortisol levels at the age of 4.5 years exhibited more severe mental health problems after enrollment in school [ibid]. This finding corresponds to the results of a Swedish study that reported a significant relationship between elevated parenting stress in children aged 1 and 8 years and the cortisol levels of children at 8 years of age [146]. Both studies indicated that long lasting levels of maternal stress during childhood may be more influential than an increase in maternal stress during only early or late childhood. Other groups have identified different trajectories of parenting stress during early childhood (stable high, decreasing or increasing stress) and both maternal, child and contextual factors accounting for stability and changes in trajectories [147]. Parenting stress has been effectively reduced by offering parenting education components [132,148], but other actions such as improving maternal-child attachment [149], mutuality [150] and responsiveness [151] have been suggested to be equally important.

The cognitive appraisal by a mother of an external event is one step in the generation of parenting stress [133], and Allen et al. reported that mothers who perceive their preterm children as very vulnerable at the time of hospital discharge tend to continue to have this perception [152]. This perception has been associated with less positive development over the first year of life of children born preterm [ibid]. A review of interventions that address the impact of parenting stress on the development of high-risk children highlighted the importance of the direct involvement of parents [132, 152]. Family-focused strategies have been reported to be more effective than interventions that mainly focus on the child, and a focus on the parent-child relationship appears to be more successful than focusing solely on the behavior of the parent [151].

1.6.3. Contribution of the children to the parent-child interactions

The contributions of infants to parent-infant interactions are important and have previously been summed up as three features of social competence: their predictability of behavior, social responsiveness and readability of cues [154]. While full-term infants are equipped to handle conditions of instability in the extrauterine environment, prematurely born infants are unable to readily adapt [155]. Immaturity, neonatal medical conditions and inappropriate environments continuously affect the physiology, behavior and integration between biology and the environment in these infants. When they interact with the environment, they may respond in either an organized or a disorganized way. The behavior of preterm infants is characterized by somewhat unpredictable fluctuations in autonomic, motor and state organizations, and they are less socially responsive than full-term infants and are less able to inform their caregivers in an understandable way about what they need [9,11]. Some responses may be appropriate according the level of maturity but are also frequently influenced by some level of disorganization. These deficits and lack of early social readability place them at risk for interactional difficulties [156].

Early alterations caused by preterm birth may also influence later development. The period during which preterm birth takes place is considered to be a critical developmental window. It's described as a disruption of organizational events that causes the brain of the preterm baby to be organized in a different/immature manner compared with that of a full-term [19].


1.7 Early interventions addressing developmental difficulties

Forty years ago, Bromwich outlined three assumptions concerning how to enhance healthy development in infants and the development of supportive parent-child interactions [124]. The first assumption was that parent-infant interactions are *reciprocal processes* in which the behavior of each participant affects the responses of the other. The second assumption was that *mutual satisfying relationships* between the parent and child are an essential premise for the later successful development of the child. Finally, the third assumption was that parental *competencies grow concurrently* with increasing responses from the child, which provides positive feedback to the parent. These assumptions are in agreement with the descriptions of transactional relationships and mechanisms in human development, which will be presented later [157].

Bromwich described six stages of maternal behavior. The first, second and third stages describe the establishment of an affective base for later interactions. The next three steps describe how the parent becomes increasingly capable of initiating developmentally supportive activities with the child, thus generalizing the impact of new activities and further generating new and appropriate activities and experiences as the child achieves new developmental levels. In her view, a core rationale is that the *interventions should try to enhance the quality of mother-infant interactions* and not merely build on the instructions and teaching of the parents. The limited effects of instruction-based interventions are already described, and researchers have searched for alternative strategies [156].

Six stages of maternal behavioral progression:

1. Enjoyment of the baby
2. Sensitivity and responsiveness to infant cues
3. Mutually satisfying time together
4. Awareness of developmentally appropriate activities
5. Ability to generalize insights about activities and devise alternatives
6. Adaptation to achieved knowledge regarding new developmental levels during infant growth.



The early history of interventions with low birth-weight infants included a variety of direct and indirect preventive approaches. The theoretical perspectives included both the causes of parenting failures and the psychosocial and developmental maladjustments of the child [158]. Several areas,

which may disturb the early parent-child relationship, have at some point been identified as targets for intervention [153,159]:

- Early separation and failure of parental bonding [5,6,160],
- Stimulus deprivation programs [158],
- Deficits in infant capacities to elicit care [10,161,162],
- Parental emotional crisis related to preterm delivery [163],
- Pre-, peri- and postnatal medical complications [21,164] and
- Adverse childrearing environments [157,165].

The Mother-Infant Transaction Program (MITP), which was tested in a sample of prematurely born children in the present study, was designed in the late 1970s [166]. At that time, the parents were only short-term visitors in the NICU, and many parents had to resume their working duties while the child was hospitalized. A majority of the parents saw their child for only a limited number of hours before discharge. The formation of the MITP program was highly influenced by contributions from Bromwich based on her description of the maternal stages of behavior [124], as well as by Als & Brazelton based on their descriptions of the hierarchy of organizational levels that are critically important for the understanding of the behavior of preterm infants [9,10,11,168].

The conceptual framework described by Als was termed synactive because *“at each stage of development and each moment of functioning, the various subsystems of functioning exist side by side, are often truly interactive, but are also often in a holding pattern, as if providing a steady substratum for one of the system differentiation processes”* [168, p. 230]. This theory describes how one could observe the way an individual child handles the experiences of the world around him/her. The “observer-window” in this framework offers the interpretation of signs from the subsystems: autonomic system, motor system, state/organizational regulation, attention and interaction system and self-regulatory system [10,11]. A basic assumption was that the subsystems were continuously interacting with one another, like a child continuously interacting with his or her environment [10,11]. This observation is in accordance with the transactional understanding of development, which has informed the design of the MITP as well as this study [166,169].

1.8 A transactional model of development

The design of the Tromsø Intervention Study on Preterms (TISP) are consistent with the original MITP study [116,166,169], in which it was understood that nature (biology) and nurture (environment) had an equally important and interconnected impact on the development of children. This theoretical framework, a transactional understanding of development, is described in the studies conducted by Sameroff and colleagues [157,170]. In our study of the development of children born prematurely, this framework informs the following: ¹⁾ both the child and its environment influence later development, ²⁾ the experiences of the child are a driving force of development together with all of the experiences facilitated by differences in the biological premises, intelligence, personality and perception of children, and ³⁾ the development of children is a product of the continuous, dynamic, bidirectional interactions between a child and the experiences provided by the social setting [170].

The differences between interactions and transactions can be explained by their effect on those involved. Interactions may be viewed as social exchanges informed by culturally appropriate responses that are consistent with the expectations of the recipients. However, transactional sequences require something new in the exchange between dialog partners because they must adapt to the responses from one another. *“Developmental changes are defined by changes in the way the child interacts with experience. They are driven by new complexities in either the individual or experience that require new adaptations in one or the other. In some areas the complexity already exists in the experience; in others it comes into play as the child reaches increasing levels of maturity”* [116, p.9]. At some age or level of development, most children show a greater continuity of behavior. This may be caused by some failure in either the child or the environment, or the adaptive process ends because the child has become an adult. Adulthood may be viewed as an equilibration in which individuals and environments are no longer in a state of adaptation. *“At this point, transactions become interactions”* [116, p.9].

Another central concept in the transactional framework is the issue of *regulation*. D’Apolito defines regulation as a state in which experiences are handled and regulation generates successful adaptation and possible development [155]. Self-regulation is defined as a cornerstone of early childhood development that cuts across all domains of behavior [1, p.26]. Others such as Feldman highlight two central features of regulation in the investigation of associations between early

neonatal biological regulation and later social regulation during the preschool years [41,171]; that is, the balance between mechanisms of excitation and inhibition and the ongoing interplay between environmental challenges and internal organization. Sameroff emphasizes the complex construct of regulation as a central phenomenon in modern developmental theory and exemplifies this balance. First, evolution is not restricted to biological change, and transactions resulting in or following regulation occurs in complex social settings. Second, the development of an individual child may be judged as a co-construction of the self-regulatory abilities and environmental “other-regulative” competencies of the child, which impede or facilitate development. Finally, Sameroff argues that some children may have severely compromised self-regulative capacities or their environments may be so chaotic that developmentally supportive transactions seem impossible [169]. Sameroff and Fiese emphasize that although the concept of self-regulation may provide an illusion that regulation is a property of the individual, it can only occur if there is a social surrounding exerting “other regulation”.

1.9 Mechanisms involved in developmental change

The importance of parent-child attachment (bonding) and the historical roots of the concept have been mentioned previously [3, 4]. New research within that field has given increased attention to the concept of bio-behavioral synchrony between attachment partners [171,172], which is described as online physiological and behavioral co-regulation. Oxytocin hormone is known to play a central role in the establishment of breastfeeding, helping to let down the milk. There has been less focus on how the regulations of hormones interplay with the formation of affiliative bonds both in infancy and later in life. Oxytocin, the so-called “hormone of love”, appears to be deeply grounded in human biology to support the formation of social bonds, care, security and healthy child development. Feldman defines affiliative bonds as “selective and enduring attachments that are formed on the basis of repeated exposure to coordination between physiological states and interactive behavior within each partner, between partners, and between the physiology of one and behavior of the other” [173,174].

This perspective, which appears to be embedded in transactions, may be of great importance for the early formation of parent-child relationships in families with children born preterm. Early

mother-infant synchrony is highly influenced by the successful maturation of internal hormonal regulation in the child [41]. From the beginning of the last trimester, both sleep-awake cycles and the regulation of arousal are undergoing important organizational development. The success of this maturational process seems to lay a foundation for the quality of mother–infant synchrony, which is described as a core activity in the social and emotional development of children [ibid].

Attachment difficulties have been described to occur frequently in preterm mother-child dyads [175,176], and gestational age has been reported as a moderate to strong predictor of maternal attachment difficulties [177]. The disturbance of early emotional bonds between parent and child may have long-lasting negative effects on the caretaking abilities of parents [177]. Interventional effects on attachment were not a focus of this study per se, but a disruption in parental attachment as a source of parent-related stress has been assessed [179]. The oxytocin regulative system is reciprocally engaged with the hypothalamic-pituitary-adrenalin axis (HPA) that mediates the stress response, and neuro-anatomical evidence suggests that there is a mutual regulation of the oxytocin and HPA systems [170,180]. This finding has shed light on Abidin's detection and inclusion of an attachment-related subscale in the parenting stress questionnaire he designed more than thirty years ago [133]. The influence of both child- and parent-related parenting stress on child development has already been outlined and is also a source of transactional processes.

The bio-behavioral regulation that was briefly described above influences moment-to-moment interactions between parents and children. Synchrony is not only a matter of physiological and behavioral timing, but it also affects the interactional quality. It seems to influence parental sensitivity and responsiveness, which are core qualities in successful parent-child interactions [123,181].

During the last decade, the impact of early adverse experiences and neonatal stress on basic neurological structures has been described through collaborative research across biological, behavioral and social sciences, including knowledge of neuro-science [1,182,183]. Als et al. reported how developmentally appropriate care could protect the immature brain of the preterm baby against structural alterations caused by neonatal distress [184], while Milgrom et al. reported how

parental sensitivity training across the NICU stay were associated with improved white matter in the brain of the preterm child at 40 w of GA [185].

The above-mentioned studies confirm that early environments shape and calibrate the functioning of biological systems [186]. American Academy of Pediatrics (AAP) have designated this phenomenon as “toxic stress” and define it as stress that is extreme and frequent, extending activation of the stress response without the buffering presence of a supportive adult [ibid]. Current evidence suggests that the early experience of stress catalyzes a series of biological adaptations that change the way the brain, neuroendocrine stress response, and immune system function. Central to this idea is the biological regulation related to the hypothalamic-pituitary adrenal (HPA) axis. The HPA axis is involved in metabolic and cardiovascular responses to acute and chronic stress. The following is concluded: *“One of the primary consequences of early life toxic stress is HPA dysregulation, as the developing neuroendocrine system is chronically pressed into action”* [ibid, p.321, 187]. Preterm infants are highly vulnerable to exposure to stress [32,35, 140,188]. During the last trimester of pregnancy (30 to 40 w of GA), these infants may be exposed to adverse environments in a NICU while brain growth and maturation occurs at a high speed (myelination, migration, and synapse formation). The belief of an early critical period in child neuro-psychological development has been suggested for a long time [158] and was subsequently critically reviewed [1,42]. The importance of the experiences of the infant during the first months and years of life has been highlighted and expressed as *“not because this period of development provides an indelible blueprint for adult well-being, but because it sets either a sturdy or fragile stage for what follows”* [1,p.5]. In other words, this knowledge may shape bridges between the known neuroanatomic and neuro-developmental vulnerabilities in children born preterm and later developmental difficulties [19]. This concept was recently discussed by Feldman and colleagues, and the synchrony in coordination between biology and behavior during social contact has been suggested as a mechanism for early periods of sensitivity via effects on the social brain, regulation of oxytocin and adult sociality [173,174].

2. Aims and questions

To examine the long-term effects of an early intervention among children born prematurely with a birth weight < 2000 grams and the influence on their surroundings and parents.

1. Behavioral-emotional development of the children up to nine years of age.

- i. Does an early intervention influence long-term behavioral-emotional development, as reported by parents and teachers up to nine years of age?
- ii. Does an early intervention influence behavioral outcomes, social competence and adaptive behavior in school, as perceived by parents and teachers by seven and nine years of age?
- iii. Are specific birth weight groups or genders particularly affected by the interventions?
- iv. How does the behavioral development of preterms (with or without intervention) compare with the behavioral development of full-term children?

2. Development of parenting stress until children's age of nine.

- i. Has the early intervention influenced the longitudinal development of parenting stress as reported by mothers and fathers?
- ii. Are there cross-sectional differences between the preterm groups in the reports of mothers and fathers regarding parenting stress up to nine years of age?
- iii. How is the development of stress reported by parents in the two preterm groups compared with that reported by the parents of term controls?

3. Child and parental proxy reports of quality of life (QoL) at nine years of age.

- i. Did the early intervention influence the self-reported QoL and parental proxy reports of QoL of preterm children at nine years of age?
- ii. Did the intervention affect the level of agreement between the child and parental proxy-reported QoL in the two preterm groups?
- iii. Was QoL, as reported by children and parents in the two preterm groups, similar to that reported by children and parents in the term reference group?

3. Materials and methods

3.1 Participants

The studies presented and discussed are part of a larger, comprehensive study: the Tromsø Intervention Study on Preterms (TISP) [169]. TISP is a randomized controlled study of preterm infants with a BW < 2000 g who were born at the University Hospital of North Norway between March 1999 and September 2002 and a replication of a previous American study [166,167,189-191]. From the counties of Finnmark and Troms, 91% of all children born prematurely were recruited for the study. Children born without severe neurological or sensory impairments and with Norwegian speaking mothers were eligible for study inclusion. Twins were consequently recruited to the same group, while triplets were excluded from the study due to the character of the intervention. Decisions on sample size were based on the results of a previous study by Achenbach et al., and a difference of 7.5 points between groups in Bayley's MDI at age 2 (corresponding to a difference of 0.5 standard deviations) with 80% probability was calculated [169,192].

When eligible infants reached an age of 32 w of GA or more, the study coordinator informed their parents about the study and asked about participation. A total of 203 infants with a birth weight below 2000 g were recruited for the study. Thirty-five of these infants were lost to participation (14 died, 13 had non-Norwegian speaking parents, six were triplets, one was diagnosed with Down syndrome and one was not asked). The parents of 22 infants refused to participate in the study, while those of the remaining 146 preterm agreed to participate in the study. The coordinator collected written informed consent from all of the parents. The preterm infants were randomized in blocks of six using computer-generated numbers to form a preterm intervention group (PI, n = 72) and a preterm control group (PC, n = 74) and stratified according to a gestational age < 28 weeks and ≥ 28 weeks. Randomization was conducted by the research department of the hospitals, and the study results were available in numbered, sealed envelopes. Three children were excluded because of blindness during the two first years of the study. In addition, three children were excluded because of deafness; however, these children were again included in the study later in childhood after they regained their hearing with a cochlea implant. Children who were identified with disabilities at later follow-ups continued in the study if they were able to participate in the

age-appropriate assessments. The flow of the study from inclusion up to nine years of age is presented in Figure 3 [Appendix, part II].

Healthy newborn infants (GA \geq 37 weeks and BW > 2800 grams) were recruited from the well-baby clinic at the same hospital to a term reference group (TR, n = 75). They were recruited by asking the parents of the first healthy newborn born after each preterm infant was recruited into the study. If the family refused to participate, subsequent families were asked sequentially. For practical reasons, all of the families recruited to the reference group lived in the area close to the city of Tromsø. Baseline information for each group is presented in Table 1 [Appendix, part I].

3.2 Study implementation

Eight experienced nurses performed the interventions. They received approximately one week of education before the pilot interventions and later implementation was started. The education included an introduction to developmental psychology and theory about state regulation, reflexes and how to initiate episodes of mutually responsive interactions with a newborn. In addition, four of the eight nurses were trained as NBAS examiners in the Neonatal Behavioral Assessment Scale (NBAS) during the study planning phase [191]. This extended knowledge throughout the entire intervention group because the development of the MITP intervention was highly inspired by the NBAS [10,193]. The initial educational introduction, training with video feedback and follow-up meetings across the 42-month inclusion period ensured a strong mutual understanding of the aims of the intervention that had been translated into Norwegian [194].

Families that were randomized to the intervention met the same intervention nurse during all of the sessions. This nurse may have met them earlier in the course of the NICU stay but was never one of the primary contacts of the family. The scheduling of intervention sessions was flexible because the nurses often postponed meetings if the child was in an inappropriate state (e.g., recently fell asleep during a session focusing on the capacity to focus alertness and social interactions). The majority of the pre-discharge sessions were performed during the final week in the hospital, but a few were conducted in the home of the family in situations with an accelerated discharge, or two sessions were given on the same day at the end of the hospital stay (e.g., morning and afternoon), which is consistent with the original study [166].

Families in the PC group were provided with the written guidelines of the NICU concerning treatment and information before discharge. This included a session with a demonstration and instructions regarding baby massage conducted by a child physiotherapist. Some parents became disappointed when their child was randomized to the preterm control group. In such cases, the parents were offered additional interactions about their baby if requested. The TR group was examined once by a pediatrician in the maternity ward and routinely on the third day of life.

3.3 The intervention program

The intervention program was a modified version of the Mother-Infant Transaction Program (hereafter referred to as MITP-M). MITP was designed by Rauh & colleagues in 1980 [166]. The modification introduced in the TISP study consisted of an initial session during which the parents of each child met their intervention nurse for a discussion in the absence of the infant present. This session provided a period to express feelings of grief and talk about experiences during pregnancy, delivery and the NICU stay. In addition, the parents were given the opportunity to become familiar and more relaxed with their intervention nurse. The mothers participated in all of the intervention sessions, and the fathers participated in 6 of the 12 sessions (54% of the intervention) with an inter-quartile range of 4 – 10 sessions.

The original MITP intervention consisted of seven one-hour sessions with the parents and their baby during the final week before discharge, and four home visits in the family's home at 1, 2, 4, and 12 weeks post-discharge. All of the hospital sessions were performed in a separate, quiet room with the baby, mother, father (when available) and intervention nurse. The interventions were adapted to the needs of each family and the natural speed with which the parents could learn and strictly follow the topics described in the MITP manual [166]. All of the sessions consisted of different agendas, and their main topics are briefly described in Figure 4. The contents of the MITP-M are provided in the Appendix, part III.

The finalization of the program at this age was justified by ending the original study [ibid] because it made up an intervention that could realistically be implemented with the financial and professional resources available. Unlike the original study, no logbooks were given to the families after the completion of the interventions. Completion of the intervention was monitored by a review of the

logbooks by the study director.

3.4 Measurements

The study focused on the number of reported behavioral problems. Behavioral development has been reported using standardized measurements that assess a broad range of competencies and areas of concern [49]. The analysis depends on reports of problems from parents and teachers and on measures of specific attainment of developmental milestones during childhood. Developmental goals can comprise specific motor, behavior or communicative skills, which have been reported previously [195].

When the socio-emotional and behavioral developments of children are analyzed through the lenses of reported behavioral difficulties, this phenomenon may be viewed as looking in a mirror. The extent of behavioral problems may indicate a greater or lesser struggle to comply with parental expectations, adapt to both one's own and environmental expectations and reach developmental milestones at age-appropriate stages.

Child Behavior Checklist (CBCL/2-3 and CBCL/4-18)

The behavioral problems of the children were reported by mothers and fathers on the CBCL/ 2-3 during toddlerhood and similarly on the CBCL/4-18 for children aged 5, 7 and 9 years [48,49]. Both questionnaires consist of statements (100 statements in the CBCL/2-3, and 113 in the CBCL/4-18). Each statement is transferred to a Likert scale of 0 to 2 (0 = not true, 1 = somewhat or sometimes true, 2 = very or often true). The parental reports are based on their perception of the child during the last 2 months across toddlerhood, while in the preschool/school years, it is extended to the perception of the behavior of the children during the last 6 months. Two broadband behavioral domains were calculated in both questionnaires (internalizing and externalizing behavior). These main dimensions are slightly different because questions about somatic complaints are based on internalizing behavior in childhood but not in toddlerhood, and the questionnaires has some different items related to behavior that are specific to the different ages. In the CBCL/2-3, two sub domains remain outside the broadband internalizing and externalizing syndromes (sleeping problems and somatic complaints), while CBCL/4-12 consists of three supplementary sub domains (social problems, thought problems and attention problems). In addition, CBCL/4-12 comprises

several questions that address the perceptions of parents regarding infant activities, social competence and competence in activities of daily living compared with their peers. The CBCL questionnaires were revised in 2001 and later modified in 2007 [47]. In the current study, the 1991 versions of the questionnaires were used across all follow-ups to facilitate the longitudinal analyses.

Teachers Report Form (TRF)

The TRF questionnaire is designed for teachers who have observed the behavior of a child over a period of at least two months [50]. Similarly to CBCL/4-18, the TRF consists of 113 statements and is scored on a Likert scale similar to CBCL. Approximately 15% of the statements are different due to differences in behavior that can be observed in a classroom/school setting compared with a home setting. Similarly to CBCL/4-18, the TRF provides standardized measurements of child and adolescent emotional/ behavioral problems and measures of competencies. Behavioral outcomes are summarized as total problems and scaled according to internalizing and externalizing dimensions. Internalizing behavior consists of the following subscales: withdrawn, anxious/depressed and somatic problems. In contrast, externalizing behavior consists of the following subscales: aggressive and delinquent problems (both in CBCL/4-18 and the TRF questionnaire). Behavioral problems are also rated on subscales related to thoughts, social life and attention.

Strengths and Difficulties Questionnaire (SDQ)

For children aged 5 and 9 years, mothers, fathers and teachers responded to selected statements and questions that were extracted from the SDQ [196]. In an attempt to avoid overlapping questions in different questionnaires, only 10 statements related to the social behavior of the children and some questions concerning the perception of parents and teachers of the difficulties of the children in everyday life were used. The statements were answered on a 3-point scale (agree, partly agree, do not agree), and the questions were answered by choosing one of four alternative answers [197]. The questionnaire is included in the Appendix, part IV.

Parenting stress index (PSI and PSI-sf)

The Parenting Stress Index-Full Form (PSI), 3rd. edition, was used for all assessments up to the age of seven years, while the Parenting Stress Index - Short Form (PSI-SF) was used for nine-year-old children [133]. The PSI consists of 120 questions covering three main dimensions of stress (child-, parent- and life-related stress). The PSI-SF consists of 36 questions that were extracted from the

parent- and child-related dimensions. Both questionnaires were translated and back-translated in cooperation with the creator (Abidin) [personal information, JAR, 2013]. A 5-point Likert scale ranging from “strongly agree” to “strongly disagree” made up the response alternatives on both questionnaires. The initial assessment of parenting stress among the mothers before randomization had limited success because many of the statements were left unanswered. A calculation of the responses on those scales sufficiently answered indicated that the initial level of parenting stress was similar in the two preterm groups. At six months, only one parent provided a report (mostly mothers), while both mothers and fathers provided separate reports on all subsequent occasions.

The PSI consists of two main dimensions that address child- and parent-related stress (covered by 101 statements/questions) and a life stress section (22 events or conditions that are often reported as stressful but may not be related to the current parenting challenges). The Child Domain contains 47 statements covering six subscales: Distractibility, Adaptability, Reinforces parent, Demandingness, Mood and Acceptability). The Parent Domain contains 54 statements covering seven subscales: Perceived competence, Isolation, Attachment, Health, Role restriction, Depression and Relation to Spouse. A total stress score was also computed based on all of the items excluding life stress questions. The theoretical model underlying the construct of the PSI is shown in Figure 2.

The PSI-SF is reported as a Total Stress score (TS) and based on three subscales, each of which consists of 12 items: Parental Distress (PD), Parent-Child Dysfunctional Interaction (P-CDI) and Difficult Child (DC). Some questions in both questionnaires are used to calculate a Defensive Responding score, which indicates the degree of potential inconsistent/denial reports by respondents.

Both the PSI-FF and PSI-SF are frequently used in research [198,199], and the correlation between TS for these two measures has been described as high (0.87) [133,200]. The PSI-SF, Difficult Child subscale consists solely of items from the Child Domain in PSI-FF, and Parental Distress subscale items are from the Parent Domain. The P-CDI subscale includes items from both the Child and Parent domains and focuses on parental perception of transactions with their child and their expectations about the behavior of the child [133]. The Norwegian versions of both PSI-FF and PSI-SF were translated by Rønning and Abidin and were used in this study with the permission of Abidin and Psychological Assessment Resources, Inc. (PAR) [201]. The questionnaires include some literal

differences such that questions in the PSI-SF may be perceived as more negative and definitive than those in the original full format of the PSI.

Kinder Lebensqualität Fragebogen (KINDL)

The Kinder Lebensqualität Fragebogen (KINDL) questionnaires have been translated and validated for use in Norwegian populations [202,203]. The translation was conducted by experienced Norwegian school teachers with university degrees in German [204]. The questionnaires are short, generic and consist of a self-report questionnaire (Kid KINDL[®]) that is appropriate for children (7 to 13 years old), and a questionnaire for the parental proxy report (KINDL[®] for parents) [203]. The questionnaires consist of 24 corresponding items that are formed equally as either positive or negative statements about different facets of the child's life. Each statement addresses experiences over the past week and is rated on a five-point scale: ¹⁾ never, ²⁾ seldom, ³⁾ sometimes, ⁴⁾ often and ⁵⁾ always. The mean scores are calculated for each of the six subscales: physical well-being, emotional well-being, self-esteem, family, friends and school and for the total scale, and they were linearly transformed to a 0 to 100 scale, on which higher scores indicated a better QoL. All of the versions of the KINDL questionnaire were supplemented with a "disease module" consisting of a filtered question and six items about possible long-lasting illness or current hospital admission. The internal consistency and reliability were tested previously and compared with the original German version of the questionnaire [109]. The translated Norwegian version displayed better internal consistency (Cronbach's α) with an increasing age of the child. Among younger children (8-9 years old), Cronbach's α was low for the subscales of emotional well-being (0.52), friends (0.49) and school (0.47), while it varied between 0.62 to 0.68 on the other three subscales and was 0.83 on the total summary scale [109,202].

Seeking help

A simple report was designed for the present study to monitor parental use of seeking help on behalf of their children. The questions included the following: ¹⁾ type of contact (physiotherapy, child habilitation, child psychiatry services, special educational services and child welfare authorities), ²⁾ the age of the children when the contact was initiated and ³⁾ duration of contact. Parents provided response to this questionnaire when the children were nine years of age [Appendix, part V].

Birth, medical risk and family-related factors

Childbirth (weight, gestational age, sex, twin) and perinatal risk factors were collected before discharge. Risk factors included the clinical risk index for babies (CRIB), presence of bronchopulmonary dysplasia (BPD) and presence of several cerebral injuries [169,192]. Family-related variables were as follows: ¹⁾ mother’s age, ²⁾ mother’s and father’s years of education, ³⁾ mother’s and father’s annual income, ⁴⁾ number of siblings, ⁵⁾ parental marital status, ⁶⁾ housing conditions, ⁷⁾ occupational status, ⁸⁾ smoking habits and ⁹⁾ ethnicity. This information was reported in separate questionnaires before discharge from the hospital [Appendix, part VI].

Figure 5. Measurements in the thesis

Measurements	Pre-rando- mization	½ y	1 y	2 y	3 y	5 y	7 y	9 y
CBCL – behavior & emotional problems – parent report				x	x	x	x	x
TRF – behavior & emotional problems – teacher report							x	x
SDQ – prosocial behavior & peer problems – parent report						x		x
SDQ – preschool and school report						x		x
KINDLE - on quality of life								x
PSI – parenting stress	x	x	x	x	x	x	x	x
Socio-Economic Status	x		x	x	x	x	x	x
Help seeking								x

y = children’s age in years.

3.5 Follow-up procedures

For the purpose of the study, all of the participating children received the same medical, developmental, and psycho-social assessments at corrected ages of 6 months and 1, 2, 3, 5, 7 and 9 years. All of the assessors outside the families were blinded to the group allocation of the children.

Questionnaires were sent to the families approximately fourteen days before the follow-up sessions and collected at the time of follow-up by the coordinator. Parents received written reports concerning the motor and cognitive development of the child after every follow-up. These included recommendations about referrals to other services if relevant that were signed by the study director. In addition, all of the parents were allowed to call to the study coordinator or director throughout the duration of the study and request advice if needed. All of the results are reported as intention to treat, and one family that was randomized to the intervention but that did not receive the intervention was included in the intervention group. The original study was planned to report the outcomes of the children at two years of age. Because the study was prolonged several times, it was approved on three occasions by the regional committee for medical ethics and the Norwegian Data Inspectorate (1999, 2005 and 2010).

3.6 Statistical analysis

The analysis of the results was dominated by group comparisons. TISP is a clinical trial, and intervention outcomes have been explored by comparisons between PI and PC outcomes and by comparing each of the preterm groups with the TR group. These analyses were generated using linear mixed models (SPSS statistics, version 20) because of repeated measures and the clustering effects of twin pairs.

In the longitudinal linear mixed-model analyses, time was treated as a continuous variable. Predicted mean group differences in each measurement occasion with 95% confidence intervals were also computed. This analysis was also based on a longitudinal model, but in the case of group comparisons, time was used as a categorical variable [205]. By changing the reference time point in the analysis, predicted group differences on each occasion could be estimated. Dichotomized variables were analyzed using generalized linear mixed models (GLMM), which generated odds ratios that were adjusted for the clustering effects of twin pairs.

Longitudinal analyses of behavioral outcomes included reports both on CBCL/2-3 and CBCL/4-18. The questionnaires contain some differences, making the use of raw scores inappropriate. Following a request to the owner of these questionnaires, we received a conversion table that made it possible to convert the existing raw scores from the CBCL/2-3 to T-scores [48].

Effect sizes were assessed in different ways in relation to the different types of statistical analyses.

¹⁾ In the LMM models, they were computed as Pseudo R-square values and based on the square of the correlation between the observed and predicted values of the dependent variable [206]. After assessing the effect of the addition of variables in a linear mixed model analysis, the change in this pseudo R-square value was determined as an effect size measurement.

²⁾ To analyze the effect sizes (ES) of treatment group comparisons, Hedges' *g* were computed [207]. Hedge' *g* is frequently recommended when sample sizes are small and is estimated by use of group-sizes, mean values and SD's to express an effect size. Higher values of Hedges' *g* indicate larger differences between groups. ³⁾ The levels of parental agreement were analyzed by intraclass correlations (ICCs). Differences between the two independent ICC coefficients for the PI and PC groups were tested as described by Alsawalmeh and Feldt [208]. This is a test of the equality of two reliability coefficients after adjustment for the unequal length of the instruments that were employed in the reliability analysis [ibid].

4. Summary of results

Paper I

Landsem IP, Handegård BH, Ulvund SE, Tunby J, Kaaresen PI, Rønning JA. Does an early intervention influence behavioral development until age nine in children born prematurely? *Child Development* 2015; 86(4):1063–1079

The longitudinal development of behavioral problems among PI and PC children followed similar trajectories across childhood from 2 until 9 years of age.

- In 23 of 24 reports on main behavioral dimensions (internalizing and externalizing), lower mean scores were reported for the PI children than for the PC children [non-significant difference].
- Slightly more PC children (18%) than PI children (11.8%) scored within a clinical range on reported problems (CBCL Total problems score > 35) [non-significant differences]. Boys were more frequent reported with high scores [non-significant difference].
- The mean T-scores on CBCL, Total problems were relatively low for all groups, ages and respondents compared to other reports.
- Across groups, mothers reported more Total Problems than fathers. [non-significant difference].

More attention problems and less successful adaptation to school were reported for PC children compared with PI children at 7 and 9 years of age.

- More attention problems were reported by mothers (ES = 0.48), fathers (ES = 0.48) and teachers (ES = 0.48) at age 7 years and by teachers (ES = 0.43) at age 9 years. Teachers reported significantly more inattention, hyper-impulsivity and somatic problems for the PC group at 9 years of age.
- More problems with thoughts were reported by PC fathers (ES = 0.52) and PC teachers (ES = 0.40) at age 9 compared with reports on PI children.
- Less successful adaptation to primary school demands were reported for PC children compared with PI children. Significant differences between preterm groups comprised measures of;
 - social competence and school performance at age 7 and the overall competence score at age 9 as reported by mothers.

- school performance at age 7 and activities and school performance at age 9 as reported by fathers.
- school achievements and perception at age 7 and competencies such as school performance, achievement, appropriate behavior and perception at age 9 as reported by teachers.

PI teachers perceived that PI children showed better adaptation to challenges in daily life compared with reports from PC teachers regarding PC children [blinded reports].

- PC children were reported to address definite or severe problems by teachers more than twice as often as PI children (OR = 2.9).
- Teacher's report of difficulties correlated significantly with parent's reports of referral to child psychiatric services, and PC children were referred to child psychiatric services more than twice as often as PI children.
- PC fathers reported problems in peer relationships were more frequently the cause of difficulties compared with PI fathers.

Some features of longitudinal development of behavioral problems on sub-domains differed significantly between the two preterm groups:

- Father's reported PC children with a BW > 1000 grams to exhibit more worrisome development in terms of externalizing problems across childhood compared with PI fathers for the similar group of PI children.
- Mothers reported that PC boys were more withdrawn during the preschool years than PI boys, but at nine years of age, they reported similar levels; an opposite tendency was reported for PI and PC girls.
- The fathers of PC boys reported a steeper increase in anxious behavior from age 5 to 9 years compared with the fathers of PI boys.

Preterm groups were compared separately with results of the TR group.

- The more worrisome development of anxious behavioral problems was reported for the PC compared with the TR group from 5 to 9 years of age by fathers.

- Significantly more behavior problems were reported in the PC compared with the TR group across all ages.
- Behavioral problems in the PI and TR groups developed with similar trajectories across childhood. At nine years of age, parents and teachers reported similar levels of behavioral problems and competencies for these groups.

Agreement between the parent's reports for internalizing and externalizing behavior differed between the two preterm groups on two occasions.

- At 2 years of age, PI parents showed greater agreement than PC parents concerning the internalizing behavior of infants.
- At 9 years of age, PC parents showed greater agreement than PI parents concerning the externalizing and total problems of children.

Differences in intervention participation influenced PI fathers reports at age 9 as fathers who had participated less reported more externalizing problems ($p = 0.001$) characterized by more aggressive behavior.

Paper II

Landsem IP, Handegård BH, Tunby J, Ulvund SE, Rønning JA. Early intervention program reduces stress in parents of preterms during childhood, a randomized controlled trial. *Trials*, 2014, 15: 387.

Mothers in the PC group reported increasing levels of child-related stress until 5 years of age that was particularly affected by limited adaptability and moodiness of the PC children. PI and TR mothers reported significantly different, declining levels across childhood compared to PC mothers.

- PC mothers reported more total and child-related stress compared with PI mothers at all follow-ups and more parent-related stress during the pre-school period.
- PC fathers reported more child-related stress at 2, 3 and 5 years of age and more total stress at 5 years of age.
- PI fathers who had participated less in the interventions reported more stress when children were 3, 5 and 9-year-old.

Mothers and fathers in the PI group reported higher correlations concerning parenting stress compared with the couples in the PC group.

Parenting stress was reported at similar levels in the PI and the TR families throughout childhood. PC group parents reported significantly more parenting stress than TR parents at all ages.

Across all of the groups, child and parent-related stress occurred at relatively low levels compared with the findings of American studies but were similar to the results of Scandinavian studies.

Paper III

Landsem IP, Handegård BH, Ulvund SE, Kaaresen PI, Rønning PI. Early intervention influences positively Quality of Life as reported by prematurely born children at age nine and their parents; a randomized clinical trial. *Health and Quality of Life Outcomes*, 2015; 13:25.

PI children reported improved physical well-being (ES = 0.57) compared with PC children at nine years of age.

- For all other features of well-being, both PI and PC children reported statistically similar results to the TR children.
- The BW and severe neonatal morbidity of the children were significant negatively correlated with reports of physical well-being.

PI parents reported better emotional well-being (ES = 0.34) and better well-being in school (ES = 0.54) than PC parents.

Agreement between children and parents was significantly lower in the PI group compared with the PC group. The pattern of agreement in the PI group was similar to that in the TR group.

PC parents reported significantly lower well-being among their children in all assessed features compared with TR parents. PI parents reported well-being at similar levels as TR parents in all assessed features except a lower score on the dimension of self-esteem.

5. Discussion

A core hypothesis of this study was that MITP-M would sensitize the parents to their infant's cues, which would lead to more adjusted and dynamic parent-infant interactions and thereby enhance the infant development. The main findings presented in this thesis support this hypothesis. The MITP-M intervention seems effective in; improving the long-term socio-emotional and behavioral development among preterm born children with BW < 2000g; reducing long-term parenting stress in parents of preterms and support self- and parent-reported QoL at children's age of nine.

Longitudinal behavior development was not affected by the intervention as the PI- and the PC children in broad outline with similar behaviorally trajectories across childhood. Even though, at seven and nine years of age the PI children were reported with less attentional problems, higher social competencies and better academic school performances than the PC children. Behavioral problems and competencies in the PI group were at similar levels as reported on the TR group from mothers, fathers and teachers at children's age of nine. The PC group continued to be reported with significantly more problems and less competencies than the TR group [paper I / 211].

PI mothers reported less child- and parent-related stress than PC mothers at all follow-ups across childhood and they reported a higher agreement with fathers than in the PC group. More child-related stress reported by the PC mothers made up a qualitatively different trajectory compared with the gradually decreasing trajectories reported from PI mothers and fathers in both groups. Both parents in the PI group reported parenting stress similar to the TR group at all follow-ups, whereas differences between the PC and the TR group increased with age throughout childhood [paper II / 179].

In the third study children and parents in the PI group reported better well-being on some measures of QoL at children's age of nine [paper III / 236].

In the following sections, the results are discussed in relation to possible influences of the MITP-M, relatedness to other results from TISP and recent research in the field of early child development and outcomes reported on preterm children.

5.1 Children's emotional behavior development

Behavioral problems of children have been reported repeatedly by TISP [209,210]. In the current study these reports were extended to include longitudinal analyses of behavior from children's age of 2 until 9 years and cross-sectional reports of problems and competencies at the ages seven and nine [211]. The trajectories reported were consistent with behavioral development that has been formerly reported for normative and at-risk samples. Even though, the overall level of problems reported were low compared to other studies [45,212]. Elgen et al. [36] reported a 40% increase in behavioral problems among children born preterm compared with full-terms at their age of eleven. Reports of behavioral problems on preterm children in TISP seem to be at a lower level. Non-significantly more PC- than PI children had high problem scores, 18% in the PC group versus 11.8% in the PI group.

Fewer problems in TISP compared to previous studies may be a positive effect of the follow-up program. Parents received written reports regarding their child's level of performance at every follow-up and could contact the study coordinator or director for talks whenever needed. This result address the importance of a well designed follow-up program for children born preterm.

Longitudinal behavior development in the PI and the PC group followed similar trajectories as in the TR group with one exception. PC fathers reported a more worrisome development of anxious behavior than TR children [211]. This corresponded with PC father's reports of a steeper increase of anxious behavior among PC boys than PI father's reports on PI boys from children's age of five until nine [ibid]. Very preterm children have previously been identified with increased risk of emotional problems in prepubescent years (6 to 13 years) and more emotional problems were especially reported on children born preterm, with male gender and those living with parents from low SES backgrounds [213]. A similar finding was reported from another large study were ELBW children were reported with a 4-fold increased odds of having emotional problems compared with term controls at six years of age [214].

This studies of emotional problems may sheds light on the different trajectories of anxious behavior reported by PC fathers compared to PI and TR fathers up to the age of nine years in TISP [209]. It is possible that the MITP-M have promoted a long-term protective effect against anxious problems in the PI group. Even though, significant differences were related to reports of more anxious problems

in relatively few PC boys in the current study and further sub-analysis could not be done because of lack of statistical power. PC parents reported that their child had been referred to specialized child-psychiatric services more than twice as often as PI parents on PI children. This information supports the hypothesis that the MITP-M may have long-term protective effects concerning children's mental health.

Children in the PI- and the PC group develop similarly to the term control children in all other aspects of longitudinal behavior development reported from TISP. However, although they develop similarly to TR children, their development may be viewed as somewhat delayed and they may need more support from their family and the society as reported in previous studies [215-219]. The variation between individuals in TISP was reflected by high standard deviations in reports of behavioral problems throughout childhood. This result confirmed previous findings reported by Gilliom & Shaw that showed significant variability in individual-level trajectories of behavior [212]. Although PI children were reported to have more problems than term controls during earlier years, they seem to reach an age-appropriate/normative level of maturation, adaptability and competence at nine years of age.

It is very promising that the MITP-M seemed to support an accelerated developmental catch-up among PI children. In agreement with the transactional understanding underlying this study and previous publications, this finding may be partly due to a decrease in parental stress [133,145, 220,221], improved parent-child interactional relationship [220] and a reduced parental perception of the child's vulnerability [219,223]. This is supposed to influence the further development of the children up to puberty and adolescence via more positive transactions and effects on the child's own, parent's- and peer's perceptions.

Parent's reports of internalizing behavior

The extent of internalizing problems reported in TISP followed the tendency reported as normative trajectories of problems by Bongers et al. [45]. The number of problems increased slightly during childhood even though the reports from different respondents displayed slightly different patterns.

The trajectory of children's behavior in terms of internalizing problems throughout childhood is reported to be highly influenced by maternal psychopathology and stress [43]. Further investigations may uncover if similar associations appear in the present study and if the intervention may have reduced, not only parenting stress but also psychopathology in the PI group. The slightly larger amount of internalizing problems reported by PC mothers compared with other groups of respondents at two years may reflect a view of their children as more withdrawn (e.g., shy, fearful, sad), which coincides with them reporting more child-related stress, especially in relation to the adaptability and mood of the children [179]. Preterm children have been described to have a reduced capacity to express positive emotions (smiles, eye contact, vocalizations) [126,224] and more dysregulated behavior during toddlerhood [80, 225,226]. In addition, 21% of VLBW children were reported to have dysregulated behavior at 2 years of age in a recent Australian study [56], and VLBW children demonstrated significantly more internalizing problems and lower socio-emotional competence at 2 years of age compared with full-terms.

Deficits in the expressivity of children born preterm are thought to affect parent-child interactions [227,228]. Parents may feel less excited by their child, which may also affect their interpretation of the child's signs and needs. Earlier reports of more intrusive parental behavior in preterm parent-child dyads [138] may be interpreted as the attempts of parents to elicit smiles and responses when the child required less stimulation, resulting in more internalizing behavior. This highlights the importance of early parental sensitization to the expressions of every unique child [42,135,229].

Preterm infants have also been described to be more dependent on maternal scaffolding behavior to establish successful emotion-regulation compared with their peers [230]. PC parents may experience difficulties in detecting this prematurity-related behavior as early as PI parents who are supported by insights from the intervention and who as early as in two-year-old children reported non-significant less internalizing behavior compared with PC parents [211]. This finding may also correspond to reports of improved maternal adaptation to the temperamental style of the child at an age of one year, as reported by Olafsen et al. [232]. Later in childhood PC parent-child dyads

appear to be less disturbed by internalizing behavior. This may indicate an establishment of improved parent-child adaptation in children aged approximately three years for PC mothers and aged five years for PC fathers.

More anxious/depressed behavior and somatic complaints have repeatedly been reported in girls than boys from the age that they enter primary school [45]. In TISP, two features of internalizing behavior made up statistically different interactional trajectories from age 2 until 9 years. They were related to preterm groups and gender: a) potentially more anxious behavior in boys in the PC group compared with the PI group from the time of school entry, and b) a shift in withdrawal problems during the first years of school compared with the preschool years. According to withdrawal behavior, the PI and the PC boys increased in similarity with increasing age, while an opposite tendency was reported for girls up to the age of nine because PC girls were reported to be slightly more withdrawn than PI girls at age nine in comparison to previous years. The impact of these results is not clear because an increased number of problems were reported for relative few children. The differences may be transient or they may be precursors of a subsequent increased risk of psychopathology in the PC group similar to the results reported on VLBW children by Treyvaud et al. [79].

Teachers reported internalizing problems in the same order as parents, which were highest in the PC group and lowest in the TR group. The PI group placed itself in the middle of these two groups. Internalizing behavior may not be as visible for teachers as for parents, and it has been suggested previously that mothers are better observers of this type of behavior than teachers [65]. However, the teachers were blinded to children's group belonging, and therefore, their reports may be more trustworthy.

Figure 6. The mean reports of internalizing behavioral problems across childhood in all three study groups from all respondents.




Figure 6. Mean T-scores for CBCL and TRF of internalizing problems reported by mothers, fathers and teachers in each study group at all follow-ups.

Parent’s reports of externalizing behavior

Longitudinal investigations of externalizing behavior revealed similar patterns of decreasing problems in all groups, which is consistent with reports on normal populations [45,233]. The significant differences in behavioral outcomes reported by Nordhov [210] at age five years disappeared in our analysis when controlling for repeated measures. However, a similar pattern to that observed for internalizing behavioral problems was evident as the PI group was consistently reported with fewer externalizing problems than the PC group by mothers, fathers and teachers [211].

Externalizing behavior, as reported in all study groups and for all groups of respondents are presented in Figure 7. TR mothers reported decreasing problems in children as early as two years of

age. PI mothers reported a corresponding drop in the average number of problems after three years of age, while PC mothers reported a decreasing trajectory after five years of age. The delayed decrease in problems may first illustrate the parenting challenges that are faced by the mothers of preterm children compared with full-term mothers. Secondly may the intervention have expedited a more successful adaptation between the mothers and children in the PI group compared with the PC group.




Figure 7. Mean T-scores for CBCL and TRF of externalizing problems reported by mothers, fathers and teachers in each study group at all follow-ups.

A surprising finding was the differences in the father’s reports of externalizing behavior across childhood for the heavier group of preterm children. PC children with a BW > 1000 g had a significantly more worrisome trajectory concerning externalizing problems than the same sub-group in the PI group. Fathers of the heavier PC children reported externalizing behavior at stable and high levels from age 2 to 9 years while fathers of the same PI sub-group reported decreasing problems similar to normal populations from age five until 9 years.

This difference may be an effect of father's restricted access to the NICU in our hospital at the time when the study took place. Firstly, fathers of heavier children were generally less present in the hospital compared with fathers of smaller and medically sicker newborns. Fathers may thus have received less knowledge about how to interact with their infants in a developmentally supportive way. Secondly, mothers of heavier preterm children may have received less guidance regarding developmentally appropriate care compared with mothers of the smallest preterms. Mothers (and some fathers) with smaller preterms were hospitalized with their children for a longer period and had thus more possibilities to learn about their infant's signs, needs and behavior from professionals in the NICU irrespective of the intervention program. PI parents of the heavier preterm children may have established a similar enhanced understanding of their infant's expressions and needs during the MITP-m sessions while PC parents of the heavier preterm children may have been more dependent on their own personal resources.

This proposal of a possible reason of the significant differences in PI and PC father's reports of externalizing behavior corresponds with the research of Spinelli et al.[234]. They reported on parental stress trajectories in premature mother-infant dyads from the age of four months to three years. One of the groups of mothers who reported high levels of stress at 4 month of age consisted of mothers who had a short hospitalization with their preterm child [ibid].

5.2 Family climate and parental agreement

Parenting behavior or family functioning has not explicitly been studied in TISP. However, an understanding of the functioning of family systems has become possible; first, through reports of parent's perceptions of children's temperament during the first year [231,232], through reports on parent's attitudes toward child-rearing across toddlerhood [235] and lastly, through the reporting of several measures of parental agreement in the papers included in this thesis concerning behavioral problems, parenting stress and perceptions of children's quality of life at nine years of age [179,211, 236].

Harsh parenting involving higher parenting control has been, together with a demanding child temperament, interparental conflicts [237] and low paternal caregiving capacities [238], described as a precursor of externalizing behavior throughout childhood [233,239]. Several of these factors have been addressed in the studies mentioned above and are suggested to be associated with the reports of slightly more behavioral problems and challenged family climate in the PC group compared with the PI group. Although no differences were uncovered between the reactivity of PI and PC children during the first year, Olafsen et al. reported that PI mothers were more affected by their children's temperamental style at 6 months of age [231]. At an age of one year, this phenomenon had disappeared, which may have occurred because the PI mothers had adjusted their parenting behavior according to the characteristics of the children. However, a strong association between maternal stress and negative reactivity was reported by PC mothers at one year of age [232]. These early differences may be one underlying reason for the consistently small and non-significant differences in reported problems between the PI and PC groups at later follow-ups [211].

Nordhov et al. reported differences in child-rearing attitudes across toddlerhood between the PI and PC group [235]. Mothers who had received the intervention reported more nurturing child-rearing attitudes toward their children at one and two years of age, and the same tendency was detected at 3 years of age. The differences reported by mothers concerning the temperaments of their children and later child-rearing attitudes may reflect some differences in parent-child transactions and family climates between the PI and PC group.

These differences may also be discussed in relation to parental reports of children's behavior and parenting stress in the current thesis. PI parents reported a higher agreement than PC parents in their assessment of internalizing behavior at two years of age [211]. This observation may indicate that both mothers and fathers were able to recognize early signs of internalizing behavior among their children after the intervention, while PC parents obtained less agreement in the family at this early stage. PC parents may have lacked important insights and possibly misinterpreted behavior such as sadness, shyness, lack of affections, and avoidance of eye contact.

The interpretation of the results above may also be supported by higher parental agreement concerning child- and parent-related stress [179]. In children aged two years, differences in parental agreement covered both aspects of parenting stress, while at age three the difference was isolated to parent-related stress. PC fathers reported fewer behavioral problems at all ages and less parenting stress throughout childhood compared with PC mothers, which is with the pattern observed in the PI and TR group. However, the differences in parental agreement between the PC and PI group at age two and three years may indicate that PC fathers were less involved in the caretaking of their child, both before and after discharge from hospital, compared with the PI fathers. They may not have been aware of the challenges faced by the PC mothers regarding children younger than three years of age, during which PC parental agreement concerning child-related stress was equivalent to that in the PI group.

PI mothers and PI fathers may have been sensitized to their infant's behavior during infancy as a result of the intervention, and even though some fathers participated less in the intervention program, they may have been more easily inspired by the mothers and caught up with them more successfully compared with the PC fathers. Agreement between parents in the TR group has not been questioned, but TR fathers appeared to be generally less aware of behavioral problems of any type compared with their partners.

5.3 Socio-emotional problems, social competence and mental health

PI children were reported by their mothers to be more socially available at the age of one year [232] and with less social problems than PC children at five years of age [210]. This tendency continued in children aged 7 and 9 years, even significant differences reported by teachers at age nine disappeared when adjusting for the presence of twins in the sample. Similarly, PI mothers reported significantly fewer social problems at age nine years compared with PC mothers after the exclusion of one extreme outlier. In general, PC fathers did not report more social problems than PI fathers. However, among those fathers who reported that his child experienced difficulties in daily life, PC fathers identified this condition to be related more often to difficulties associated with peer relationships. Finally, PI mothers rated their children with higher social competence compared with PC mothers at seven years of age, and a similar non-significant difference was reported by both PI mothers and PI fathers at nine years of age.

Developmental goals concerning social functioning in early school years are described as the ability to form dyadic friendships with peers and the diminution of physical aggression [37], handling of frustration and self-regulating [242]. The intervention may have strengthened the ability of the PI children to reach such goals at an age-appropriate time. This result makes sense because children born preterm have been reported to need a longer duration than full-terms to achieve similar levels of performance [215,242].

Socio-emotional problems are frequently identified as precursors of later complex school- or psycho-social problems. VLBW children without major handicaps were reported to display more socially inappropriate behavior from age 6 to 8 years compared to full-terms in a longitudinal study from the 1990s [66]. Recently, it was shown that 25% of a sample of ELBW children was identified with socio-emotional delays at two years of age [242]. Increased social problems among VLBW children have also been described to be predictive of later lower social competence and poorer school performance at eleven years of age [241]. In another study, 33% of the problems reported by teachers in elementary schools were presaged in parental reports for children aged 12 to 36 months [44]. Thus, socio-emotional problems may start early and be relatively stable throughout childhood.

Later studies continued to report a greater number of socio-emotional problems in preterm populations compared with full-terms [243]. They discovered socio-emotional problems at age five years to be predicted by emotional problems at two years of age [ibid]. Next, socio-emotional problems at five years of age and children with a higher social risk at seven years of age were more likely to meet the criteria for a psychiatric disorder at an age of 7 years [79]. Compared with terms, preterm children had a three times increased odds of meeting any psychiatric diagnosis at 7 years of age (24% of preterm children)[ibid]. The most common diagnoses were anxiety, attention deficit/hyperactivity and autism spectrum disorders [244]. This finding is consistent with two Norwegian studies that investigated preterm populations. In a sample of VLBW children from the central region of Norway, 46% of children were found to have psychiatric problems at the age of 14 [245]. Among the children, 25% met the criteria for a psychiatric disorder [99], and the prevalence of psychiatric morbidity displayed a similar level at the age of 20 years [245]. In the western region of Norway, Elgen et al. reported that 27% of a VLBW sample exhibited a psychiatric disorder at the age of 11 years [36,98]. These results can be seen in relation to those of a Nordic study of impaired psycho-social functioning/problems that estimated the normative prevalence to be approximately 10% [246]. In TISP, parents reported children in the PC group as more than twice as often referred to specialized child-psychiatric services at children's age of nine compared to reports on PI children. This information was highly correlated with teachers reports of more difficulties in everyday life in the PC group compared with the PI group. The parental reports included not information about severity of symptoms or diagnosis but the findings above may be an indication of more severe socio-emotional problems in the PC group than the PI group in middle childhood.

How may the findings of higher social competencies among PI children compared with PC children be influenced by the early MITP-M intervention? One possible explanation addresses a more successful down-regulation of infant stress/distress from infancy and across preschool years in the PI group. PI parents may have learned how they can adjust their interactions and demands to the individual capacities of the child [80,239,247]. A reduction of stress might correspond to better cortisol regulation [146,220,248], higher performance in the anterior attentional system and improved control [249]. Another plausible explanation is the bidirectional relationship between the parent and child and the influence of less parenting stress in PI families [32,221,228]. Parenting stress

has been described to predict later child coping competence, and child coping competence predicted later parenting stress [222].

5.4 Attention problems and perceived academic performance

Preterm children aged seven and nine years exhibited two main differences; teachers and parents reported fewer attention-related problems and improved school achievement in PI compared with PC children [223]. PI children adapted more easily to school demands and new environments compared to PC children. A similar pattern was reflected by parental reports of stress at seven and nine years of age. PI parents reported less stress than PC parents, with significant differences observed between mothers of seven and nine-year-old children [ibid.].

There are several possible explanations why the intervention may have promoted better attentional capacities among preterm children in the PI group compared with the PC group. The maturation and development of children's attention throughout childhood may have been supported by a more successful down-regulation of initial neonatal distress and a subsequent consistent reduction of parenting stress in PI families [179]. In infancy, children may have greater incentives to impart robustness into the processing of the orienting system and subsequently the alerting system, positively impacting the ability of children to focus attention. Attention is a basic condition for the ability of a child to orient. Early orienting attentional capacity at 5 months of age has been described as a precursor, a "building block" for later attention that is needed for learning and performance until four years of age [250]. Finally, it is plausible to suggest that intervention-generated, well-tuned parent-child interactions in the PI group may have fostered more scaffolding behavior among PI parents. This could have supported the development of the effortful control system in children and more goal-directed attention in toddlerhood. During early childhood, executive functions are linked both with school readiness, academic performance and early-onset disorders (e.g., ADHD, inattention) [250,251].

The heightened risk for learning and attention problems among children born preterm has been well documented [83,84,117,252-254]. Groen-Blockhuis et al. reported strong evidence for a causal relationship between a lower BW and later attention problems [78]. Furthermore, children born preterm have been reported to have a more than doubled risk of developing ADHD disorders compared with full-term children and a 50% increased risk of requiring specialized pedagogical services [26]. In the study by Indredavik et al. [99], VLBW children reported attention problems that were similar to their full-term peers at 14 years of age, while both mothers and fathers of preterms,

on average, reported a three-fold increase in attention problems compared with parents of full-terms. As many as 25% of the VLBW children were identified with attention problems [ibid.]. This finding has later been associated with white matter abnormalities at an age of 15 years, and the authors suggested that a higher order of cognitive function, such as attention, depends on the intact communication between several cortical areas [256].

Attention is both a complex growing competency in the developing children and a prerequisite for developing other competencies such as social functioning and school achievements. However, the assessment of attention is poorly integrated in standardized cognitive measures. This finding may be a reason why the cognitive results reported in TISP at children's age of nine did not reveal any differences between the PI and PC group in contrast to the differences reported previously at children's five years of age [257,258]. Hauglann et al. reported similar intelligence coefficients (IQ) in the PI and PC group at the age of nine years and with marginal differences at seven years of age. They did not compare outcomes of IQ with the term control group. A Finnish follow-up study from the beginning of this millennium reported better cognitive development among VLBW children than reported in earlier publications but still they differed significantly from full-term controls already at the age of two [260]. It has also been noted that IQ scores did not represent a sufficient measure of the mental resources and capabilities of the children [259]. Aylward has stated that the sole use of IQ scores may mask the complex profiles of children with multiple areas of weakness that may negatively affect later functioning in school. This finding was supported by a study that identified problems in features such as attention, executive functioning and memory among preterm children who were assessed as normal in terms of IQ [25].

Proximal environmental factors, such as difficult parent-child interactions, and distal environmental factors, such as reduced social-economic status (SES), in families are supposed to influence the school performance of children [61,165,259]. This is reflected in the results of the current study, the intervention may to some degree have made the PI families more resilient. Still, at children's age of nine, a greater amount of stress related to difficult interactions with children was reported by PC parents compared with PI and TR parents [179]. We suppose that the patterns of interactions within each family may have been well consoled after all those years. Both parents and children may have formed fixed pictures regarding whom and how the other is concerned. Thus, more stress in the PC

families may be viewed as a hidden disturbance, a proximal and relational environmental factor that influences the overall performance of these children.

PI children were viewed as comparable with full-term peers by their teachers in terms of school and academic performance. However, a more intense use of pedagogical specialized services may have been necessary to reach this level. A total of 34% of the PC children, 28% of the PI children and only 3% of the TR children had received additional school services during the preschool/school years, according to parent's reports at nine years of age.

A slight increase in parenting stress among PI mothers and fathers of children aged seven years may be a sign of the elevated parental support and involvement needed in this group compared with the TR group. The catch-up within the PI group may have been supported by both additional school services and an increase in parental investments.

5.5 Parenting stress and possible influences on children's behavioral problems

Long-lasting stress that is present early in development has been described with detrimental effects on the well-being of parents, children and the relationship between them [199]. In line with previous reports from TISP [192, 209], more parenting stress were reported in the PC group compared with the PI group at all follow-ups until children's age of nine [paper II]. Reports of stress have been analyzed longitudinally from infancy up to the age of nine years and especially PC mothers seem affected by increasingly more child related stress across childhood. The same aspects of parenting stress created the biggest differences between the preterm groups throughout childhood. This was primarily reports of more distractibility, more moodiness and maternal perception of less competence and attachment.

The sub-dimensions mentioned above cover aspects of parenting stress associated with difficulties in parent-infant interactions [133] which continued to be reported as significantly more frequent by PC mothers compared with PI mothers at children's age of nine [paper II]. This corresponds with the study by Gray et al. where the largest difference in parenting stress between a preterm- and a term group was related to difficulties in parent-child interactions [139]. They did not discover differences between groups in early infancy (4 months) [261], while significantly more maternal stress was reported in the preterm group at age one [139]. In that study, symptoms of maternal depression and infant temperament were detected as independent risk factors at both follow-ups (4 & 12 months) [139,261]. Stress related to dysfunctional parent-child interactions has also been reported to show specificity to child internalizing problems in a study of referred children between the ages of 5 and 17 years [262]. The association was independent of psychopathology among the parents. Mechanisms of causality have not yet been investigated in TISP. It seem very important to continue the investigations and explore to what degree differences in child outcomes between preterm groups are caused by enhanced parent-child interactional relationships from infancy until middle school age.

Child-related stress

PC mothers reported more child-related stress across childhood compared with PI mothers, and this made up a different trajectory in children up to the age of seven years compared with reports from other groups and PC fathers [179]. Child-related stress is thought to reflect how parents perceive the attributes of their child [133] and how much they struggle with their care-taking responsibilities in light of the child's illness, behavioral problems or emotional disturbances [132]. As we all become stressed when faced with big challenges in life (e.g. feeling our life is threatened, our children are in danger), child-related stress may be perceived as an answer to a challenging situation, and it is a rational response to a demanding situation [132,133].

PC mothers reported also the most internalizing, externalizing, attention and social behavioral problems among all of the respondents across all of the assessments [223]. Together with their reports of the most parenting stress among all of the respondents, these results is consistent with the findings of a Canadian study in which parenting stress caused by high levels of distractibility in the preschool years predicted behavioral problems at seven years of age in a sample of full-term and moderately preterm children [32].

The areas that created differences in child-related stress between PC and PI mothers have been previously reported as challenging for children born preterm; these children are less likely than full-terms to instigate interactions, are less attentive, show less positive emotions and are sometimes regarded as less attractive social partners [154,263]. These features were addressed in MITP-M in which the parents were supported to detect the social identity of their child and to be emotionally affected by the early signs of responsivity in the behavior of their infant [264]. This may have promoted feelings of love, acceptance and positive attitudes in the PI parents.

In addition, MITP-M provided information regarding how parents could effectively co-regulate their infant and modify their actions to facilitate a more effective adaption of the child to life in the home environment. Preterm children experience high levels of stress during their stay in the NICU [265-267]. Their altered behavior and reactions to stimuli may be viewed as consequences of their adaptation to non-appropriate environments. Parents of preterms must help their infants re-adapt to more nurturing environments and thus help them to become less distressed. This phenomenon has been confirmed in recent research showing that positive parenting behavior and reduced parenting stress appear to ameliorate the negative effects of early pain among very preterm

children [188]. These parental factors have also been associated with less internalizing behavior in children aged 18 months [53]. Another study described associations between neonatal pain and stress and an altered HPA axis functionality up to 7 years of age in children who were born very preterm [267]. These studies underscore that children born preterm require successful parental co-regulation throughout childhood.

Parent-related stress

MITP-M seems to have subdued the pressure of parent-related stress on children's mothers. PI mothers reported significantly less parent-related stress from infancy until 5 years of age and again borderline less stress compared to PC mothers at nine years. In contrast to child-related stress, parent-related stress may be more subtle because it reflects the perception of the parents regarding their own parenting capabilities and the resources available to meet these demands [132]. Differences between the preterm groups were most pronounced in the subscales Attachment and Competence and to some degree in the scale focusing on marital satisfaction (PSI-Spouse).

The PSI subscale of Attachment focuses on parent's perception of emotional closeness to their child, which is an important factor in the fostering of their intrinsic motivation of being parents [133]. The disruptive experience of becoming parents weeks and month before planning, the limited possibilities to stay intimately close (skin to skin), the unfamiliar environments in the NICU and the continuous presence of strangers are all risk factors of elevated attachment-related stress among these parents [5,40,227]. In one study, only 20% of mothers of preterm children were described as having a secure attachment to their children at 6 months of age compared with 53% of mothers of full-terms [227]. However, a relatively recent Finnish study did not report greater attachment difficulties in preterm mother-child dyads compared with dyads of full-term newborns [268]. Attachment difficulties may also be difficult to observe. Bienfait et al. observed low correlations between the reports of mothers and NICU nurses regarding mother-child bonding difficulties after two days in the NICU [269]. She concluded that mothers must have the possibility of expressing their feelings because their need for help may not be easy to observe by others. In TISP, this need was especially addressed by the first session added to the original MITP [169].

The powerful experience of being emotionally touched by the meeting with one's own newborn baby is an important story - rewritten in prose literature, in philosophy and in writings about becoming a parent [132,270,271]. These sometimes overwhelming passionate feelings may be viewed as a nature-given gift to new parents, supported by a biological-endocrine system to insure the caretaking of newborn infants [171-174]. The bio-behavioral synchrony between attachment partners, who describe connections between behavior and physiological systems, represents a powerful new knowledge that has not yet been fully investigated [173]. Nevertheless, this understanding has partly grown from studies of preterm infants for whom the quality of synchrony

may be reduced due to both early separation and difficult interpretation of the behavior of the child. Strong associations have been reported between physiological stability (heart rate variability and sleep-awake organization) in the third trimester [41], parent-child interactive synchrony later in infancy [40] and the ability of the child to self-regulate him or herself during the preschool years [41,80].

In some way, the MITP-M intervention may have restored or strengthened parental attachment perceptions and thereby created significantly fewer attachment-related stresses in the PI group compared with the PC group. After the interventions, the PI parents may have been able to “re-interpret” the behavior of the infant so that expressions of emotional instability became a “call for support” or a changeable mood of the infant “a call for paucity” in interactions and activities.

Perceptions of competence were the second parent-related dimension of stress making up differences between the PI and PC group throughout childhood. The MITP-M is described as a sensitizing intervention, but many teaching elements are included in this intervention. Parents must be aware of and recognize the type of signs, behavior, and expressions that may guide the daily caretaking of their child. Parenting Self-Efficacy (PSE) is thought to be a central element in the beliefs and expectation of parents regarding their ability to parent successfully [272,273]. Parenting stress is strongly associated with PSE [274], and PSE may be viewed as an important coping mechanism in the process characterizing the development of stress [276,276]. PSE is commonly viewed as a strengthening factor in parenting competence, but studies of preterm children have revealed that high levels of PSE without knowledge of child development were inversely associated with outcomes. The mothers with high PSE but little developmental knowledge were the least sensitive in interactions with their children [277]. Thus, a naive sense of self-efficacy may be potentially negative if parents are responsible for a preterm born child who displays unusual reactions to overstimulation or inappropriate handling. Low levels of self-efficacy may not be more common among parents of very preterm infants compared with parents of more mature children [278]. However, these researchers found that PSE regarding parenting tasks mediated the relationship between psychological symptoms and self-perceived parental competence. In TISP, there is limited information about the self-efficacy of the parents, but it seems relevant to mention these relationships as one possible mechanism underlying the differences reported for parent-related stress (paper II).

5.6 Influences on children's quality of life

In children aged nine years, TISP generated the first result in which children themselves reported outcomes independently of their parents [236]. Both PI children and their parents reported a somewhat higher quality of life, on average, than the PC families, and we suggest that the MITP-M can have long-lasting positive effects on well-being in families rearing preterm born children. Group differences became evident in different aspects of QoL in the children's and parent's reports, namely as bodily sensations in children and as a perception of emotional well-being and well-being in school among parents.

The results have already been discussed thoroughly in relation to comparable research [279-285] as well as previous and current reports from TISP in paper III [236]. Thus, a brief summary of the main finding will be presented herein. The PI children reported a higher level of bodily well-being than the PC group, and they also rated their bodily well-being as non-significantly higher than that of the TR children. Although a low BW and neonatal illness were negatively associated with QoL, neither these nor other birth, medical or socio-demographic factors explained the significant difference between the PI and PC group. It may be questioned whether the improved bodily well-being in the PI group was caused by better, early parent-child emotional co-regulation as reported by Treyvaud et al. [244] and as such created a more nourishing family climate with less stress in the PI families [179]. This finding may correspond with the strong associations detected between children's report of bodily wellbeing at nine years of age, and mother's, father's and teacher's reports of behavioral problems at seven years of age [236].

PI parents perceived their children to have a higher QoL than the PC group in the dimensions of emotional well-being and the ability of the child to thrive in school. Corresponding with other studies [283,286], a strong association was found between parents' proxy reports of emotional well-being and maternal reports of stress. Associations were evident at all ages, and the strongest association was observed between parental reports of emotional well-being and maternal stress reported at seven years of age.

PI parents also rated their children as enjoying a significantly higher school-related QoL, in line with the CBCL and TRF results [paper I & III]. This difference seemed to mostly affect boys but was highly related to attention problems reported by teachers at nine years of age [236]. The congruence

between reports of attention problems and QoL is not surprising. Being able to stay focused and assimilate messages are essential skills for all children, enabling them to experience well-being, social belonging and learning in school.

Figure 8 shows the means for different subscales and respondents.




Figure 8. Mean total and dimensionally separated QoL reported by children and parents in each study group at nine years of age.

In general, both children and parents rated QoL in the children as relatively high. PC children reported lower bodily well-being than PI children but the same level as TR children across all dimensions.

However, PC parents rated their children as having a lower QoL in all dimensions compared with TR parents, while PI parents reported a QoL similar to the reports provided by parents of full-terms on all scales, excluding slightly lower self-esteem. As repeatedly stated, minor difficulties related to prematurity may first become visible when children enter the school system [57,83]. Parents in both preterm groups may face greater challenges in this phase, as reflected by reports of somewhat greater parenting stress [paper II] and withdrawn behavior of the child [paper I]. This may also affect the parent’s experience of their children’s self-esteem.

6. Strengths and limitations

TISP has been acknowledged for its strong design [287] and repeated continuation of the study, permitting long-term follow up into middle childhood. Although many of the reported outcomes are influenced by several mapped and unmapped factors, the randomization process revealed a high level of equality between the PI and PC group.

Next, the high participation and low drop-out rates of the families in all three study groups has been a success factor. This may be a result of the participating families, ^{a)} a loyalty and idealistic belief in the value of the study, ^{b)} perceived personal benefits of the follow-up program, ^{c)} reports about one's own child development and, especially, ^{d)} easy available communication with and advice from the study coordinator, who met them at all sessions from the time of study recruitment until the children were nine years of age. Another strength that must be mentioned is the use of multiple informants which is especially recommended when assessing children's affective symptomatology [288].

Subsequently, the collection, punching and scientific processing of the data were conducted by the same small team (study coordinator and statistician) across all years to form a qualitative guarantee of the data material.

However, some weaknesses must be mentioned. Research reported in this thesis was solely based on data from questionnaires. Information about children's behavior, QoL or parenting stress was not assessed more directly. A main aim of the study was to compare outcomes between groups, for which the use of validated and well-designed questionnaires seemed appropriate.

In most studies that continue for more than a decade, such as TISP, new knowledge may inform the research field about factors that should be monitored because they may influence children in each group differently. In TISP it had been beneficial if exposure to painful events and the use of skin-to-skin contact (e.g. kangaroo care) with parents had been registered for all study participants during the NICU stay. This information is not available. The final years demonstrated that exposure to neonatal pain has a substantial negative impact on the cortisol regulation and behavior at school of children [289], while skin-to-skin contact with parents has been reported to support long-term developmental outcomes [267,290].

7. Clinical implications/ implications for later research

In the last 9 years, TISP has reported promising results for children born preterm as well as their families [192,209,210,234,258,229,230,257,291,292]. Although the MITP is more than 30 years old, the modified version tested in TISP (MITP-M) verified that children born preterm may catch-up with their peers born full-term in the middle of childhood. This was provided by parents who obtained the appropriate support in a phase that was important for the parent-child relationships. These results are consistent with the latest theories introduced by Feldman who suggested that early periods of sensitivity can be determined for the development of children and introduced as parent-child synchrony as a core mechanism for the early regulation and coordination of biology and behavior during social contact [173].

Although several possible reasons for the effects of the MITP-M are mentioned in this thesis, they must be tested scientifically. It will be important to investigate whether parenting stress is a precursor of behavioral problems in children or whether the behavior of children is a more primary trigger of a relationship between stress and behavior, which has been suggested to be influenced bi-directionally [1,68,122,234,294]. We have not yet analyzed to what degree early externalizing behavior is followed by later internalizing behavior in our sample, but this is a relevant topic because previous research has suggested that disruptive young children are likely to experience coexisting internalizing difficulties or difficulties that appear during later childhood [212].

The success and quality of parental bonding to the newborn child lays a foundation for the attachment quality of the child [294-296]. All children develop close emotional bonds to those who take care of them, but not all bonds (attachments) provide equal amounts of security [1,297,298]. Sherman et al. did not find associations between negative reactivity and attachment classifications in children aged 5 months, but at the age of one year, those children identified with an insecure and ambivalent attachment style were most reactive [299]. The attachment security of children was not exclusively investigated in the present study, but higher levels of maternal and paternal attachment-related stress were reported in the PC group from the age of 6 months and throughout childhood. This finding may indicate that the intervention strengthened parental attachment and subsequently the attachment quality of the children in the PI group. Fearon & Belsky found significant associations between early attachment quality and externalizing behavioral problems in

middle childhood, especially among children living with additional environmental risk and in boys [300]. Attachment quality may be enhanced by the MITP-M, and this hypothesis should be further investigated.

Modifying the MITP-M for the future

The MITP-M intervention is designed as a pre- and post-discharge intervention and was started the last week of the child's hospitalization [166,194]. During the last 15 years, the progress observed in TISP in terms of new knowledge has challenged Norwegian NICU's to facilitate higher degrees of parental presence and participation across the period of hospitalization of preterm infants. These changes have been inspired by knowledge concerning developmental care (especially Newborn Individualized Development Care and Assessment Program, NIDCAP) [301-307], previous reports and presentations from TISP and studies focusing on developmental benefits for the preterm child provided with intimate and sustained skin-to-skin contact with primary adults (mother/father) [301,305].

Similar processes have been documented in many countries, and the importance of guiding the parents of preterms throughout the entire NICU stay has been frequently underlined [38,308]. Key elements in such interventions are described as ^{a)} including parents in caregiving activities, ^{b)} introducing and maintaining shared attention of the child's development, ^{c)} shape-structured settings for learning, ^{d)} building bridges between familiar and new knowledge, and gradually transferring responsibility to parents in terms of suggesting problem solutions and next performing activities [308]. This is also in line with recommendations from Fegran after her study in a Norwegian NICU [2,7] and in line with the "old" recommendations by Bromwich [124].

Research supporting this statement may be exemplified by an Australian study [309]. In that study, they modified the MITP to an intervention with 9 parental-guiding sessions that were delivered across the NICU stay followed by one single home visit [309]. This intervention is known as the "Premie Start Program" and was implemented by a team of psychologists that had worked extensively with preterm populations. They reported positive interventional effects as less distressed preterm children at a term-equivalent age (40 weeks GA) [ibid.], enhanced mother-infant interactions, more social and self-regulative behavior among the children and less stressed mothers

at 3 months of age [310] and improved social behavior competencies among the children at 6 months of age [309,310].

A review of interventions aimed at enhancing the mother-child interactional relationship detected eight different interventions that exhibited some efficacy [227,311]. Interventions that promote “cue-based care”, defined as maternal care given to the infant in response to the infant’s behavioral cues, in combination with sensitive responsive mothering, appeared to be the most effective. The early introduction of interventional support has also focused on children becoming less responsive to interventions as they grow older [15]. Others have recently reviewed the use of home-visiting programs for high-risk populations and documented them as positively associated with enhanced parent-child interactions [312].

The MITP-M has given TISP outstanding results with respect to long-lasting positive effects, both related to the development of children, parental well-being and parent-child interactions. The features mentioned above have been largely incorporated into the MITP-M. Further modifications should focus on how parts of the intervention could be administered earlier than the final week before discharge and next on how the sessions after discharge could be administered in a collaboration between the family, MITP-M interventionists in the NICU and community health care providers.

The Norwegian national guidelines, which include a description of the recommended follow-ups of children born prematurely, were published in 2007 [164], which is one year after a summary of knowledge regarding possible effective interventions in the follow-ups of children born preterm [312]. Even the guidelines were influenced by early results from TISP and other national/international research investigations the conclusions and recommendations in the guidelines do not include a systematic pre- and post-discharge intervention such as the MITP-M [164]. This may be because the most promising results from TISP have been published over the last five years. Several elements from the MITP-M intervention were even incorporated into the general recommendations. To date, no research can confirm the sufficiency of these recommendations or to what degree they are followed across the country.

Four home visits during the first 3 months were included in the MITP-M intervention. In regular early follow-up in all municipalities, home visits are performed by public health nurses. The forwarding of

knowledge, reliance and family-nurse relationships of the NICU professionals to primary health care professionals may vary due to routines, work load and attitudes concerning cooperation across health-care levels. Some studies have reported a lack of knowledge regarding the development of children born preterm among public health nurses [314,315]. In particular, in areas with one or two nurses working alone this is frequently mentioned as a factor for concern.

An early follow-up program should be based on an approach in which families in need of different types of support (above standard discharge information) can be ensured in a systematic but also flexible and multiprofessional way. This would imply structured cooperation between professionals and the families and between health care actors at different health care levels. This cooperation should take into account how much knowledge and experience may be available at each family's home. At present, most families want to leave the NICU and hospital as soon the child is sufficiently physiologically regulated, and the nutrition may be given safely at home. Many children leave the hospital more than one month before they reach full-term according to the corrected age (40 w of GA). This observation implies great responsibilities on the parents of children during the transition to the home environments, and a lack of well-experienced support during the first weeks and months may heighten the risk of interactional problems and subsequent developmental risks [311].

As stated by Watson [38], "Leaving the NICU is only the first step". Premies show additional problems as the complexity of the tasks increase, and many problems do not become apparent until later in childhood. Some parents may find it difficult to build parental competencies that allow them to reach the 2-3 highest steps of the staircase described by Bromwich [124,156]. As concluded by several reports, a longer follow-up is highly important for many children born preterm, providing them with the possibility of developing into the best possible version of themselves [316,317].

Long-term follow-up

Some developmental difficulties and the need for specialized treatment/education were detected at all of the follow-ups in TISP and in all of the study groups. Those children/families who asked for it were supported by the study coordinator or director to contact specialized services. In general, this phenomenon may have generated a child-family supportive effect; families did not need to waste time and energy in the search for adequate help when needed. Parents from families in all groups repeatedly mentioned the usefulness of and their thankfulness for the follow-up program in TISP. The gap between the TISP follow-up program and regular services in the Norwegian health-care system concerning children born preterm is wide, and in the future follow-up must be more defined and revised.

In the previously mentioned guidelines [164], recommendations for follow-up routines for children born prematurely are separated between the specialized health care system and the primary health care system. Children with an extremely low birth weight, below 28 w of GA, or children with medical complications are defined as in need of specialized health care follow-up, whereas others are recommended to receive follow-up in the regular primary health care system [ibid]. Recently, The Norwegian Knowledge Centre for the Health Services (Kunnskapssenteret) was asked to summarize prognostic studies of ELBW infants who received acute lifesaving treatment [318]. No Norwegian studies focusing on the evaluation of post-discharge follow-up programs have been found, and thus attention is needed in this area of study.

The results reported in TISP demonstrates the absence of a precise association between birth conditions, neonatal risk conditions and later socio-emotional developmental outcomes, despite the high importance of bio-medical factors. Other factors such as the distress of the children, parental bonding, the mental health conditions and self-efficacy of the parents, access to social support and several other socio-economic factors may greatly influence the developmental outcomes of the children. Research has confirmed that children born preterm (BW < 2000 grams) must be followed up throughout childhood [316,317]. Both proximal and distal environmental factors should be measured in the follow-up after preterm delivery because birth and medical information only predict some of the problems that may be experienced by these children throughout childhood [259].

8. Conclusions

The MITP-M intervention supported a process of gradual normalization of the behavioral-emotional development of children born preterm from birth until the age of nine years.

On average, the children born preterm (BW < 2000 g) in families who received the intervention reported more problems than their healthy, full-term peers throughout childhood. However, at nine years, their teachers and parents rated them as having similar social and academic competencies compared with children who were born full-term and to have similar levels of socio-emotional behavioral problems.

On the other hand, children born preterm (BW < 2000g) in families who received standard information and guidance before and after discharge from the hospital continued to be reported with more problems and less competencies than the two other groups of children.

References

1. Shonkoff J, & Phillips D (Eds.) *From neurons to neighborhoods: The science of early childhood development*. Washington DC, National Academy Press; 2000. 588 p.
2. Fegran L, Helseth S, Slettebø Å. Nurses as moral practitioners encountering parents in neonatal intensive care units. *Nurs Ethics* 2006; 13(1):52-64.
3. Alsop-Shields L, Mohay H. John Bowlby and James Robertson: theorists, scientists and crusaders for improvements in the care of children in hospital. *J Adv Nurs* 2001;35(1): 50-58.
4. Bowlby J. *Attachment and loss*. New York, Basic Books; 1969 & 1982. 496 p.
5. Kennel JH, Gordon D, Klaus M. The effects of early mother-infant separation on later maternal performance. *Pediatr Res* 1970; 4:473-474.
6. Field TM. Effects of early separation, interactive deficits, and experimental manipulation of mother-infant face to face interaction. *Child Dev* 1977; 48:763-772.
7. Fegran L. *Parents and nurses in a neonatal intensive care unit – the development of a mutual beneficial partnership in the care of the infant*. [Thesis]. Faculty of Medicine, University of Oslo; 2009. 80 p.
8. Davis L, Mohay H, Edwards H. Mother's involvement in caring for their premature infants: an historical overview. *J Adv Nurs* 2003; 42(6):578-586.
9. Als H, Tronick E, Adamson L, Brazelton TB. The behavior of the full-term but underweight newborn infant. *Dev Med Child Neurol* 1976; 18(5):590-602
10. Als H, Tronick E, Lester B. The Brazelton Neonatal Behavioral Assessment Scale (BNBAS), *J Abnorm Child Psych* 1977; 5:214-231.
11. Als H. Toward a Synactive Theory of Development: promise for the assessment and support of infant individuality. *Infant Ment Health* 1982;3(4):229-243.
12. Hutchfield K. Family-centered care: a concept analysis. *J Adv Nurs* 1999; 29:1178-87.
13. Institute for patient and family-centered care (IPFCC). Washington, USA. Established 1982. <http://www.ipfcc.org/>.
14. Rikstrygdeverket. Pleiepenger til et medlem med omsorg for et alvorlig sykt barn, 1998 med senere tilføyelser. Rundskriv paragraf 9-11, Downloaded from NAV 06.07.15. <https://www.nav.no/rettskildene/Rundskriv/%C2%A7%9-11+Pleiepenger+til+et+medlem+med+omsorg+for+et+alvorlig+sykt+barn.147958.cms>

15. Benzie, KM, Magill-Evans JE, Hayden KA, Ballantyne M. Key components of early intervention programs for preterm infants and their parents: a systematic review and meta-analysis. *BMC Pregnancy Childbirth* 2013; 13 (suppl 1):S10.
16. Miles MS & Holditch-Davis D. Parenting the prematurely born child: pathways of influence. *Semin Perinatol* 1997; 21(3):254-266.
17. Melnyk BM, Feinstein NF, Alpert-Gillis L, Fairbanks E, Crean HF, Sinkin RA, Stone PW, Small L, Tu X, Gross SJ. Reducing premature infant's length of stay and improving parents' mental health outcomes with the Creating Opportunities for Parent Empowerment (COPE) Neonatal intensive care unit program. *Pediatrics* 2006; 118:e1414-1428.
18. Montirosso R, Fedeli C, Prete AD, Calciolari G, Borgatti R, NEO-ACQUA study group. Maternal stress and depressive symptoms associated with quality of developmental care in 25 Italian neonatal intensive care units: a cross sectional observational study. *Int J Nurs Stud* 2014; 51:994-1002.
19. Aylward GP. Neurodevelopmental outcomes of infants born prematurely. *Dev & Behav Pediatr* 2005; 26(6):427-441.
20. Markestad T, Kaarensen PI, Ronnestad A, Reigstad H, Lossius K, Medbø S, Zanussi G, Englund IE, Skjaerven R, Irgens LM. Early death, morbidity, and need of treatment among extremely preterm infants. *Pediatrics* 2005; 115(5):1289-1298.
21. Eichenwald EC, Stark AR. Management and outcomes of very low birth weight. *New Engl J Med* 2008; 358(16):1700-1711.
22. WHO media centre. Preterm birth, Downloaded November 2014. <http://www.who.int/mediacentre/factsheets/fs363/en/>
23. Blencowe H, Cousens S, Oestergaard MZ, Chou D, Moller A-B, Narwal R, Adler A, Garcia CV, Rohde S, Say L, Lawn JE. National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. *Lancet* 2012; 379:2162-2172.
24. Aarnoudse-Moens, C SH, Weisglas-Kuperus N, van Goudoever JB & Oosterlaan J. Meta-analysis of neurobehavioral outcomes in very preterm and/or very lowbirth weight children. [Review Article] *Pediatrics* 2009; 124(2):717-728.
25. Hack M, Taylor HG, Drotar D, Schluchter M, Cartar L, Andreias L, Wilson-Costello D, Klein N. Chronic conditions, functional limitations, and special health care needs of school-aged children born with extremely low-birth-weight in the 1990s. *JAMA* 2005; 294(3):319-325.
26. Bhutta AT, Cleves MA, Casey PH, Cradock MM, Anand KJS. Cognitive and behavioral

- outcomes of school-aged children who were born preterm. *JAMA* 2002, 288(2):728-737.
27. McCormick MC, Workman-Daniels K, Brooks-Gunn J. The behavioral and emotional well-being of school-age children with different birth weights. *Pediatrics* 1996; 97(1):18-25.
 28. Medisinsk fødselsregister. Folkehelseinstituttet, Oslo.
Downloaded 30.07.15. <http://www.fhi.no/helseregistre/medisinsk-fodselsregister>
 29. March of Dimes, PMNCH, Save the Children, WHO. *Born To Soon: The global action report on preterm birth*. [Edited by Howson CP, Kinney MV, Lawn JE]. WHO, Geneva. 2012.
 30. Bastek JA, Sammel D, Pare`E, Srinivas SK, Posencheg MA, Elovitz MA. Adverse neonatal outcomes: examining the risks between preterm, late preterm, and term infants. *Am J Obstret Gynecol* 2008; 199(367):e1-e8.
 31. Mulder H, Pitchford NJ, Hagger MS, Marlow N. Development of executive function and attention in preterm children: A systematic review. *Dev Neuropsychol* 2009; 34 (4):393-421.
 32. Benzie KM, Harrison MJ, Magill-Evans J. Parenting stress, marital quality, and child behavior problems at age 7 years. *Publ Health Nurs* 2004; 21(2):111-121.
 33. Shirley MA. A behavioral-syndrome characterizing prematurely born children. *Child Dev* 1939; 10:115-128.
 34. Drillien CM. Abnormal neurological signs in the first year of life in low-birthweight infants: possible prognostic significance. *Dev Med Child Neurol* 1972; 14:575-584.
 35. Gray RF, Indurkha A, McCormick MC. Prevalence, stability, and predictors of clinically significant behavior problems in low birth weight children at 3, 5, and 8 years of age. *Pediatrics* 2004; 114(3):736-743.
 36. Elgen I, Sommerfelt K, Markestad T. Population-based, controlled study of behavioural problems and psychiatric disorders in low birthweight children at 11 years of age. *Arch Dis Child Fetal Neonatal Ed* 2002; 87(2):128-132.
 37. Denham SA, Wyatt TM, Bassett HH, Echeverria D, Knox SS. Assessing socio-emotional development in children from a longitudinal perspective. *J Epidemiol Community Health* 2008; 63(Suppl I):i37-i52.
 38. Watson A. Understanding neurodevelopmental outcomes of prematurity. *Adv Neonatal Care* 2010; 10(4):188-193.
 39. Rose-Krasnor L. The nature of social competence: a theoretical review. *Soc Dev* 1997; 6(1):111-135.

40. Feldman R, Greenbaum CW, Yirmiya N. Mother-infant affect synchrony as an antecedent of the emergence of self-control. *Dev Psychol* 1999; 35(5):223- 231.
41. Feldman R. From biological rhythms to social rhythms: Physiological precursors of mother-infant synchrony. *Dev Psychol* 2006; 42(1):175-188.
42. Bakermans-Kranenburg MJ, van IJzendoorn MH, Juffer F. Less Is More: Meta-analyses of sensitivity and attachment interventions in early childhood. *Psychol Bull* 2003; 129(2):195-215.
43. Sterba SK, Prinstein MJ, Cox MJ. Trajectories of internalizing problems across childhood: Heterogeneity, external validity, and gender differences. *Dev Psychopathol* 2007; 19:345-366.
44. Briggs-Gowan MJ, Carter A, Bosson-Hennan J, Guyer AE, Horwitz SM. Are infant-toddler social-emotional and behavioral problems transient? *J Am Acad Child Adolesc Psychiat* 2006; 45(7):849-858.
45. Bongers IL, Koot HM, van der Ende J, Verhulst FC. The normative development of child and adolescent problem behavior. *J Abnorm Psychol* 2003; 112(2):179-192.
46. Tandon M, Cardeli E, Luby J. Internalizing disorders in early childhood: a review of depressive and anxiety disorders. *Child Adolesc Psychiatr Clin N Am* 2009; 18(3):593-610.
47. Achenbach TM. Achenbach System of Empirically Based Assessment (ASEBA). 30.07.15 <http://www.aseba.org/>
48. Achenbach TM. *Manual for the Child Behavior Checklist/2-3 and 1992 Profile*. University of Vermont, Burlington, VT. 1992.
49. Achenbach TM. *Manual for the Child Behavior Checklist/4-18 and 1991 Profile*. Vol. 88. University of Vermont, Burlington, VT. 1991.
50. Achenbach TM. *Manual for the Teacher's Report Form and 1991 Profile*. Vol. 214. University of Vermont, Burlington, VT. 1991.
51. Reyes ADE, Kazdin AE. Informant discrepancies in the assessment of childhood psychopathology: a critical review, theoretical framework, and recommendations for further study. *Psychol Bull* 2005; 131(4): 483-509.
52. Ranger M, Synnes AR, Vinall J, Grunau RE. Internalizing behaviours in school-age children born very preterm are predicted by neonatal pain and morphine exposure. *Eur J Pain* 2014; 18:844-852.
53. Vinall J, Miller SP, Synnes AR, Grunau RE. Parent behaviors moderate the relationship

between neonatal pain and internalizing behaviors at 18 months corrected age in children born very prematurely. *Pain* 2013; 154:1831-1839.

54. Loe IM, Lee ES, Luna B, Feldman HM. Behavior problems of 9 – 16 year old preterm children: Biological, sociodemographic, and intellectual contributions. *Earl Hum Dev* 2011; 87:247-252.
55. Farooqi A, Hägglöf B, Sedin G, Gothefors L, Serenius F. Mental health and social competencies of 10- to 12-year-old children born at 23 to 25 weeks of gestation in the 1990s: A Swedish national prospective follow-up study. *Pediatrics* 2007; 120:118-133.
56. Spittle AJ, Treyvaud K, Doyle LW, Roberts G, Lee KJ, Inder TE, Cheong JLY, Hunt RW, Newnham CA, Anderson PA. Early emergence of behavior and social-emotional problems in very preterm infants. *J Am Acad Child Adolesc Psychiatry* 2009; 48(9):909-918.
57. Van Baar AL, Vermaas J, Knots E, de Kleine MJK, Soons P. Functioning at school age of moderately preterm children born at 32 to 36 weeks' gestational age. *Pediatrics* 2009; 124:251-257.
58. Bartels M, Hudziak JJ, Boomsma DI, Rietveld MJH, van Beijsterveldt TCEM, van den Oord EJC. A study of parent ratings of internalizing and externalizing problem behavior in 12-year-old twins. *J Am Acad Child Adolesc Psychiatry* 2003; 42:1351-1359.
59. Duhig AM, Renk K, Epstein MK, Phares V. Interparental agreement on internalizing, externalizing and total behavior problems: a meta-analysis. *Clin Psychol* 2000; 7(4):435-453.
60. Van der Ende J, Verhulst FC, Tiemeier H. Agreement of informants on emotional and behavioral problems from childhood to adulthood. *Psychol Assessment* 2012; 24(2):293-300.
61. Leijten P, Raaijmakers MAJ, Orobio de Castro B, Matthys W. Does socioeconomic status matter? A meta-analysis on parent training effectiveness for disruptive child behavior. *J Clin Child Adolesc Psychol* 2013; 42(3):384-392.
62. Heiervang E, Goodman A, Goodman R. The Nordic advantage in child mental health: separating health differences from reporting style in a cross-cultural comparison of psychopathology. *J Child Psychol Psychiatry* 2008; 49(6):678-685.
63. Reitz E, Decovic M, Meijer AM. Relations between parenting and externalizing and internalizing problem behavior in early adolescence: Child behavior as moderator and predictor. *J Adolesc* 2006; 29:419-436.
64. Hølling H, Kurth BM, Rothenberger A, Becker A, Schlack R. Assessing psychopathological problems of children and adolescents from 3 to 17 years in a nationwide representative sample: results of the German health interview and examination survey for children and

adolescents (KiGGS). *Eur Child Adolesc Psychiatry* 2008; 17:34-41.

65. Keiley MK, Bates JE, Dodge KA, Pettit GS. A cross-domain growth analysis: Externalizing and internalizing behaviors during 8 years of childhood. *J Abnorm Child Psychol* 2000; 28(2):161-179.
66. Ross G, Lipper EG, Auld PAM. Social competence and behavior problems in premature children at school age. *Pediatrics* 1990; 86:391-397.
67. Campell SB, Shaw DS, Gilliom M. Early externalizing behavior problems: Toddlers and preschoolers at risk for later maladjustment. *Dev Psychopathol* 2000; 12:467-488.
68. Choe DE, Olson SL, Sameroff AJ. Effects of maternal distress and parenting on the development of children's self-regulation and externalizing behavior. *Dev Psychopathol* 2013; 25:437-453.
69. Weijer-Bergsma E, Wijnroks L, Jongmans MJ. Attention development in infants and preschool children born preterm: a review. *Infant Behav Dev* 2008; 31(3): 333–351.
70. Hart EL, Lahey BB, Loeber R, Applegate B, Frick PJ. Developmental change in attention-deficit hyperactivity disorder in boys: a four-year longitudinal study. *J Abnorm Child Psychol* 1995; 23(6):729-749.
71. Ribeiro LA, Zachrisson HD, Schjolberg S, Aase H, Rohrer-Baumgartner N, Magnus P. Attention problems and language development in preterm low-birth-weight children: Cross-lagged relations from 18 to 36 month. *BMC Pediatrics* 2011; 11:59-73.
72. Kievet de JF, van Elburg RM, Lafeber HN, Oosterlaan J. Attention problems of very preterm children compared with age-matched term controls at school-age. *J Pediatr* 2012; 161(5): 824-829.
73. Posner MI, Petersen SE. The attention system of the human brain. *Annu Rev Neurosci* 1990; 13:25-42.
74. Butcher PR, Kalverboer AF, Geuze RH, Stremmelaar EF. A longitudinal study of the development of shift of gaze to a peripheral stimulus in preterm infants with transient periventricular echogenicity. *J Exp Child Psychol* 2002; 82:116-140.
75. Lawson K & Ruff H. Early focused attention predicts outcome for children born prematurely. *J Dev Behav Pediatr* 2004; 25:399-406.
76. Hunnius S. *Through the eyes of an infant. The early development of visual scanning and disengagement of attention.* [Thesis]. Rijksuniversiteit Groningen, The Netherlands. 2004. 169 p. Downloaded 30.07.15. <https://www.rug.nl/research/portal/files/2925110/thesis.pdf>

77. Rose SA, Feldman JF, Jankowski JJ. Attention and recognition memory in the first year of life: a longitudinal study of preterm and full-term infants. *Dev Psychol* 2001; 37:135-151.
78. Groen-Blokhuis MM, Middeldorp CM, van Beijsterveldt CEM, Boomsma DI. Evidence for a causal association of low birth weight and attention problems. *J Am Acad Child Adolesc Psychiat* 2011; 50(12):1247-1254.
79. Treyvaud K, Ure A, Doyle LW, Lee KJ, Rogers CE, Kidokoro H, Inder TE, Anderson PJ. Psychiatric outcomes at age seven for very preterm children: rates and predictors. *J Child Psychol Psychiatry* 2013; 54(7):772-779.
80. Feldman R. The development of regulatory functions from birth to 5 years: Insights from premature infants. *Child Dev* 2009; 80(2):544-561.
81. Bul KCM, van Baar AL. Behavior problems in relation to sustained selective attention skills of moderately preterm children. *J Dev Phys Disabil* 2012; 24:111-123.
82. Delobel-Ayoub M, Arnaud C, White-Koning M, Casper C,... Larroque B and for the EPIPAGE study group. Behavioral problems and cognitive performance at 5 years of age after very preterm birth: The EPIPAGE study. *Pediatrics* 2009; 123:1485-1492.
83. Elgen I, Sommerfelt K. Low birthweight children: coping in school? *Acta Paediatr Scand* 2002; 91(8):939-945.
84. Elgen I, Lundervold AJ, Sommerfelt K. Aspects of inattention in low birth weight children. *Pediatr Neurol* 2003; 30 (2):92-97.
85. Cavell TA. Social adjustment, social performance, and social skills: a tri-component model of social competence. *J Clin Child Psychol* 1990, 19:111-122.
86. Ritchie K, Bora S, Woodward LJ. Social development of children born very preterm: a systematic review. *Dev Medicine Child Neurol* 2015 Apr 27. doi: 10.1111/dmcn.12783.[Epub ahead of print]
87. Carter AS, Briggs-Gowan MJ, Davis NO. Assessment of young children's social-emotional development and psychopathology: recent advances and recommendations for practice. *J Child Psychol Psychiatry* 2004; 45(1): 109-134.
88. Briggs-Gowan MJ, Carter AS, Skuban EM, Horwitz SM. Prevalence of social-emotional and behavioral problems in a community sample of 1- and 2 year-old children. *J Am Acad Child Adolesc Psychiatry* 2001; 40: 811-819.
89. Horwitz SM, Gary LC, Briggs-Gowan MJ, Carter AS. Do needs drive services use in young children? *Pediatrics* 2003; 112:1373-1378.

90. Feldman R, Dollberg D, Nadam R. The expression and regulation of anger in toddlers: Relations to maternal behavior and mental representations. *Infant Behav Dev* 2011; 34: 310-320.
91. Winchester SB, Sullivan MC, Marks AK, Doyle T, DePalma J, McGrath MM. Academic, social and behavioral outcomes at age 12 of infants born preterm. *Western J Nurs Research* 2009; 31(7): 853-871.
92. Bora S, Pritchard VE, Moor S, Austin NC, Woodward LJ. Emotional and behavioral adjustment of children born very preterm at early school age. *J Paediatr Child Health* 2011; 47:863-869.
93. Jones KM, Champion PR, Woodward LJ. Social competence of preschool children born very preterm. *Early Hum Dev* 2013; 89:795-802.
94. Larroque B, Ancel PY, Marchand-Martin L, et al. Special care and school difficulties in 8-year old very preterm children: the Epipage cohort study. *PLoS ONE* 2011, 6:e21361.
95. Feldman R, Eidelman AI, Rotenberg N. Parenting stress, infant emotion regulation, maternal sensitivity, and the cognitive development of triplets: A model for parent and child influences in a unique ecology. *Child Dev* 2004; 75(6):1774-1791.
96. Bernier A, Carlson SM, Whipple N. From external regulation to self-regulation: early parenting precursors of young children's executive functioning. *Child Dev* 2010; 81(1): 326-339.
97. Berg-Nielsen TS, Vikan A, Dahl AA. Parenting related to child and parental psychopathology: a descriptive review of the literature. *Clin Child Psychol Psychiatry* 2002; 7: 529-552.
98. Elgen IB, Holsten F, Odberg MD. Psychiatric disorders in low birthweight young adults. Prevalence and association with assessments at 11 years. *Eur Psychiat* 2012; 28 (7):393-396.
99. Indredavik MS, Vik T, Heyerdahl S, Kulseng S, Brubakk A-M. Psychiatric symptoms in low birth weight adolescents, assessed by screening questionnaires. *Eur Child Adolesc Psychiatry* 2005; 14: 226-235.
100. Johnson S, Marlow N. Preterm birth and childhood psychiatric disorders. *Pediatric Research* 2011; 69: 11R-18R.
101. Nosarti C, Reichenberg A, Murray RM, Cnattingius S, Lambe MP, Yin L, MacCabe J, Rifkin L, Hultman CM. Preterm birth and psychiatric disorders in young adult life. *Arch Gen Psychiatry* 2012; 69(6):E1-8.
102. Saigal S, Rosenbaum PL, Feeny D, Burrows E, Furlong W, Stoskopf BL, Hoult L. Parental perspectives of the health status and health-related quality of life of teen-aged children who

- were extremely low birth weight and term controls. *Pediatrics* 2000; 105: 569-574.
103. Saigal S, Tyson J. Measurement of quality of life of survivors of neonatal intensive care: critique and implications. *Semin Neonatol* 2008; 32: 59-66.
 104. Zwicker JG, Harris SR. Quality of life of formerly preterm and very low birth weight infants from preschool age to adulthood: A systematic review. *Pediatrics* 2008, 121(2):366-376.
 105. World Health Organization. The world health organization QoL assessment (WHOQOL): position paper from the World Health Organization. *Soc Sci Med* 1995; 41(10):1403-1409.
 106. Aggernæs A: *Livskvalitet*. FADL's forlag, København. 1988.
 107. Dinesen SJ, Greisen G: Quality of life in young adults with very low birth weight. *Arch Dis Child Fetal Neonatal Ed* 2001; 85: 165-169.
 108. Matthejat F, Simon B, König U, Quaschner K, Barchewitz C, Felbel D et al. Lebensqualität bei psychisch kranken children und jugendlichen. (Quality of life in children and adolescents with mental health disorders). *Z Kinder Jug-Psych* 2003; 31(4): 293-303.
 109. Jozefiak T. *Quality of life and mental health in children and adolescents: Child and parent perspectives*. [Thesis]. Norwegian University of Science and Technology, NTNU, Trondheim, Norge. 2009.78 p.
 110. Lund LK, Vik T, Lydersen S, Løhaugen GCC, Skranes J, Brubakk A-M, Indredavik MS. Mental health, quality of life and social relations in young adults born with low birth weight. *Health Qual Life Out* 2012; 10:146-155.
 111. Saigal S, Feeny D, Furlong W, Rosenbaum P, Burrows E, Torrance G. Comparison of the health-related quality of life of extremely low birth weight children and a reference group of children at age eight years. *J Pediatr* 1994; 125: 418-425.
 112. Vederhus BJ, Markestad T, Eide GE, Graue M and Halvorsen T. Health-related quality of life after extremely preterm birth: a matched controlled cohort study. *Health Qual Life Out* 2010, 8:53.
 113. Verrrips G, Brouwer L, Vogels T, Taal E, Drossaert C, Feeny D, Verheijden, Verloove-Vanhorick P. Long term follow-up of health-related quality of life in young adults born very preterm or with a very low birth weight. *Health Qual Life Out* 2012; 10:49.
 114. Dahl LB, Kaaresen PI, Tunby J, Handegård BH, Kvernmo S, Rønning JA. Emotional, behavioral, social and academic outcomes in adolescents born with very low birth weight. *Pediatrics* 2006; 118:e449-e459.
 115. Darlow BA, Horwood J, Pere-Bracken HM, Woodward LJ. Psychosocial outcomes of young

adults born very low birth weight. *Pediatrics* 2013; 132: e1521-1528.

116. Sameroff A. [Ed] *The transactional model of development; how children and contexts shape each other*. American Psychological Association, Washington,DC. 2009. 290 p.
117. Aarnoudse-Moens CS, Oosterlaan J, Duivenvoorden HJ, van Goudoever JB, Weisglas-Kuperus N. Development of preschool and academic skills in children born very preterm. *J Pediatrics* 2011; 158:51-56.
118. Wolke D. Psychological development of prematurely born children. *Arch Dis Child* 1995; 78:567-570.
119. Adams-Chapman I. Neurodevelopmental Outcome of the Late Preterm Infant. *Clin Perinatol* 2006; p:947-964.
120. Pihlakoski L, Sourander A, Aromaa M, Rønning JA, Rautava P, Helenius H, Silanpää M. Do antenatal and postnatal parental psychological distress, and recognized need of help predict preadolescent's psychiatric symptoms? The Finnish Family Competence Cohort Study. *Child Psychiatr Hum Dev* 2012; 44(2):305-319.
121. Treyvaud K, Anderson VA, Lee KJ, Woodward LJ, Newnham C, Inder TE, Doyle LW, Anderson PJ. Parental mental health and early socio-emotional development of children born very preterm. *J Pediatrics* 2010; 35(7):768-777.
122. Williford AP, Calkins SD, Keane SP. Predicting change in parenting stress across early childhood: child and maternal factors. *J Abnorm Child Psychol* 2007; 35:251-263.
123. Laucht M, Esser G, Schmidt MH. Developmental outcome of infants born with biological and psychological risks. *J Child Psychol. Psychiat.* 1997; 38(7):843-853.
124. Bromwich RM. Focus on maternal behavior in infant intervention. *Am J Orthopsych.* 1976, 46(3):439-446.
125. Barnard KE, Bee HL, Hammond MA. Developmental changes in maternal interactions with term and preterm infants. *Infant Behav Dev* 1984; 7:101-113.
126. Brachfeld S, Goldberg S, Sloman J. Parent-infant interaction in free play at 8 and 12 months: effects of prematurity and immaturity. *Infant Behav Dev* 1980; 3:289-305.
127. Bustan D, Sagi A. Effects of early hospital-based intervention on mothers and their preterm infants. *J Appl Dev Psychol* 1984; 5:305-317.
128. Forcada-Guex M, Pierrehumbert B, Borghini A, Moessinger A, Muller-Nix C. Early dyadic patterns of mother-infant interactions and outcomes of prematurity at 18 months. *Pediatrics* 2006; 118:e107-e114.

129. Schmid B, Blomeyer D, Buchmann A, Trautmann-Villalba P, Zimmermann U, Schmidt M, Laucht M. Quality of early mother-child interaction associated with depressive psychopathology in the offspring: A prospective study from infancy to adulthood. *J Psychiat Res* 2011; 5:1387-1394.
130. Cronin CMG, Shapiro CR, Casiro OG, Cheang MS, Math M. The impact of very low-birth-weight infants on the family is long lasting. *Arch Pediatr Adolesc Med* 1995; 149: 151-158.
131. Lazarus RS. From psychological stress to the emotions: A history of changing outlooks. *Annu Rev Psychol* 1993; 44:1-21.
132. Deater-Deckard K. Parenting stress and child adjustment: Some old hypotheses and new questions. *Clin Psychol* 1998, 3(2):314-332.
133. Abidin RR. *Parenting Stress Index, Professional Manual*, Third Edition. PAR Psychological Assessment Resources, Inc. 1995.
134. Lovejoy CM, Graczyk PA, O'Hare E, Neuman G: Maternal depression and parenting behavior: a meta-analytic review. *Clin Psychol Rev* 2000; 20(5):561-592.
135. Guralnick MJ: Preventive interventions for preterm children: effectiveness and developmental mechanisms. [Review article]. *J Dev Behav Pediatr* 2012; 33(4):353-364.
136. Kazdin A. Child, parent and family dysfunction as predictors of outcome in cognitive-behavioral treatment of anti-social children. *Behav Res Ther* 1995; 33:383-395.
137. Ong LC, Chandran v, Boo NY. Comparison of parenting stress between Malaysian mothers of four-year old very low birthweight and normal weight children. *Acta Paediatr* 2001; 90: 1464-1469.
138. Halpern LF, Brand KL, Malone AF. Parenting stress in mothers of very-low -birth-weight (VLBW) and full-term infants: A function of infant behavioral characteristics and child-rearing attitudes. *J Pediatr Psychol* 2001; 26(2):93-104.
139. Gray PH, Edwards DM, O'Callaghan MJ, Cuskelly M, Gibbons K. Parenting stress in mothers of very preterm infants - Influence of development, temperament and maternal depression. *Early Hum Dev* 2013; 89:625-629.
140. Voigt B, Brandl A, Pietz J, Pauen S, Kliegel M, Reuner G. Negative reactivity in toddlers born prematurely: indirect and moderated pathways considering self-regulation, neonatal distress and parenting stress. *Infant Behav Dev* 2012; 36: 124-138.
141. Field T. Postpartum depression effects on early interactions, parenting, and safety practices. A review. *Infant Behav Dev* 2010; 33:1-6.

142. Huhtula M, Korja R, Lehtonen L, Haataja L, Lapinleimu H, Rautava P, on behalf the PIPARI study group. Parental Psychological well-being and behavioral outcome of very low birth weight infants at 3 years. *Pediatrics* 2012; 129(4):e937-e944.
143. Huhtula M, Korja R, Lehtonen L, Haataja L, Lapinleimu H, Rautava P, on behalf the PIPARI study group. Associations between parental psychological well-being and socio-emotional development in 5 year-old preterm children. *Early Hum Dev* 2014; 90(3): 119-124.
144. Wilson S, Durbin EC. Effects of paternal depression on fathers' parenting behavior: a meta-analytic review. *Clin Psychol Rev* 2010; 30:167-180.
145. Essex MJ, Klein MH, Cho E, Kalin NH. Maternal stress beginning in infancy may sensitize children to later stress exposure: Effects on cortisol and behavior. *Biol Psychiat* 2002; 52:776-784.
146. Koch F-S, Ludvigsson J, Sepa A. Parents' Psychological stress over time may affect children's cortisol at age 8. *J Pediatr Psychol* 2010; 35(9):950-959.
147. Chang Y, Fine MA. Modeling parenting stress trajectories among low-income young mothers across the child's second and third years: Factors accounting for stability and change. *J Fam Psychol* 2007; 21(4):584-594.
148. Vanderveen JA, Bassler D, Robertson CMT, Kirpalani H. Early interventions involving parents to improve neurodevelopmental outcomes of premature infants: a meta-analysis. *J Perinatol* 2009; 29:343-351.
149. Willinger U, Diendorfer-Radner G, Willnauer R, Jörgl G, Hager V. Parenting stress and parental bonding. *Behav Med* 2005; 31(2):63-69.
150. Deater-Deckard K, Atzaba-Poria N, Pike A. Mother- and father-child mutuality in Anglo and Indian British families: a link with lower externalizing problems. *J Abnorm Child Psychol* 2004; 32(6): 609-620.
151. Smith M. Good parenting: Making a difference. *Early Hum Dev* 2010; 86: 689-693.
152. Allen EC, Laneen MC, Legault C, Naughton MJ, Pivor C, O'Shea TM. Perception of child vulnerability among mothers of former premature infants. *Pediatrics* 2004; 113(2):267-273.
153. Deater-Deckard K & Bulkley J. Parent concerns in long-term follow-up. *Semin Neonatol* 2000; 5:171-178.
154. Goldberg S, DeVitto B. Parenting children born preterm. M.H. Bornstein [Ed] *Handbook of parenting: children and parenting* (2nd. Ed.) p.329-354, Mahwah, NJ: Lawrence Erlbaum Associates. 1995.

155. D'Apolito K. What is an organized infant? *Neonat Network* 1991; 10(1):23-29.
156. Bromwich RM. The interaction approach to early intervention. *Inf Ment Health J* 1990; 11(1):66-79.
157. Sameroff AJ & Chandler MJ. Reproductive risk and the continuum of caretaking causality. In FD Horowitz [Ed] *Review of child development research* (vol 4.) Chicago: University of Chicago Press, 1975.
158. Rauh VA, Nurcombe B, Ruoff P, Jette A, Howell D. The Vermont infant studies project: The rationale for a Mother-Infant Transaction Program. In Bond & Joffe [Ed] *Facilitating Infant and Early Childhood Development*. University Press of New England Hanover and London, 1982.
159. Feldman R. Intervention programs for premature infants: considering potential mechanisms for change. *The Signal* 2002; 10(2&4):1-11.
160. Barnett CR, Leiderman PH, Grobstein R. Neonatal separation, the maternal side of interactional deprivation. *Pediatrics* 1970; 45: 197-205.
161. Klein M & Stern L. Low birthweight and the battered child syndrome. *Am J Dis Child* 1971; 122:15-18.
162. Fraiberg S. Blind infants and their mothers: An examination of the sign system. In M. Levis and LA Rosenblum (Eds.). *The effect of the infant on its caregiver*. New York: John Wiley and Sons, 1974.
163. Caplan G. Patterns of parental response to the crisis of premature birth: A preliminary approach to modifying mental health outcome. *Psychiatry* 1968; 23: 365-375.
164. Sosial og Helsedirektoratet. Faglige retningslinjer for oppfølging av for tidlig fødte barn. IS-1419, Oslo, 2007.
165. Bronfenbrenner U. Is early intervention effective? In BZ Friedlander, GM Sterritt and GE Kirk (Eds.) *Exceptional Infant* (vol 3) New York: Brunner/Mazel. 1975.
166. Rauh VA, Nurcombe B, Achenbach T, Howell C. The Mother-Infant Transaction Program. The content and implications of an intervention for the mothers of low-birthweight infants. *Clin Perinatol* 1990; 17(1): 31-45.
167. Nurcombe B, Howell D, Rauh VA, Teti DM, Ruoff P, Brennan J. An intervention program for mothers of low-birthweight infants: preliminary results. *J Am Acad Child Psychiat* 1984; 23(3): 319-325.
168. Als H. 1986. A synactive model of neonatal behavioral organization: Framework for the

assessment of neurobehavioral development in the premature infant and for support of infants and parents in the neonatal intensive care environment. Part 1: *Theoretical framework*. The Haworth Press. Downloaded from informahealthcare.com 27.09.2013.

169. Rønning JA, Ulvund SE, Kaaresen PI, Dahl LB. Prosjekt tidlig intervensjon 2000. Kan tidlig intervensjon forebygge psykologiske utviklingsforstyrrelser/forsinkelser og forbedre helsetilstanden hos for tidlig fødte barn? University of Tromsø and the University Hospital of North Norway, 1998. [Unpublished protocol].
170. Sameroff AJ, Fiese B. Transactional regulation: the developmental ecology of early intervention. In Shonkoff J, Meisel S. [Ed] *Handbook of Early Childhood Intervention*. p.135-159. Cambridge: Cambridge University Press. 2000.
171. Feldman R. Bio-behavioral Synchrony: A model for integrating biological and microsocial processes in the study of parenting. *Parenting: Science and practice*. Special Issue: The arc of parenting from epigenomics to ethics 2012; 12(2-3):93-267. Psychology Press: New York.
172. Swain JE. Becoming a parent – biobehavioral and brain science perspectives. *Curr Probl Pediatr Adolesc Health Care* 2011; 41:192-196.
173. Feldman, R. Sensitive periods in human social development: New insights from research on oxytocin, synchrony, and high-risk parenting. *Dev Psychopathol* 2015; 27:369-395.
174. Feldman R. Parent-infant synchrony: a biobehavioral model of mutual influences in the formation of affiliative bonds. *Monogr Soc Res Child Dev* 2012; 2:42-51.
175. Evans T, Whittingham K, Boyd RN. What helps the mother of a preterm infant become securely attached, responsive and well-adjusted? *Infant Behav Dev* 2012; 35:1-11.
176. Mehler K, Wendrich D, Kissgen R, Roth B, Oberthuer A, Pillekamo F, Kribs A. Mothers seeing their VLBW infants within 3h after birth are more likely to establish a secure attachment behavior: evidence of a sensitive period with preterm infants? *J Perinatol* 2011; 31: 404-410.
177. Yarcheski A, Mahon NE, Yarcheski TJ, Hanks MM, Cannella BL. A meta-analytic study of predictors of maternal-fetal attachment. *Int J Nurs Studies* 2009; 46: 708-715.
178. Coppola G, Cassibba R, Costatini A. What can make the difference? Premature birth and maternal sensitivity at 3 months of age: The role of attachment organization, traumatic reaction and baby's medical risk. *Infant Behav Dev* 2007; 30: 679-684.
179. Landsem IP, Handegaard BH, Tunby J, Ulvund SE, Rønning JA. Early intervention program reduces stress in parents of pre-terms during childhood, a randomized controlled trial. *Trials* 2014; 15(1): 387.

180. Dabrowska J, Hazra R, Ahern TH, Guo J-D, McDonald AJ, Masgagni F, Muller JF, Young LJ, Rainne DG. Neuroanatomical evidence for reciprocal regulation of the corticotrophin-releasing factor and oxytocin systems in the hypothalamus and the bed nucleus of the stria terminalis of the rat: Implications for balancing stress and affect. *Psychoneuroendocrinology* 2011; 36(9):1312-1326.
181. Tully LA, Arseneault L, Caspi A, Moffitt TE, Morgan J. Does maternal warmth moderate the effects of birth on twins' Attention-Deficit/Hyperactivity Disorder (ADHD) symptoms and low IQ? *J Consult Clin Psych* 2004; 72(2):218-226.
182. Garner AS, Schonkoff JP, Siegel BS, Dobbins MI, Earls MF, McGuinn L, Pascoe J, Wood DL. Early childhood adversity, toxic stress, and the role of the pediatrician: translating developmental science into lifelong health. *Pediatrics* 2012; 129:e224.
183. Warhol JG (Ed). New perspectives in early emotional development. *Pediatrics* 1998; 102(5): 1225-1331.
184. Als H, Duffy FH, McAnulty GB, Rivkin MJ, Vajapeyam S, Mulkern S, Warfield SK, Huppi PS, Butler SC, Conneman N, Fischer C, Eichenwald EC. Early experiences alter brain function and structure. *Pediatrics* 2004; 113: 846-857.
185. Milgrom J, Newnham C, Anderson PJ, Doyle LW, Gemmill AW, Lee K, Hunt RW, Bear M, Inder T. Early sensitivity training for parents of preterm infants: impact on the developing brain. *Pediatr Res* 2010; 67(3):330-335.
186. Johnson SB, Riley AW, Granger DA, Riis J. The science of early life toxic stress for pediatric practice and advocacy. *Pediatrics* 2013; 131:319.
187. Gunnar MR, Quevedo K. The neurobiology of stress and development. *Annu Rev Psychol* 2007; 58:145-173.
188. Tu MT, Grunau RE, Petrie-Thomas J, Haley DW, Weinberg J, Whitfield MF. Maternal stress and behavior modulate relationships between neonatal stress, attention, and basal cortisol at 8 month in preterm infants. *Dev Psychobiol* 2007; 49(2):150-164.
189. Achenbach TM, Phares V, Howell CT, Rauh VA, Nurcombe B. Seven-year outcome of the Vermont intervention program for low-birthweight infants. *Child Dev* 1990; 61(6):1672-1681.
190. Achenbach TM, Howell CT, Aoki M & Rauh VA. Nine-year outcome of the Vermont Intervention Program for low birth-weight infants. *Pediatrics* 1993; 91(1):45-55.
191. Rauh VA, Achenbach TM, Nurcombe B, Howell CT, Teti DM. Minimizing adverse effects of low-birthweight: four-year results of an early intervention program. *Child Dev* 1988, 59:544-553.

192. Kaaresen PI, Rønning JA, Ulvund SE, Dahl LB. A randomized, controlled trial of the effectiveness of an early-intervention program in reducing parenting stress after preterm birth. *Pediatrics* 2006; 118(1):e9-19.
193. Brazelton TB, Nugent K. *The Neonatal Behavioral Assessment Scale*, 3rd ed. MacKeith Press, London. 2011.
194. Smith L: Norwegian translation of the Mother-Infant Transaction Program (MITP). Tromsø Intervention Study on Preterms (TISP) homepage. Downloaded 30.07.15
http://uit.no/Content/365882/MITP_Vermontmodellenn.pdf
195. Horovitz M, Matson JL. Developmental milestones in toddlers with atypical development. *Res Dev Dis* 2011; 32: 2278-2282.
196. Goodman R, Scott S. Comparing the Strengths and Difficulties Questionnaire and the Child Behavior Checklist: is small beautiful? *J Abnorm Psychol* 1999; 27(1):17-24.
197. Goodman A, Goodman R. Strengths and difficulties questionnaire as a dimensional measure of child mental health. *J Am Acad Child Adolesc Psychiat* 2009; 48(4):400-403.
198. Abidin RR. Bibliography for the Parenting Stress Index. online resources, Downloaded 01.11.2013. http://www4.parinc.com/webuploads/productinfo/PSI_Bibliography.pdf.
199. Crnic KA, Gaze C, Hoffman C. Cumulative parenting stress across the preschool period: Relations to maternal parenting and child behavior at Age 5. *Inf Child Dev* 2005; 14:117-132.
200. Haskett ME, Ahern LS, Ward CS, Allaire JC. Factor structure and validity of the Parenting Stress Index-Short Form. *J Clin Child & Adoles Psychol* 2006; 35(2):302-312.
201. Psychological Assessment Resources, Inc. (PAR). <http://www4.parinc.com/>
202. Jozefiak T, Larsson B, Wichstrøm L, Mattejat F and Ravens-Sieberer U. Quality of life as reported by school children and their parents: a cross-sectional survey. *Health Qual Life Out* 2008; 6:34.
203. Ravens-Sieberer U, Bullinger M. Manual for the KINDL-R Questionnaire for measuring health-related Quality of Life in children and adolescents, Revised Version. 2000.
[<http://www.kindl.org>]
204. Helseth S, Jozefiak T. Norwegian version of the KINDL® questionnaire.
[<http://www.kindl.org/daten/pdf/ManEnglish.pdf>]. Downloaded 30.07.15.
205. Twisk JWR. *Applied Multilevel Analysis*. Cambridge: Cambridge University Press, 2006.
206. Singer JD & Willett JB. *Applied longitudinal data analysis, modeling change and event*

occurrence. New York, Oxford University Press, 2003.

207. Hedges LV, Olkin L. *Statistical methods for meta-analysis*. Academic Press, Orlando, FL. 1985.
208. Alsawalmeh YM, Feldt LS. Test of the hypothesis that the intraclass reliability coefficient is the same for two measurement procedures. *Appl Psych Meas* 1992, 16:195-205.
209. Kaaresen PI, Rønning JA, Tunby J, Nordhov SM, Ulvund SE, Dahl LB. A randomized controlled trial of an early intervention program in low birth weight children: outcome at 2 years. *Early Hum Dev* 2008; 84(3):201-209.
210. Nordhov SM, Rønning JA, Ulvund SE, Dahl LB, Kaaresen PI. Early intervention improves behavioral outcomes for preterm infants: randomized controlled trial. *Pediatrics* 2012; 129:9-16.
211. Landsem IP, Handegård BH, Ulvund SE, Tunby J, Kaaresen PI, Rønning JA. Does an early intervention influence behavioral development until age nine of children born prematurely? *Child Dev* 2015; 86(4):1063–1079.
212. Gilliom M, Shaw DS. Codevelopment of externalizing and internalizing problems in early childhood. *Dev Psychopathol* 2004; 16:313-333.
213. Samara M, Marlow N, Wolke D. Pervasive behavior problems at 6 years of age in a total-population sample of children born at ≤ 25 weeks of gestation. *Paediatrics* 2008; 122:562-573.
214. Hall J, Wolke D. A comparison of prematurity and small for gestational age as risk factors for age 6-13 year emotional problems. *Early Hum Dev* 2012; 88(10):797-804.
215. Caravale B, Mirante N, Vagnoni C, Vicari S. Change in cognitive abilities over time during preschool age in low risk preterm children. *Early Hum Dev* 2012; 88:363-367.
216. Stoelhorst GM, Rijken M, Martens SE, van Zwieten PH, Feenstra J, Zwinderman AH, Wit JM, Veen S, Leiden follow-up project on prematurity. Developmental outcomes at 18 and 24 months of age in very preterm children: a cohort study from 1996 to 1997. *Early Hum Dev* 2003; 72(2):83-95.
217. Mürner-Lavanchy I, Steinlin M, Kiefer C, Weisstanner C, Ritter BC, Perrig W, Everts R. Delayed development of neural language organization in very preterm born children. *Dev Neuropsychol* 2014; 39(7):529-542.
218. Wright BA, Zecker SG. Learning problems, delayed development, and puberty. *Proc Acad Sci USA* 2004; 101(26):9942-9946.
219. Tallandini MA, Morsan V, Gronchi G, Macagno F. Systematic and meta-analytic review:

Triggering agents of parental perception of child's vulnerability in instances of preterm birth. *J Ped Psychol* 2015; 40(6): 545-553.

220. Essex MJ, Klein MH, Cho E, Kalin NH. Maternal stress beginning in infancy may sensitize children to later stress exposure: effects on cortisol and behavior. *Soc Biol Psychiatry* 2002; 52: 776-784.
221. Morgan J, Robinson D, Aldridge J. Parenting stress and externalizing child behavior. *Child Fam Sos Work* 2002; 7:219-225.
222. Cappa KA, Begle AM, Conger JC, Dumas JE, Conger AJ. Bidirectional relationships between parenting stress and child coping competence: findings from the pace study. *J Child Fam Stud* 2011; 20:334-342.
223. Allen EC, Manuel JC, Legault C, Naughton MJ, Pivor C, O'Shea TM. Perception of child vulnerability among mothers of former premature infants. *Pediatrics* 2004; 113(2):267-273.
224. Harel H, Gordon I, Geva R, Feldman R. Gaze behaviors of preterm and full-term infants in nonsocial and social contexts of increasing dynamics: visual recognition, attention regulation, and gaze synchrony. *Infancy* 2010; 16(1):69-90.
225. Asok A, Bernard K, Roth TL, Rosen JB, Dozier M. Parental responsiveness moderates the association between early-life stress and reduced telomere length. *Dev Psychopathol* 2013; 25(3): 577-585.
226. Poehlmann J, Schwichtenberg AJM, Schlafer RJ, Hahn E, Bianchi J-P, Warner R. Emerging self-regulation in toddlers born preterm or with low birth weight: Different susceptibility to parenting? *Dev Psychopathol* 2011; 23:177-193.
227. Borghini A, Pierrehumbert B, Miljkovicitch, Muller-Nix C, Forcada-Guex M & Ansermet F. Mothers attachment representations of their premature infant at 6 and 18 months after birth. *Infant Ment Health J* 2006; 27(5):494-508.
228. Fite PJ, Greening L, Stoppelbein L. Relation between parenting stress and psychopathic traits among children. *Behav Sci Law* 2008; 26:239-248.
229. Evans T, Whittingham K, Sanders M, Colditz P, Boyd RN. Are parenting interventions effective in improving the relationship between mothers and their preterm infants? *Infant Behav Dev* 2014; 37:131-154.
230. Erikson SJ, Duvall EW, Fuller J, Schrader R, McLean P, Lowe JR. Differential associations between maternal scaffolding and toddler emotion regulation in toddlers born preterm and fullterm. *Early Hum Dev* 2013; 89:699-704.

231. Olafsen KS, Kaaresen PI, Handegard BH, Ulvund SE, Dahl LB, Rønning JA. Maternal ratings of infant regulatory competence from 6 to 12 months: influence of perceived stress, birth-weight, and intervention: a randomized controlled trial. *Infant Behav Dev* 2008; 31(3):408-421.
232. Olafsen KS, Rønning JA, Handegård BH, Ulvund SE, Dahl LB, Kaaresen PI. Regulatory competence and social communication in term and preterm infants at 12 mnd corrected age. Results from a randomized controlled trial. *Infant Behav Dev* 2012; 35:140-149.
233. Petersen IT, Bates JE, Dodge KA, Lansford JE, Pettit GS. Describing and predicting developmental profiles of externalizing problems from childhood to adulthood. *Dev Psychopathol* 2015; 27(3):791-818.
234. Spinelli M, Poehlman J, Bolt D. Predictors of parenting stress trajectories in premature infant-mother dyads. *J Fam Psychol* 2013; 27(6):873-883.
235. Nordhov SM, Kaaresen PI, Rønning JA, Ulvund SE, Dahl LB. A randomized study of the impact of a sensitizing intervention on the child-rearing attitudes of parent of low birth weight preterm infants. *Scand J Psychol* 2010; 51:385-391.
236. Landsem IP, Handegård BH, Ulvund SE, Kaaresen PI, Rønning JA. Early intervention influences positively quality of life as reported by prematurely born children at age nine and their parents; a randomized clinical trial. *Health Qual Life Out* 2015; 13:25.
237. Buehler C, Anthony C, Krishnakumar A, Stone G, Gerard J, Pemberton S. Interparental conflict and youth problem behaviors: a meta-analysis. *J Child Fam Stud* 1997; 6:233-247.
238. Sloan K, Rowe J, Jones L. Stress and coping in fathers following the birth of a preterm infant. *J Neonat Nurs* 2008; 14:108-115.
239. Poehlmann J, Hane A, Burnson C, Maleck S, Hamburger E, Shah PE. Preterm infants who are prone to distress: differential effects of parenting on 36-month behavioral and cognitive outcomes. *J Child Psychol Psychiatry* 2012; 53(10):1018-1025.
240. Hauser-Cram P, Warfield ME, Shonkoff JP, Krauss MW, Sayer A, Upshur CC, Hodapp RM. Children with disabilities: a longitudinal study of child development and parent well-being. *Monogr Soc Res Child* 2001; 66 (3). Published by Wiley on behalf of the Society for Research in Child Development. <http://www.jstor.org/stable/3181571> . UNN, downloaded 15.01.14.
241. Schothorst PF, van Engeland H. Long-term behavioral sequelae of prematurity. *J Am Acad Child Adolesc Psychiatry* 1996; 35(2):175-183.
242. Boyd LAC, Msall ME, O'Shea TM, Allred EN, Hounshell G, Leviton A. Social-emotional delays at 2 years in extremely low gestational age survivors: Correlates of impaired

- orientation/engagement and emotional regulation. *Early Hum Dev* 2013; 89:925-930.
243. Treyvaud K, Doyle LW, Lee KJ, Roberts G, Lim J, Inder TE, Anderson PJ. Social-emotional difficulties in very preterm and term 2 years olds predict specific social-emotional problems at the age of 5 years. *J Ped Psychiatr* 2012; 37(7): 779-785.
244. Treyvaud K, Lee KJ, Doyle LW, Anderson PJ. Very preterm birth influences parental mental health and family outcomes seven years after birth. *J Pediatr* 2014; 164(3):515-521.
245. Lund LK, Vik T, Skranes J, Lydersen S, Brubakk AM, Indredavik MS. Low birth weight and psychiatric morbidity; stability and change between adolescence and young adulthood. *Early Hum Dev* 2012; 88(8):623-629.
246. Bilenberg N, Petersen D, Hoerder K, Gillberg C. The prevalence of child-psychiatric disorders among 8-9-year-old children in Danish mainstream schools. *Acta Psych Scand* 2005; 111:59-67.
247. Landry SH, Smith KE, Swank PR. Responsive parenting: Establishing early foundations for social, communication, and independent problem-solving skills. *Dev Psychol* 2006; 42(4):627-642.
248. Tu MT, Gruneau RE, Petrie-Thomas J, Haley DW, Weinberg J, Whitfield MF. Maternal stress and behavior modulate relationships between neonatal stress, attention, and basal cortisol at 8 month in preterm infants. *Dev Psychobiol* 2007; 49(2):150-164.
249. Davis EP, Bruce J, Gunnar MR. The Anterior Attention Network: Associations with temperament and neuroendocrine activity in 6-years old children. *Dev Psychobiol* 2002; 40:43-56.
250. Cuevas K, Bell MA. Infant attention and early childhood executive function. *Child Dev* 2014; 85(2):397-404.
251. Brogan E, Cragg L, Gilmore C, Marlow N, Simms V, Johnson S. Inattention in very preterm children: implications for screening and detection. *Arch Dis Child* 2014; 99:834-839.
252. Saigal S, den Ouden L, Wolke D, Hoult L, Paneth N, Streiner DL, Whitaker A, Pinto-Martin J. School age outcomes in children who were extremely low birth weight from four international population-based cohorts. *Pediatrics* 2003; 112(4):942-950.
253. Jeyaseelan D, O'Callagan M, Neulinger K, Shum D, Burns Y. The association between early minor motor difficulties in extreme low birth weight infants and school age attentional difficulties. *Early Hum Dev* 2005; 82:249-255.
254. Keller-Margulis M, Dempsey A, Llorens A. Academic outcomes for children born preterm: A

Summary and call for research. *Early Child Educ* 2011; 39:95-102.

255. Reijneveld SA, de Kleine MJK, van Baar AL. Behavioral and emotional problems in very preterm and very low birth weight infants at age 5 years. *Arch Dis Fetal Neonatal Ed* 2006; 91: F423-F428.
256. Skranes J, Vangberg TR, Kulseng S, Indredavik MS, Evensen KAI, Martinussen M, Dale AM, Haraldseth O, Brubakk AM. Clinical findings and white matter abnormalities seen on tensor imaging in adolescents with very low birth weight. *Brain* 2007; 130:654-666.
257. Hauglann L, Handegård BH, Ulvund SE, Nordhov M, Rønning JA, Kaaresen PI. Cognitive outcome of early intervention in preterms at 7 and 9 years of age: a randomized controlled trial. *Arch Dis Child Fetal Neonatal Ed.*, 2015, 100(1):pp.F11-6.
258. Nordhov SM, Rønning JA, Dahl LB, Ulvund SE, Tunby J, Kaaresen PI. Early intervention improves cognitive outcomes for preterm infants: randomized controlled trial. *Pediatrics* 2010; 126(5): e1088-1094.
259. Aylward GP. Cognitive and neuropsychological outcomes: more than IQ scores. *Ment Retard Dev D R* 2002; 8:234-240.
260. Munck P, Haataja L, Maunu J, Parkkola R, Rikalainen H, Lapinleimu H, Lethonen L, PIPARI Study Group. Cognitive outcome at 2 years of age in Finnish infants with very low birth weight born between 2001 and 2006. *Acta Pædiatrica* 2010; 99:359-366.
261. Gray PH, Edwards DM, O'Callaghan MJ, Cuskelly M. Parenting stress in mothers of preterm infants during early infancy. *Early Hum Dev* 2012; 88:45-49.
262. Costa NM, Weems CF, Pellerin K, Dalton R. Parenting stress and childhood psychopathology: an examination of specificity to internalizing and externalizing symptoms. *J Psychopathol Behav Ass* 2005; 28(2):113-122.
263. Harrison MJ, Magill-Evans J. Mother and father interactions over the first year with term and preterm infants. *Res Nurs Health* 1996; 19:451-459.
264. Lavelli M, Fogel A. Developmental changes in the relationship between the infant's attention and emotion during early face-to-face communication: the 2 month transition. *Dev Psychol* 2005; 41(1): 265-280.
265. Brummelte S, Chau CMY, Cepeda IL, Degenhardt A, Weinberg J, Synnes AR, Grunau RE. Cortisol levels in former preterm children at school age are predicted by neonatal procedural pain-related stress. *Psychoneuroendocrino* 2015; 51:151-163.
266. Smith GC, Gutovich J, Smyser C, Pineda R, Newnham C, Tjoeng TH, Vavasseur C, Wallendorf M, Neil J, Inder T. Neonatal intensive care unit stress is associated with brain development in

preterm infants. *Ann Neurol* 2011; 70:541-549.

267. Feldman R, Rosenthal Z, Eidelman AI. Maternal-preterm skin-to-skin contact enhances child physiologic organization and cognitive control across the first 10 years of life. *Biol Psychiatry* 2014; 75:56-64.
268. Korja R, Latva R, Lehtonen L. The effects of preterm birth on mother-infant interaction and attachment during the infant's first two years. *Acta Obstet Gynecol Scand* 2012; 91:164-173.
269. Bienfait M, Maury M, Haquet A, Faille JL, Franc N, Combes C, Daude H, Picaud JC, Rideau A, Cambonie G. Pertinence of the self-report mother-to-infant bonding scale in the neonatal unit of maternity ward. *Early Hum Dev* 2011; 87(4):281-287.
270. Stern D, Brushweiler-Stern N, Freeland A. *The birth of a mother: How the Motherhood experience changes you forever*. Basic Books, 1 Ed. 1998. 231 p.
271. Nugent KJ, Keefer CH, Minear S, Johnson LC, Blanchard Y. *Understanding newborn behavior relationships*. Paul H. Brookes Publishing Co., Baltimore. 2007. 256p.
272. Bandura A. Self-efficacy: Toward a unifying theory of behavioral change. *Psychol Rev* 1977; 84(2):191-215.
273. Jones TL & Prinz RJ. Potential roles of parental self-efficacy in parent and child adjustment: a review. *Clin Psychol Rev* 2005; 25:341-363.
274. Wells-Parker E, Miller DI, Topping JS. Development of control-of-outcome scales and self-efficacy scales for woman in four life roles. *J Pers Asses* 1990; 54(3-4):564-575.
275. Lazarus RS. Towards better research on stress and coping. *APA* 2000; 55(6); 665-673. Downloaded 29.07.15 from <http://psycnet.apa.org/journals/amp/55/6/665.pdf>
276. Lazarus RS. From Psychological stress to emotions: A history of changing outlooks. *Annu Rev Psychol* 1993; 44:1-21.
277. Hess CR, Teti DM, Hussey-Gardner B. Self-efficacy and parenting of high-risk infants: The moderating role of parent knowledge of infant development. *Appl Dev Psychol* 2004; 25:423-437.
278. Pennell C, Whittingham K, Boyd R, Sanders M, Colditz P. Prematurity and parental self-efficacy: The preterm parenting & self-efficacy checklist. *Infant Behav Dev* 2012; 35:678-688.
279. Jozefiak T, Larsson B, Wichstrøm L, Wallander J, Matthejat F: Quality of life as reported by children and parents: a comparison between students and child psychiatric outpatients. *Health Qua Life Out* 2010; 8:136.

280. Jozefiak T, Larsson B, Wichstrom L, Rimehaug T. Competence and emotional/behavioural problems in 7-16-year-old Norwegian school children as reported by parents. *Nordic J Psychiatry* 2012; 66(5): 311-319.
281. Jardine J, Glinianaia SV, McConachie H, Embleton ND, Rankin J. Self-reported quality of life of young children with conditions from early infancy: a systematic review. *Pediatrics* 2014; 134: e1129.
282. Eiser C, Eiser JR, Mayhew AG, Gibson AT. Parenting the premature infant: balancing vulnerability and quality of life. *J Child Psychol Psych* 2005; 46:1169-1177.
283. Lee CF, Hwang FM, Chen CJ, Chien LY. The interrelationships among parenting stress and quality of life of caregiver and preschool child with very low birth weight. *Fam Com Health* 2009; 32(3):228-237.
284. Lund LK, Vik T, Lydersen S, Løhaugen GCC, Skranes J, Brubakk AM and Indredavik MS. Mental health, quality of life and social relations in young adults born with low birth weight. *Health Qual Life Out* 2012; 10:146.
285. Kelly MM. Assessment of quality of life after prematurity in 9- to 10-year old children. *MCN* 2014; 39(1):26-32.
286. Østberg M, Hagekull B. Parenting stress and external stressors as predictors of maternal ratings of child adjustment. *Scand J Psychol* 2013; 54:213-221.
287. Spittle A, Ortin J, Anderson P, Boyd R, Doyle RW. Early developmental intervention programmes post-hospital discharge to prevent motor and cognitive impairments in preterm infants [Review]. *Cochrane Database Syst Rev* 2012; 12:CD005495.
288. Tavares TP, Puka K, Smith ML. Emotional functioning: long-term outcomes after pediatric epilepsy surgery. *Epilepsia* 2015; 56(5): 745-753.
289. Doesburg SM, Chau CM, Cheung TPL, Moiseev A, Gruneau RE. Neonatal pain-related stress, functional cortical activity and visual-perceptual abilities in school-age children born at extremely low gestational age. *Pain* 2013; 154:1946-1952.
290. Lyngstad LT, Tandberg BS, Storm H, Ekeberg BL, Moen A. Does skin-to-skin contact reduce stress during diaper change in preterm infants? *Early Hum Dev* 2014; 90:169-172.
291. Olafsen KS, Rønning JA, Kaaresen PI, Ulvund SE, Handegard BH, Dahl LB. Joint attention in term and preterm infants at 12 months corrected age: the significance of gender and intervention based on a randomized controlled trial. *Infant Behav Dev* 2006; 29(4):554-563.
292. Olafsen KS, Rønning JA, Dahl LB, Ulvund SE, Handegård BH, Kaaresen PI: Infant responsiveness and maternal confidence in the neonatal period. *Scand J Psychol* 2007; 48(6):

499-509.

293. Deater-Deckard K, Atzaba-Poria N, Pike A. Mother- and father-child mutuality in Anglo and Indian British families: a link with lower externalizing problems. *J Abnorm Child Psychol* 2004; 32(6):609-620.
294. Vaughan EL, Feinn R, Bernard S, Brereton M, Kaufman JS. Relationships between child emotional and behavioral symptoms and caregiver strain and parenting stress. *J Fam Issues* 2013; 34(4):534-556.
295. Nygren M, Carstensen J, Ludvigsson J, Frostell AS. Adult attachment and parenting stress among parents of toddlers. *J Reprod Infant Psych* 2012; 30(3):289-302.
296. Pohlmann J, Fiese BH. The interaction of maternal and infant vulnerabilities on developing attachment relationships. *Dev Psychopathol* 2001; 13:1-11.
297. Walsh J. Definitions matter: if maternal-fetal relationships are not attachment, what are they? *Arch Wom Ment Health* 2010; 13:449-451.
298. Muller-Nix C, Forcada-Guex M, Pierrehumbert B, Jaunin L, Borghini A, Ansermet F. Prematurity, maternal stress and mother-child interactions. *Early Hum Dev* 2004; 79:145-158.
299. Sherman LJ, Stupica B, Dykas MJ, Ramos-Marcuse F, Cassidy J. The development of negative reactivity in irritable newborn as a function of attachment. *Infant Behav Dev* 2013; 36:139-146.
300. Fearon RMP, Belsky J. Infant-mother attachment and the growth of externalizing problems across the primary-school years. *J Child Psychol Psychiat* 2011; 52(7):782-791.
301. Lode, IU, Andersen, RD, Ericsson M. Bruk av hud-mot-hud kontakt I norske nyfødte intensivavdelinger. *Sykepleien Forskn* 2015; 10(2):152-160.
302. Ministry of Health and Care Services. Regulations on children in health care institutions. Oslo, 2013. <https://lovdata.no/dokument/SF/forskrift/2000-12-01-1217>. UNN, downloaded 15.06.2015.
303. Kleberg A, Hellström-Westas L, Widström A-M. Mothers' perception of Newborn Individualized Developmental Care Assessment Program (NIDCAP) as compared to conventional care. *Early Hum Dev* 2007; 83(6):403-411.
304. Ombudsman for Children. *Helse på barns premisser*. Report from the Ombudsman 2013, Oslo. http://barneombudet.no/wp-content/uploads/2014/03/Helse_p%C3%A5_barns_premisser.pdf Downloaded at UNN 30.07.15

305. Blomqvist YT, Nyqvist KH. Initiation and extent of skin-to-skin care at two Swedish neonatal intensive care units. *Acta Paediatr* 2013;102:22–8.
306. Landsem IP, Tunby J, Cheetham NB. Utvikling av klinisk kompetanse og kunnskap – lange veier og mange faser! *Tidsskr Barnesykepl* 2012; 16(3):18-20.
307. Kleberg A. Omsorgsprogrammet. Karolinska Training Center, Stokholm.[unpublished]
308. Schroeder M, Pridham K. Development of relationships competencies through guided participation for mothers of preterm infants. *JOGNN* 2006; 35:358-368.
309. Milgrom J, Newnham C, Martin PR, Anderson PJ, Doyle LW, Hunt RW, Achenbach TM, Ferretti C, Holt CJ, Inder TE, Gemmill AW. Early communication in preterm infants following intervention in the NICU. *Early Hum Dev* 2013; 89:755-762.
310. Newnham CA, Milgrom J, Skouteris H. Effectiveness of a modified Mother-Infant Transaction program on outcomes for preterm infants from 3 to 24 months of age. *Inf Behav Dev* 2009; 32:17-26.
311. Herd M, Whittingham K, Sanders M. Efficacy of preventive parenting interventions for parents of preterm infants on later child behavior: A systematic review and meta-analysis. *Infant Ment Health J* 2014; 35(6):630-641.
312. Goyal NK, Teeters A, Ammerman RT. Home visiting and outcomes of preterm infants: a systematic review. *Pediatrics* 2013; 132(3):502-516.
313. Reinart LM, Kornør H, Langengen IW, Markestad T. Tiltak for oppfølging av for tidlig fødte barn, Nasjonalt kunnskapssenter for helsetjenesten. Rapport 7-2006, 34 sider. Downloaded 03.08.15. <http://www.kunnskapssenteret.no/publikasjoner/tiltak-for-oppfolging-av-for-tidlig-fodte-barn>
314. Olavesen ES. Helsesøsterkompetanse og oppfølging av premature barn, en kartleggingsundersøkelse. [Master-Thesis]. Institutt Sykepl.vitenskap og Helsefag, Medisinsk fakultet, Universitetet i Oslo. 2010, Downloaded 03.08.15 <https://www.duo.uio.no/bitstream/handle/10852/28188/SolstadxOlavesenxMasteroppgave.pdf?sequence=2>
315. Erdal B. Bedre oppfølging av premature. *Sykepleien* 2012; 100(11):53-56.
316. Doyle LW, Anderson PJ, Battin M, Bowen JR, Brown N, Callanan C Spittle AJ, Wolke D, Woodward LJ. Long term follow-up of high risk children: who, why and how? *BMC Pediatrics* 2014; 14:279.
317. Sherman MP, Aylward GP, Lauriello NF. Follow-up of the NICU patient. *Medscape*. Article updated Sep 25, 2013. Downloaded 03.08.15 <http://emedicine.medscape.com/article/1833812->

overview#a6

318. Nasjonalt kunnsapsenter for helsetjenesten. Prognose og oppfølging av ekstremt premature barn. Prosjektbeskrivelse, downloaded 04.08.15
<http://www.kunnsapsenteret.no/prosjekter/prosjektplan-for-prognose-og-oppfolging-av-ekstremt-premature-barn>

APPENDIX

Table 1. Birth, medical and demographic characteristics of infants and parents

	PI Group N = 72	PC Group N = 74	TR Group N = 75
<i>Infant characteristics</i>			
BW, mean \pm SD, g	1396 \pm 429	1381 \pm 436	3619 \pm 490
400 - 1000 g, n (%)	20 (28)	20 (27)	
1001 - 1500 g, n (%)	15 (21)	20 (27)	
1501 - 2000 g, n (%)	37 (51)	34 (46)	
GA, mean \pm SD, wk	30.2 \pm 3.1	29.9 \pm 3.5	39.3 \pm 1.3
< 28 wk, n (%)	17 (24)	19 (27)	
28 - 32 wk, n (%)	36 (50)	37 (50)	
\geq 33 wk, n (%)	19 (26)	18 (24)	
Boy, n (%)	38 (53)	39 (53)	40 (54)
Twin, n (%)	16 (22)	14 (19)	0
Prenatal steroid use, n (%)	53 (74)	57 (77)	
SNAP II, mean \pm SD	8.3 \pm 10.9	10.4 \pm 11.3	
CRIB score, mean \pm SD, N = 85	3.2 \pm 2.8	2.7 \pm 2.9	
Received ventilation, n (%)	29 (40)	37 (50)	
Duration of ventilation, n (%)	7.0 \pm 18.6	7.1 \pm 17.3	
Postnatal steroid use, n (%)	9 (13)	10 (14)	
Oxygen therapy at 38 wk GA, n (%)	11 (15)	14 (19)	
Abnormal cerebral ultrasound, n (%)			
IVH grade 1 or 2	7 (10)	8 (11)	
IVH grade 3 or 4	3 (4)	5 (7)	
Periventricular leukomalacia	4 (6)	8 (11)	
<i>Maternal and social characteristics</i>			
Mother's age, mean \pm SD, y	30.8 \pm 6.1	29.1 \pm 6.4	29.7 \pm 6.1
Firstborn child, n (%)	40 (56)	37 (54)	27 (37)
Mother's education, mean \pm SD, ^{a)}	14.6 \pm 2.8	13.5 \pm 3.2	14.9 \pm 2.8
Father's education, mean \pm SD, ^{a)}	13.8 \pm 3.1	13.5 \pm 3.2	14.4 \pm 3.2
Mother's monthly income, ^{b)}	15.8 \pm 7.7	14.6 \pm 6.7	15.9 \pm 8.0
Father's monthly income, ^{b)}	21.1 \pm 8.7	19.9 \pm 8.1	21.9 \pm 9.8

^{a)} = education in years

^{b)} = in Norwegian 1000 kroner, calculated for 131 families due to 15 twins

CRIB = Clinical Risk Index for Babies

SNAP II = Score of Acute Neonatal Physiology II

IVH = Intraventricular Hemorrhage

Figure 3. Study Flow from randomization up to nine years of age




Figure 4. Contents of the modified Mother-Infant Transaction Program (MITP-M)

1. Ventilation session.

A talk about parent's experiences, feelings and thoughts related to the pregnancy, delivery and stay in the neonatal care unit while the child has been hospitalized.

2. Helping parents to know their baby.

The first session is dedicated to elicit the best alertness possible from the baby. Parents receive a demonstration of how their child may be, at least briefly, responsive to faces, voices and objects and interested in social interactions. Using elements from a NBAS assessment, parents are invited to look, ask and become attached to and proud of their baby's individuality.

3. How does the infant's body inform us about levels of arousal or balance?

This session focus on the expressions of the child's homeostatic reflexive system and how signs of stress or stability become visible in the skin, breathing patterns, automatic movements and visceral system. Parents are taught to understand how they can adapt to this language by adjusting their actions.

4. How do the infant's body movements inform us about levels of arousal or balance?

This session focuses on the infant's movements, tone and posture. Parents are instructed in how these expressions contribute information about the infant's level of organization/distress.

5. Understanding the infant's expression of and transitions between sleep and awake states.

The parents are introduced to their infant's unique expression of states and how states define the infant's availability and capacities.

6. How parents can help the infant become alert and available for interactions

This session builds upon the previous ones by incorporating knowledge about physiological and motor signs of stability or distress during the exploration of infant alertness.

7. How can parents use this knowledge in daily caretaking

Through activities such as waking, diaper changing, feeding and bathing, parents are guided to use knowledge from the previous sessions to provide care with minimal stress and disturbance regarding the infant's organization.

8. An overview of the first six sessions, preparing for discharge

In the last hospital session, all of the topics are discussed, the parents asks questions and elements are repeated if requested.

9. Home visit 1: Adaptation to the new environment at home

How the family may have made adjustments with the infant's needs in mind

10. Home visit 2: Expanding the family repertoire concerning play and mutual engagement

Parent-child social interactions are addressed, and parents' stories about new activities are applauded.

11. Home visit 3: Recognizing the behavioral style of the infant

The third visit, one month after discharge, focuses on parents' observations of their infant's behavioral style and on how they can adapt to the rhythms and capacities of their infant.

12. Home visit 4: Summarizing the program and recognizing developmental change

The program is summarized, and changes that have appeared are reviewed with the parents. The program is terminated, and the relationship between the family and the interventionist is closed in a solemn way.

Questionnaire with extracted elements from SDQ

For administrasjon

ID-nr.: 7 år 9 år

Sterke og svake sider (SDQ-Nor) utvalgte spørsmål

Vennligst kryss av for hvert utsagn: Stemmer ikke, Stemmer delvis eller Stemmer helt. Prøv å svare på alt selv om du ikke er helt sikker eller synes utsagnet virker rart. Svar på grunnlag av barnets oppførsel de siste 6 månedene.

Barnets fornavn

Gutt/Jente

Besvares av: Mor Far Lærer

	Stemmer ikke	Stemmer delvis	Stemmer helt
1. Omtenkksom, tar hensyn til andre menneskers følelser	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4. Deler gjeme med andre barn (godter, leker, andre ting)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6. Ganske ensom, leker ofte alene	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9. Hjelpsom hvis noen er såret, lei seg eller føler seg dårlig	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
11. Har minst en god venn	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
14. Vanligvis likt av andre barn	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
17. Snill mot yngre barn	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
19. Plaget eller mobbet av andre barn	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
20. Tilbyr seg ofte å hjelpe andre (foreldre, lærere, andre barn)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
23. Kommer bedre overens med voksne enn med barn	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Har du andre kommentarer eller bekymringer?

Vær så snill å snu arket - det er noen få spørsmål til på den andre siden

Samlet, synes du at barnet ditt har vansker på ett eller flere av følgende områder:
med følelser, konsentrasjon, oppførsel eller med å komme overens med andre mennesker?

Nei	Ja- små vansker	Ja – tydelige vansker	Ja – alvorlige vansker
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Hvis du har svart "Ja", vennligst svar på følgende spørsmål:

•Hvor lenge har disse vanskene vært tilstede?

Mindre enn en måned	1-5 måneder	6-12 måneder	Mer enn ett år
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

•Blir barnets selv forstyrret eller plaget av vanskene?

Ikke i det hele tatt	Bare litt	En god del	Mye
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

•Påvirker vanskene barnets dagligliv på noen av de følgende områder?

Ikke i det hele tatt	Bare litt	En god del	Mye
Forhold til jevnaldrende <input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Læring av ferdigheter/ leker etc. <input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

•Er vanskene en belastning for deg eller familien som helhet?

Ikke i det hele tatt	Bare litt	En god del	Mye
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Underskrift Dato

Mor / Far / Andre (vennligst beskriv):

Tusen takk for hjelpen

Help-seeking Questionnaire**”Prosjekt tidlig intervensjon 2000”**

For administrasjon

ID-nr : 9 år

--	--	--

Har barnet hatt behov for hjelp/veiledning fra instansene nederst på arket?

Ja Nei

Hvis svaret er ja; sett en strek for når kontakten startet og når den opphørte.

Eks: 1 4
 Fysioterapi |-----|

ÅR	1	2	3	4	5	7	9
Fysioterapi							
PPT							
Habilitering							
Barnevern							
BUP							

**Parents report of family and socio-economic factor prior to discharge
from the hospital**

**Tidlig intervensjon 2000
Spedbarn, Barneavdelingen
RiTØ**

Deltager nr.....

Mors navn.....Født.....

Barnets navn.....Født.....



SOSIALE OG ØKONOMISKE FAKTORER

Sivil status

- Gift.....
Samboende.....
Enslig.....
Annet.....

Mors utdanning

Totalt antall års skolegang
(inkludert grunnskole).....

Gått på høyskole/Universitet

- Nei
 Ja
Hvis ja, antall år.....

Mors yrke (før svangerskapspermisjon)

- Hjemmeværende.....
Trygdet.....
Arbeidsledig.....
Student/skole-elev.....
Deltidsarbeid.....
I så fall, antall timer pr. uke.....
Fulltids yrkesaktiv.....
Yrke.....

Fars utdanning

Totalt antall års skolegang
(inkludert grunnskole).....

Gått på høyskole/Universitet

- Nei
 Ja
Hvis ja, antall år.....

Fars yrke

- Hjemmeværende.....
Trygdet.....
Arbeidsledig.....
Student/skole-elev.....
Deltidsarbeid.....
I så fall, antall timer pr. uke.....
Fulltids yrkesaktiv.....

Yrke.....

SNU ARKET !

BoligEnebolig/rekkehus..... Leilighet

Annet:

 Selveid LeierCam²**Økonomi**Mors inntekt siste måned (for
svangerskapspermisjon). Inkludert evt.
trygdeytelserFars inntekt siste måned (inkludert evt.
trygdeytelser)