

Wheezes, crackles and rhonchi: simplifying description of lung sounds increases the agreement on their classification: a study of 12 physicians' classification of lung sounds from video recordings

Hasse Melbye,¹ Luis Garcia-Marcos,^{2,3} Paul Brand,^{4,5} Mark Everard,⁶ Kostas Priftis,⁷ Hans Pasterkamp,⁸ The ERS task force for lung sounds

To cite: Melbye H, Garcia-Marcos L, Brand P, *et al.* Wheezes, crackles and rhonchi: simplifying description of lung sounds increases the agreement on their classification: a study of 12 physicians' classification of lung sounds from video recordings. *BMJ Open Resp Res* 2016;**3**:e000136. doi:10.1136/bmjresp-2016-000136

Received 11 March 2016
Revised 8 April 2016
Accepted 10 April 2016

ABSTRACT

Background: The European Respiratory Society (ERS) lung sounds repository contains 20 audiovisual recordings of children and adults. The present study aimed at determining the interobserver variation in the classification of sounds into detailed and broader categories of crackles and wheezes.

Methods: Recordings from 10 children and 10 adults were classified into 10 predefined sounds by 12 observers, 6 paediatricians and 6 doctors for adult patients. Multirater kappa (Fleiss' κ) was calculated for each of the 10 adventitious sounds and for combined categories of sounds.

Results: The majority of observers agreed on the presence of at least one adventitious sound in 17 cases. Poor to fair agreement ($\kappa < 0.40$) was usually found for the detailed descriptions of the adventitious sounds, whereas moderate to good agreement was reached for the combined categories of crackles ($\kappa = 0.62$) and wheezes ($\kappa = 0.59$). The paediatricians did not reach better agreement on the child cases than the family physicians and specialists in adult medicine.

Conclusions: Descriptions of auscultation findings in broader terms were more reliably shared between observers compared to more detailed descriptions.

The stethoscope is the quintessential iconic symbol of the medical profession. However, the reputation of this 200-year-old instrument as a useful diagnostic tool in lung disease has been declining since chest radiography became available.¹ Reports on the limited diagnostic value of chest auscultation in conditions like pneumonia² and heart failure³ have contributed to the low standing of chest auscultation among medical experts today. Guidelines from the Global initiative for Obstructive Lung Disease (GOLD) give little credit to lung sounds, and auscultation

KEY MESSAGES

- ▶ How can we improve the agreement on lung sound descriptions?
- ▶ Although auscultation of the lungs is important in medical diagnosis and decision-making, disagreement on the use of terms describing the sounds weakens the diagnostic value of the adventitious lung sounds for chest diseases.
- ▶ Poor agreement was found when 12 observers classified lung sounds from video recordings of 20 patients with lung diseases into detailed categories, whereas acceptable agreement was obtained when the terms were combined into broader categories.

findings are not even listed among clues to early diagnosis of chronic obstructive pulmonary disease (COPD).⁴ In the Global Initiative for Asthma (GINA) guidelines, however, wheezing is mentioned to be a main symptom of asthma.⁵

Clinical studies have shown that lung auscultation is far from useless. Crackles predict radiographically confirmed pneumonia more strongly than any single respiratory symptom,⁶ and wheezes are heard more frequently with increasing severity of bronchial airflow limitation.⁷ However, the results of such studies may be challenged when one takes into consideration the interobserver variation in the description of lung sounds between clinicians. Efforts to standardise the terminology led to a statement from the International Lung Sound Association (ILSA) in 1987.⁸ The European Respiratory Society (ERS) established a Task Force on Lung sounds in 1999, and a report aiming at



CrossMark

For numbered affiliations see end of article.

Correspondence to
Dr Hasse Melbye;
hasse.melbye@uit.no

the standardisation of computerised analysis of lung sounds, including a chapter on terminology, was published in 2000.⁹ An overview of our present knowledge on lung sounds and recommended terminology has recently been published.¹⁰

In a review on interobserver agreement on chest findings and their diagnostic value published in 2010, the authors found great variation between the published studies in terms of lung sounds, with mostly fair to moderate agreement in the use of terminology.¹¹ In a study from primary care in 12 European countries in 2007, great variation between the countries was found in the use of lung sound terms in patients with acute cough.¹² It seems, therefore, that earlier attempts to standardise lung sounds terminology have failed to improve agreement between physicians on how to describe the lung sounds they hear during chest auscultation. This is important because lung auscultation is still commonly used in clinical practice, and the findings have an impact on the treatment of patients. Hearing crackles, for instance, strongly predicts antibiotic prescribing.^{13 14} The current nomenclature system with its distinction between coarse and fine crackles and between rhonchi and wheezes¹⁰ may make it more difficult to reach agreement on the use of terms.¹⁵ The differences in terminology between different languages hampers meaningful exchange of chest auscultation findings between clinicians and researchers from different countries. These considerations prompted the institution of another ERS Task Force in 2012 to further standardise the use of lung sound terminology between clinicians and researchers from countries with different languages, based on a repository of audiovisual recordings of lung auscultation.

An initial reference collection of 20 audiovisual recordings has recently been made available online as part of the ERS site Learning resources. We wanted to determine the interobserver variation in the classification of these sounds, with particular attention to more or less detailed descriptions of adventitious sounds. We also wanted to assess a potential influence of the professional background, that is, paediatrician versus family physician or adult medical specialist, on the classification of recordings from children and adults.

MATERIAL AND METHODS

This is a study of interobserver variation between medical doctors in classifying lung sounds from video recordings performed at respiratory units in Spain, Greece, Norway, the Netherlands and Canada during 2013. The recording equipment was standardised, using video cameras with external microphone input (Canon Legria HF series, Panasonic HC-X900) and quality electret microphones which were placed in the tubes of standard stethoscopes. More details have already been published.¹⁶ Each centre obtained ethics approval at their institution and prepared consent forms in their required formats.

The quality of 80 video recordings were rated by five of the six task force members, leaving out the member who had done the actual recordings, and they recommended whether or not to include the recording on the repository. Twenty recordings, 10 of children and 10 of adults, were judged to be of sufficient quality. The recordings, each of approximately 15 s duration, were classified independently by the six task force members on an online questionnaire (FluidSurveys.com, Ottawa, Canada). Five of the task force members were paediatricians and one a general practitioner (GP). Their age ranged from 52 to 64 years. Subsequently, the 20 recordings were classified by a convenient sample of six additional physicians who used the same online questionnaire. Three were internationally recognised lung sound researchers aged 59–71 years, among whom one was a paediatrician, and three Norwegian GPs aged 38–49 years. The latter had postgraduate clinical experience of 6 years or more, but no exposure to lung sound research. The observers downloaded the video files to their computers and were asked to classify the recordings according to recommended English language nomenclature (fine and coarse crackles, high-pitched and low-pitched wheezes).⁸ We also included the category of rhonchi, although it is used interchangeably with low-pitched wheezes.¹⁰ Identification by respiratory phase thus offered 10 non-exclusive choices. The questionnaire also offered free-text options to describe adventitious sounds by other terms.

All observers reported normal hearing capacity, except for one 71-year-old expert who reported some trouble with speech perception at higher frequencies, but not with lung sounds that were typically below 1000 Hz. No instructions were given regarding the volume setting for audio playback.

Statistical analysis

Agreement among the majority of observers (seven or more) on each of the 10 terms in each of the 20 cases was identified. To further evaluate this interobserver agreement, Fleiss' multirater kappa (κ) with 95% CIs was calculated. The κ values were interpreted as follows: 0–0.20 slight, 0.21–0.40 fair, 0.41–0.60 moderate, 0.61–0.80 substantial, and 0.80–1.0 almost perfect agreement.¹⁷ After calculating agreement on the 10 predefined categories, multirater κ was calculated after combining fine and coarse crackles in a common category, rhonchi with high-pitched and low-pitched wheezes, and finally the respective inspiratory and expiratory sounds into simply crackles and wheezes. Agreement in reporting other sounds than the 10 fixed options were elevated for sounds reported by seven or more observers to be present in the same case. We also calculated Fleiss' κ among the paediatricians and the observers familiar with examining adults in subsamples of children and adult patients. SPSS V.22 and R statistical package were used in the analyses.

RESULTS

Frequencies of agreement

High-pitched expiratory wheeze was the predefined sound category most frequently reported by the observers, and in 5 of the 20 cases the majority (seven or more) of the observers reported this sound. The majority of observers never reached this level of agreement on the terms expiratory fine crackles, inspiratory or expiratory rhonchi, and inspiratory low-pitched wheezes. The term low-pitched wheezes was more frequently used than rhonchi and when these interchangeable terms were combined, better agreement was reached (figure 1), and it was even better when combined with high-pitched wheezes. Likewise, when fine and coarse crackles were combined into one category, agreement among the majority of the task force members occurred more frequently (figure 1). Such agreement on the presence of one or more of the four sound categories (inspiratory and expiratory crackles and wheezes) was reached in 16 of the 20 cases. The majority agreed on more than one of the four categories in 8 of the 20 cases, in 2 adult cases and 6 child cases. In one case, the majority of observers reported pleural rub (table 1).

Multirater κ agreements

Slight multirater kappa agreement was found for 5 of the 10 basic descriptions of lung sounds ($\kappa \leq 0.20$), fair agreement for four of the categories ($\kappa 0.21-0.40$) and moderate agreement for one category only, high-pitched inspiratory wheezes with $\kappa=0.43$ (figure 2). After combining fine and coarse crackles and high-pitched and low-pitched wheezes together with rhonchi, moderate agreement was reached for three of the four categories (figure 3). An even better agreement was reached after further lumping inspiratory and expiratory sounds, with kappas for crackles and wheezes of 0.62 and 0.59, respectively (figure 3). The agreement on pleural rub reached a κ of 0.52.

Impact of age of cases and kind of physician

The agreement tended to be stronger on the adult cases than on the child cases. The paediatricians did not

reach better agreement on the child cases than the doctors familiar with examining adults (figure 4).

DISCUSSION

Only slight to fair agreement was found for detailed descriptions of the adventitious sounds. In contrast, moderate to substantial agreement was reached for the combined categories of crackles and wheezes. For the wheezes, there was also moderate to substantial agreement when differentiating between inspiratory and expiratory sounds. The paediatricians did not agree better than the other doctors on the lung sounds from children. Overall, agreement on paediatric lung sounds was poorer than that on lung sounds from adults.

Similar or somewhat stronger agreements on the presence of crackles and wheezes were found in this study compared to most previous studies.¹⁸⁻²⁹ In most previous studies, the observers listened to real patients and registered crackles and wheezes without specifying the respiratory phase. Moderate agreement with κ values between 0.41 and 0.60 have usually been found, also in a study where a teaching stethoscope was used, allowing four observers to listen simultaneously.²⁵ In a few studies, the observers have listened to audiotapes and reached similar or somewhat better agreements.^{29,30} No previous interobserver study of auscultation findings has been based on high-quality audiovisual recordings with a microphone inserted into the tubing of a regular stethoscope. The most comparable previous study had observers watching video recordings and registering wheezing heard without a stethoscope. In that study, the multirater kappa agreement was 0.36.³¹ The moderate to substantial agreement found in this and previous interobserver studies on lung sounds is not inferior to agreement reached in other examinations frequently used in pulmonary medicine, like consolidation on chest radiography ($\kappa 0.4-0.6$)³² and the CT patterns of bronchiectasis, emphysema and honeycombing ($\kappa 0.42-0.59$).³³

In many previous studies, detailed descriptions of the sounds have been based on computerised analysis. Fine

Figure 1 The number among the 20 video recorded cases on which 7 or more of the 12 observers reached agreement on the presence of detailed and combined categories of crackles and wheezes.




Table 1 Number of the 12 observers who agreed on auscultation findings in 20 video cases

Case	Age, years	Sex	Diagnosis	Inspiratory crackles	Expiratory crackles	Inspiratory wheezes	Expiratory wheezes	Pleural rub
1	7	M	Asthma	9	8	11	8	0
2	9	F	Asthma	0	0	10	10	0
3	10	M	Bronchitis	9	7	2	7	0
4	½	M	Bronchiolitis	1	1	5	11	0
5	52	F	COPD	0	0	1	12	0
6	4	M	Bronchiectasis	1	1	11	12	0
7	5	F	Asthma, atelectasis	4	4	3	10	0
8	61	F	Emphysema, lung cancer	1	1	5	11	0
9	78	M	Lung cancer	9	7	7	4	0
10	88	F	Asthma and COPD	0	1	11	12	0
11	2	M	Pneumonia	11	11	1	1	0
12	78	M	Pulmonary fibrosis	11	1	1	1	0
13	6	F	Recurrent LRTI	0	0	1	7	0
14	3	F	Acute LRTI	3	1	7	8	0
15*	3	F	Pneumothorax	0	0	1	1	0
16	77	M	Pulmonary fibrosis, pneumonia	11	2	1	4	0
17	69	M	Pleural haemorrhage	5	5	1	1	7
18*	71	M	Pleural effusion	0	0	1	1	0
19*	79	M	Lung cancer	1	0	0	0	0
20	66	F	Radiation pneumonitis	12	0	0	0	0

Bold font is used when 7 or more of the 12 observers agreed.

*Eleven out of the 12 observers agreed on no adventitious sound in this case.

COPD, chronic obstructive pulmonary disease; LRTI, lower respiratory tract infection.

and coarse crackles have been identified by the duration of each crackle as shown on a phonogram.³⁴ It is more difficult to differentiate the types of crackles by listening. It is, for instance, possible that the amplitude of crackles plays a role in how the sound is perceived.³⁵ Loud

crackles may be perceived to be coarser than faint crackles of the same duration.

It can also be difficult to differentiate between low-pitched and high-pitched wheezes by listening. Separation based on a 200 Hz cut-off has been

Figure 2 Multirater agreement (Fleiss' κ) between 12 observers on detailed descriptions of lung sounds from 20 video recordings.





Figure 3 Multirater agreement (Fleiss' κ) between 12 observers on combined categories of lung sounds from 20 video recordings.



recommended.³⁶ Few observers can perfectly determine the pitch, and observer disagreement on this subclassification is therefore not surprising. Agreement on the presence of rhonchi was particularly poor in our study. This is in accordance with previous studies, where the agreement on rhonchi was much weaker than the agreement on crackles and wheezes.^{24 37} Since the terms 'rhonchi' and 'low-pitched wheezes' are used interchangeably,¹⁰ low agreement could be expected. The task force decided anyway to include 'rhonchi' as a separate category, thinking the term could be preferred for low-pitched wheezes that do not sound musical but more like snoring. With our results, it seems difficult to agree on this division of low-pitched continuous sounds.


Other characteristics of crackles and wheezes may be more important and also easier to identify than those applied in our classification. The number of crackles may be of importance and also the timing during inspiration.³⁴ In terms of wheezes, both sound intensity and the duration of the wheezes in each respiratory phase are of importance. Shim and Williams³⁸ found that the three characteristics high pitch, intensity and spanning

the entire phase were linked to decreased peak expiratory flow rate in patients with asthma.

The 20 cases were from real patients with various lung diseases, both children and adults, and in eight cases more than one category of adventitious sound was presented. Six of the eight complex cases were recordings from children, and these might have been more difficult to classify than the recordings with only one kind of adventitious sound. This may have led to poorer agreement in the paediatric cases than in the adult cases.

To avoid misunderstanding on chest findings by auscultation, the use of combined terms of crackles and wheezes, or similar categories in other languages, should be encouraged when health workers communicate with each other. This does not mean that more precise terms should be discarded. Distinguishing between fine and coarse crackles and high-pitched wheezes and low-pitched wheezes/rhonchi may be important for some diagnoses,³⁴ for example, during early stages of interstitial lung fibrosis when fine inspiratory crackles are heard.³⁹ This may also be relevant in various obstructive airway diseases of young children

Figure 4 Multirater agreement (Fleiss' κ) between 6 paediatricians and 6 doctors for adults on the presence of crackles and wheezes (inspiratory or expiratory) from 10 video recordings of children and 10 video recordings of adults.



where ‘wheeze’ is too broad a term to adequately characterise their lung sounds.⁴⁰ However, large studies with relevant outcomes that take other easily available information into account need to be carried out to prove the clinical usefulness of such differentiation.

Strengths and limitations

The video cases were selected from a larger group of files to ensure only high-quality recordings with few artefacts. Although artefacts are also common in real life, we expect that agreement on all recordings without the application of quality criteria would have been poorer than the results presented here.

The mixture of observers increased the probability of transferable results. The task force members had some clinical information on the cases they had contributed to, but although this could have had some influence on their rating of these cases, this knowledge could not have had any significant effects on the agreements.

The statistical strength of the differences in κ values between detailed and lumped categories of sounds may be questioned. The CI of the κ for inspiratory crackles (figure 3) did not include the κ values for fine or coarse inspiratory crackles (figure 2), and that of expiratory crackles did not include the κ of fine expiratory crackles. Likewise, the CIs for inspiratory and expiratory wheezes did not include the respective κ values for rhonchi and low-pitched wheezes (figures 2 and 3). This indicates that improved agreements after lumping were statistically significant.

CONCLUSION

We found only slight to fair agreement for detailed descriptions of crackles and wheezes. Broader terms were more reliably shared between the observers.

Author affiliations

¹Faculty of Health Sciences, General Practice Research Unit, UIT the Arctic University of Norway, Tromsø, Norway

²Pediatric Respiratory and Allergy Units, Arrixaca University Children's Hospital, University of Murcia, Murcia, Spain

³IMIB-Arrixaca Biohealth Research Institute, Murcia, Spain

⁴Princess Amalia Children's Center, Isala Hospital, Zwolle, The Netherlands

⁵Postgraduate School of Medicine, University Medical Centre and University of Groningen, Groningen, The Netherlands

⁶School of Paediatrics, University of Western Australia, Princess Margaret Hospital, Subiaco, Western Australia, Australia

⁷Children's Respiratory and Allergy Unit, Third Dept of Paediatrics, "Attikon" Hospital, University of Athens Medical School, Athens, Greece

⁸Section of Respirology, Dept of Pediatrics and Child Health, University of Manitoba, Winnipeg, Manitoba, Canada

Acknowledgements The authors would like to thank the patients who have taken part in the video recording, the medical student Raimonda Einarsen and Stian Andersen for shooting videos at the University Hospital of North Norway, Drs Steve Kraman, Andrew Bush, Päivi Pirilä, Peder A. Halvorsen, Magnus Hjortdahl and Anne H. Davidsen for the classification of the sounds, Dr Juan Carlos Aviles Solis for statistical assistance and Dr Peter M. A. Calverley for promoting the establishment of the ERS Task Force for Lung Sounds.

Contributors HM takes final responsibility for the content of this manuscript, including the data and analysis. HP designed the questionnaire, and all authors classified the sounds. HM, LG-M, HP and PB contributed to data analysis, and HM, LG-M, HP, PB, ME and KP contributed to the interpretation of results and writing of the manuscript. All authors approved the final version.

Funding Financial support has been given by The European Respiratory Society

Competing interests None declared.

Ethics approval Each centre obtained ethics approval at their institution and prepared consent forms in their required formats.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. Loudon R, Murphy RL Jr. Lung sounds. *Am Rev Respir Dis* 1984;130:663–73.
2. Osmer JC, Cole BK. The stethoscope and roentgenogram in acute pneumonia. *South Med J* 1966;59:75–7.
3. Wang CS, FitzGerald JM, Schulzer M, et al. Does this dyspneic patient in the emergency department have congestive heart failure? *JAMA* 2005;294:1944–56.
4. Vestbo J, Hurd SS, Agustí AG, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 2013;187:347–65.
5. Reddel HK, Bateman ED, Becker A, et al. A summary of the new GINA strategy: a roadmap to asthma control. *Eur Respir J* 2015;46:622–39.
6. van Vugt SF, Verheij TJ, de Jong PA, et al. Diagnosing pneumonia in patients with acute cough: clinical judgment compared to chest radiography. *Eur Respir J* 2013;42:1076–82.
7. Oshaug K, Halvorsen PA, Melbye H. Should chest examination be reinstated in the early diagnosis of chronic obstructive pulmonary disease? *Int J Chron Obstruct Pulmon Dis* 2013;8:369–77.
8. Mikami R, Muro M, Cugell DW, et al. International Symposium on Lung Sounds. Synopsis of proceedings. *Chest* 1987;92:342–5.
9. ERS Task Force Report, ed Sovijarvi A VJEJ. Computerized Respiratory Sound Analysis (COSA): recommended standards for terms and techniques. *Eur Respir Rev* 10. 2000;77:585–649.
10. Bohadana A, Izbicki G, Kraman SS. Fundamentals of lung auscultation. *N Engl J Med* 2014;370:744–51.
11. Benbassat J, Baumal R. Narrative review: should teaching of the respiratory physical examination be restricted only to signs with proven reliability and validity? *J Gen Intern Med* 2010;25:865–72.
12. Francis NA, Melbye H, Kelly MJ, et al. Variation in family physicians' recording of auscultation abnormalities in patients with acute cough is not explained by case mix. A study from 12 European networks. *Eur J Gen Pract* 2013;19:77–84.
13. Jakobsen KA, Melbye H, Kelly MJ, et al. Influence of CRP testing and clinical findings on antibiotic prescribing in adults presenting with acute cough in primary care. *Scand J Prim Health Care* 2010;28:229–36.
14. Patra S, Singh V, Pemde HK, et al. Antibiotic prescribing pattern in paediatric in patients with first time wheezing. *Ital J Pediatr* 2011;37:40.
15. Pasterkamp H, Montgomery M, Wiebcke W. Nomenclature used by health care professionals to describe breath sounds in asthma. *Chest* 1987;92:346–52.
16. Pasterkamp H, Brand PL, Everard M, et al. Towards the standardisation of lung sound nomenclature. *Eur Respir J* 2016;47:724–32.
17. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics* 1977;33:159–74.
18. Spiteri MA, Cook DG, Clarke SW. Reliability of eliciting physical signs in examination of the chest. *Lancet* 1988;1:873–5.
19. Mulrow CD, Dolmatch BL, DeLong ER, et al. Observer variability in the pulmonary examination. *J Gen Intern Med* 1986;1:364–7.

20. Holleman DR, Jr, Simel DL, Goldberg JS. Diagnosis of obstructive airways disease from the clinical examination. *J Gen Intern Med* 1993;8:63–8.
21. Badgett RG, Tanaka DJ, Hunt DK, *et al*. Can moderate chronic obstructive pulmonary disease be diagnosed by historical and physical findings alone? *Am J Med* 1993;94:188–96.
22. Badgett RG, Tanaka DJ, Hunt DK, *et al*. The clinical evaluation for diagnosing obstructive airways disease in high-risk patients. *Chest* 1994;106:1427–31.
23. Wang EE, Milner RA, Navas L, *et al*. Observer agreement for respiratory signs and oximetry in infants hospitalized with lower respiratory infections. *Am Rev Respir Dis* 1992;145:106–9.
24. Wipf JE, Lipsky BA, Hirschmann JV, *et al*. Diagnosing pneumonia by physical examination: relevant or relic? *Arch Intern Med* 1999;159:1082–7.
25. Brooks D, Thomas J. Interrater reliability of auscultation of breath sounds among physical therapists. *Phys Ther* 1995;75:1082–8.
26. Elphick HE, Lancaster GA, Solis A, *et al*. Validity and reliability of acoustic analysis of respiratory sounds in infants. *Arch Dis Child* 2004;89:1059–63.
27. Gajdos V, Beydon N, Bommenel L, *et al*. Inter-observer agreement between physicians, nurses, and respiratory therapists for respiratory clinical evaluation in bronchiolitis. *Pediatr Pulmonol* 2009;44:754–62.
28. Prophan P, Dela Rosa RS, Shubina M, *et al*. Wheeze detection in the pediatric intensive care unit: comparison among physician, nurses, respiratory therapists, and a computerized respiratory sound monitor. *Respir Care* 2008;53:1304–9.
29. Puder LC, Fischer HS, Wilitzki S, *et al*. Validation of computerized wheeze detection in young infants during the first months of life. *BMC Pediatr* 2014;14:257.
30. Kiyokawa H, Greenberg M, Shiota K, *et al*. Auditory detection of simulated crackles in breath sounds. *Chest* 2001;119:1886–92.
31. Bekhof J, Reimink R, Bartels IM, *et al*. Large observer variation of clinical assessment of dyspnoeic wheezing children. *Arch Dis Child* 2015;100:649–53.
32. Williams GJ, Macaskill P, Kerr M, *et al*. Variability and accuracy in interpretation of consolidation on chest radiography for diagnosing pneumonia in children under 5 years of age. *Pediatr Pulmonol* 2013;48:1195–200.
33. Walsh SL, Calandriello L, Sverzellati N, *et al*. Interobserver agreement for the ATS/ERS/JRS/ALAT criteria for a UIP pattern on CT. *Thorax* 2016;71:45–51.
34. Piirila P, Sovijarvi AR, Kaisla T, *et al*. Crackles in patients with fibrosing alveolitis, bronchiectasis, COPD, and heart failure. *Chest* 1991;99:1076–83.
35. Piirila P, Sovijarvi AR. Crackles: recording, analysis and clinical significance. *Eur Respir J* 1995;8:2139–48.
36. Meslier N, Charbonneau G, Racineux JL. Wheezes. *Eur Respir J* 1995;8:1942–8.
37. Wilkins RL, Dexter JR, Murphy RL Jr, *et al*. Lung sound nomenclature survey. *Chest* 1990;98:886–9.
38. Shim CS, Williams MH Jr. Relationship of wheezing to the severity of obstruction in asthma. *Arch Intern Med* 1983;143:890–2.
39. Shirai F, Kudoh S, Shibuya A, *et al*. Crackles in asbestos workers: auscultation and lung sound analysis. *Br J Dis Chest* 1981;75:386–96.
40. Elphick HE, Ritson S, Rodgers H, *et al*. When a “wheeze” is not a wheeze: acoustic analysis of breath sounds in infants. *Eur Respir J* 2000;16:593–7.