



# Journal of Mathematical Inequalities

*With Compliments  
of the Author*

Zagreb, Croatia

Volume 13, Number 3, September 2019

*O. O. Fabelurin, J. A. Oguntuase and L.-E. Persson*

*Multidimensional Hardy-type inequalities on time  
scales with variable exponents*

# JOURNAL OF MATHEMATICAL INEQUALITIES

## AIMS AND SCOPE

Journal of Mathematical Inequalities (**JMI**, J. Math. Inequal.) presents carefully selected original research articles from all areas of pure and applied mathematics, provided they are concerned with mathematical inequalities. **JMI** will also periodically publish invited survey articles with interesting results treating the theory of inequalities, as well as relevant book reviews.

**JMI** is published quarterly, in March, June, September, and December.

## SUBMISSION OF MANUSCRIPTS

Authors are requested to submit their articles electronically as  $\text{\TeX}$  or  $\text{\LaTeX}$  files, with enclosed Adobe Acrobat PDF format through the provided web interface on journal web page.

The author who submitted the article for publication will be denoted as a corresponding author. He/She manages all communication and correspondence with the **JMI** regarding the article, makes any revisions, and reviews and releases the author proofs. Authors may indicate a member of the Editorial Board whom they consider appropriate for the article. However, assignment to that particular editor is not assured.

## COPYRIGHT

The acceptance of the article automatically implies the copyright transfer to the **JMI**. Manuscripts are accepted for review with the understanding that the same work has not been published (except in the form of an abstract), that it is not under consideration for publication elsewhere, that it will not be submitted to another journal while under review for the **JMI**, and that its submission for publication has been approved by all of the authors.

## OPEN ACCESS

Journal of Mathematical Inequalities is published as open access journal. All interested readers are allowed to view, download, print, and redistribute any article without paying for subscription. Offprints of each article may be ordered from **JMI** prior to publication.

## PREPARATION OF MANUSCRIPTS

Manuscripts should be written in English, using prepared journal  $\text{\LaTeX}$  style.

The first page should contain the article title, authors' names (complete first names and family names), complete affiliations (name of institution, address, city, state, and zip code), e-mail addresses, proposed running head (not more than 40 characters), a short abstract (not more than 150 words), a list of key words and phrases, and the AMS 2010 Mathematics Subject Classification primary (and secondary) codes. Avoid abbreviations, mathematical symbols and formulas, diagrams, and reference to the text of the article.

Figures should be prepared in a digital form suitable for direct reproduction, at resolution of 300 dpi or higher, and in EPS, TIFF or JPEG format.

Bibliographic references should be listed alphabetically at the end of the article, each numbered by an Arabic number between square brackets. The following information should be provided for references to journals: names of authors, full title of the article, abbreviated name of the journal, volume, year of publication, and range of page numbers. For standard abbreviations of journal names, the authors should consult the latest Abbreviations of Names and Serials reviewed in Mathematical Reviews.

## SOURCE FILE, PROOFS

Upon acceptance of the article, the authors will be asked to send the related  $\text{\LaTeX}$  source file to the Editorial Office, in accordance to the journal style. In order to accelerate the publication process, the AMS  $\text{\LaTeX}$  package is strongly preferred. PDF proofs will be sent by e-mail to the corresponding author.

## FORTHCOMING PAPERS

Papers accepted and prepared for publication will appear in the forthcoming section of Journal Web page. They are identical in form as final printed papers, except volume, issue and page numbers.

**JMI** is published by Publishing House **ELEMENT**, Zagreb, Croatia.

All correspondence and subscription orders should be addressed to the Editorial Office:

[www.ele-math.com](http://www.ele-math.com)  
e-mail: [jmi@ele-math.com](mailto:jmi@ele-math.com)

Journal of Mathematical Inequalities  
Editorial Office  
Menceticeva 2, 10000 Zagreb, Croatia  
Fax: +385 1 6008799

## MULTIDIMENSIONAL HARDY-TYPE INEQUALITIES ON TIME SCALES WITH VARIABLE EXPONENTS

O. O. FABELURIN, J. A. OGUNTUASE AND L.-E. PERSSON

(Communicated by J. Pečarić)

*Abstract.* A new Jensen inequality for multivariate superquadratic functions is derived and proved. The derived Jensen inequality is then employed to obtain the general Hardy-type integral inequality for superquadratic and subquadratic functions of several variables.

### 1. Introduction

Hardy's discrete inequality reads: if  $p > 1$  and  $\{a_k\}_{k=1}^{\infty}$  is a sequence of nonnegative real numbers, then

$$\sum_{n=1}^{\infty} \left( \frac{1}{n} \sum_{k=1}^n a_k \right)^p \leq \left( \frac{p}{p-1} \right)^p \sum_{n=1}^{\infty} a_n^p. \quad (1.1)$$

Furthermore, G. H. Hardy [9] announced (without proof) that if  $p > 1$  and the function  $f$  is nonnegative and integrable over the interval  $(0, x)$ , then

$$\int_0^{\infty} \left( \frac{1}{x} \int_0^x f(t) dt \right)^p dx \leq \left( \frac{p}{p-1} \right)^p \int_0^{\infty} f^p(x) dx. \quad (1.2)$$

Inequality (1.2) was finally proved by Hardy [10] in 1925. Thus, inequality (1.2) is usually referred to in the literature as the classical Hardy integral inequality while inequality (1.1) is its discrete analogue. The constant  $\left( \frac{p}{p-1} \right)^p$  on the right hand sides of both inequalities (1.1) and (1.2) is the best possible.

Note that (1.1) follows from (1.2), which was pointed out by Hardy [9] but there he also informed that a proof of (1.1) was given to him already in a private letter from E. Landau in 1921. More information concerning the interesting prehistory of Hardy's inequality can be found in [15].

In the last five decades, the Hardy inequality (1.2) has been extensively studied and generalized. A lot of information as regarding applications, alternative proofs, variants, generalizations and refinements abound in the literature (see e.g. the books [11, 16, 17] and the references cited therein).

*Mathematics subject classification* (2010): 26D10, 26D20, 26E70.

*Keywords and phrases:* Multidimensional inequalities, Jensen's inequality, Hardy-type inequalities, time scales, superquadratic functions.

In his PhD thesis, S. Hilger [12] (see also [6, 13, 14]) initiated the calculus of time scales in order to create a theory that will unify discrete and continuous analysis. This new concept has inspired researchers to study Hardy inequalities on time scales. The first known work in this direction is probably due to P. Řehák [19] who in 2005 derived Hardy integral inequality on time scales. Indeed, he showed that

$$\int_a^\infty \left( \frac{1}{\sigma(x)-a} \int_a^{\sigma(x)} f(t) \Delta t \right)^p \Delta x < \left( \frac{p}{p-1} \right)^p \int_a^\infty f^p(x) \Delta x,$$

where  $a > 0, p > 1$  and  $f$  is a nonnegative function.

For notations here and in the sequel see Section 2.

In 2001, R. P. Agarwal et al. [1] obtained the following Jensen's inequality on time scales

$$\Phi \left( \frac{1}{b-a} \int_a^b f(x) \Delta x \right) \leq \frac{1}{b-a} \int_a^b \Phi(f(x)) \Delta x.$$

Moreover, T. Donchev et al. [8] employed the above result to derive the following Hardy-type inequality involving multivariate convex functions on time scales:

**THEOREM 1.1.** *Let  $(\Omega_1, \mathcal{M}, \mu_\Delta)$  and  $(\Omega_2, \mathcal{L}, \lambda_\Delta)$  be two time scale measure spaces and  $U \subset \mathbb{R}^n$  be a closed convex set. Let  $K : \Omega_1 \rightarrow \mathbb{R}$  be defined by  $K(x) := \int_{\Omega_2} k(x, y) \Delta y < \infty$ ,  $x \in \Omega_1$ , where  $k(x, y) \geq 0$  is a kernel. Moreover, let  $\zeta : \Omega_1 \rightarrow \mathbb{R}$  and the weight  $w = w(y)$  be defined by*

$$w(y) := \int_{\Omega_1} \left( \frac{k(x, y) \zeta(x)}{K(x)} \right) \Delta x, \quad y \in \Omega_2.$$

*Then for each convex function  $\Phi$ ,*

$$\int_{\Omega_1} \zeta(x) \Phi \left( \frac{1}{K(x)} \int_{\Omega_2} k(x, y) f(y) \Delta y \right) \Delta x \leq \int_{\Omega_2} w(y) \Phi(f(y)) \Delta y \quad (1.3)$$

*holds for all  $\lambda_\Delta$ -integrable functions  $f : \Omega_2 \rightarrow \mathbb{R}^n$  such that  $f(\Omega_2) \subset U \subset \mathbb{R}^n$ .*

In a recent paper, Oguntuase and Persson [18] presented a number of Hardy-type inequalities on time scales using superquadraticity technique which is based on the application of Jensen dynamic inequality. For some recent developments on Hardy-type inequalities on time scales and related results we refer interested reader to the book [3].

Motivated by the above results, our main aim in this paper is to first establish a Jensen inequality for multivariate superquadratic functions and then employ it to derive some new general Hardy-type inequalities for multivariate superquadratic functions involving more general kernels on arbitrary time scales.

The paper is organized as follows: In Section 2, we recall some basic notions, definitions and results on multivariate superquadratic functions on time scales. In Section 3 we state and prove our main results.

## 2. Preliminaries, definitions and some basic results

First, we recall that a time scale (or measure chain)  $\mathbb{T}$  is an arbitrary nonempty closed subset of the real line  $\mathbb{R}$  with the topology of the subspace  $\mathbb{R}$ . Examples of time scales are the real numbers  $\mathbb{R}$  and the discrete time scale  $\mathbb{Z}$ . Since a time scale  $\mathbb{T}$  may or may not be connected, we need the concept of jump operators. For  $t \in \mathbb{T}$ , we define the forward jump operator  $\sigma : \mathbb{T} \rightarrow \mathbb{T}$  by

$$\sigma(t) = \inf\{s \in \mathbb{T} : s > t\}$$

and the backward jump operator by

$$\rho(t) = \inf\{s \in \mathbb{T} : s < t\}.$$

If  $\sigma(t) > t$ , we say that  $t$  is right-scattered and if  $\rho(t) < t$  we say that  $t$  is left-scattered. The points that are both right-scattered and left-scattered are called isolated. If  $\sigma(t) = t$ , then  $t$  is said to be right-dense, and if  $\rho(t) = t$  then  $t$  is said to be left-dense. The points that are simultaneously right-dense and left-dense are called dense. The mapping  $\mu : \mathbb{T} \rightarrow [0, \infty)$  defined by

$$\mu(t) = \sigma(t) - t$$

is called the graininess function. If  $\mathbb{T}$  has a left-scattered maximum  $M$ , then define  $\mathbb{T}^k = \mathbb{T} \setminus \{M\}$ ; otherwise  $\mathbb{T}^k = \mathbb{T}$ . Let  $f : \mathbb{T} \rightarrow \mathbb{R}$  be a function. Then we define the function  $f^\sigma : \mathbb{T} = \mathbb{R}$  by  $f^\sigma(t) = f(\sigma(t))$  for all  $t \in \mathbb{T}$ . Also, for a function  $f : \mathbb{T} \rightarrow \mathbb{R}$ , the delta derivative is defined by

$$f^\Delta(t) := \lim_{s \rightarrow t, \sigma(s) \neq t} \frac{f^\sigma(s) - f(t)}{\sigma(s) - t}.$$

A function  $f : \mathbb{T} \rightarrow \mathbb{R}$  is called rd-continuous provided it is continuous at all right-dense points in  $\mathbb{T}$  and its left-sided limits exists (finite) at all left-dense points in  $\mathbb{T}$ . We refer interested readers to the books [2], [6] and [7] for more details concerning the calculus of time scales. Note that we have

$$\sigma(t) = t, \mu(t) = 0, f^\Delta = f', \int_a^b f(t) \Delta t = \int_a^b f(t) dt, \text{ when } \mathbb{T} = \mathbb{R},$$

$$\sigma(t) = t + 1, \mu(t) = 1, f^\Delta = \Delta f, \int_a^{b-1} f(t) \Delta t = \sum_{t=a}^{b-1} f(t), \text{ when } \mathbb{T} = \mathbb{Z}.$$

The following Fubini's theorem on time scale in [5] will be needed in the proof of our results in Section 3:

**LEMMA 2.1.** *Let  $(\Omega, \mathcal{M}, \mu_\Delta)$  and  $(\Lambda, \mathcal{L}, \lambda_\Delta)$ , be two finite dimensional time scale measures spaces. If  $f : \Omega \times \Lambda \rightarrow \mathfrak{R}$  is a  $\mu_\Delta \times \lambda_\Delta$ -integrable function and define the function  $\phi(y) = \int_\Omega f(x, y) \Delta x$  for a.e.  $y \in \Lambda$  and  $\varphi(x) = \int_\Lambda f(x, y) \Delta y$  for a.e.  $x \in \Omega$ , then  $\phi$  is  $\lambda_\Delta$ -integrable on  $\Lambda$ ,  $\varphi$  is  $\mu_\Delta$ -integrable on  $\Omega$  and*

$$\int_\Omega \Delta x \int_\Lambda f(x, y) \Delta y = \int_\Lambda \Delta y \int_\Omega f(x, y) \Delta x. \quad (2.1)$$

Moreover, M. Anwar et al. [4] result on the Jensen inequality for convex functions in several variables on time scales will also be needed.

**THEOREM 2.2.** *Let  $(\Omega_1, \Sigma_1, \mu_\Delta)$  and  $(\Omega_2, \Sigma_2, \lambda_\Delta)$  be two time scale measure spaces. Suppose  $U \subset \mathbb{R}^n$  is a closed convex set and  $\Phi \in C(U, \mathbb{R})$  is convex. Moreover, let  $k : \Omega_1 \times \Omega_2 \rightarrow \mathbb{R}$  be nonnegative such that  $k(x, .)$  is  $\lambda_\Delta$ -integrable. Then*

$$\Phi \left( \frac{\int_{\Omega_2} k(x, y) \mathbf{f}(y) \Delta y}{\int_{\Omega_2} k(x, y) \Delta y} \right) \leq \frac{\int_{\Omega_2} k(x, y) \Phi(\mathbf{f}(y)) \Delta y}{\int_{\Omega_2} k(x, y) \Delta y} \quad (2.2)$$

holds for all functions  $\mathbf{f} : \Omega_2 \rightarrow U$ , where  $f_j(y)$  are  $\mu_{\Delta_2}$ -integrable for all  $j \in \{1, 2, \dots, n\}$ , and  $\int_{\Omega_2} k(x, y) \mathbf{f}(y) \Delta(y)$  denotes the  $n$ -tuple

$$\left( \int_{\Omega_2} k(x, y) f_1(y) \Delta(y), \int_{\Omega_2} k(x, y) f_2(y) \Delta(y), \dots, \int_{\Omega_2} k(x, y) f_n(y) \Delta(y) \right).$$

In the sequel, we make the following definitions, assumptions and notations.

- (A1.)  $\Omega_1 = \Omega_2 = [\mathbf{a}, \mathbf{l}] = [a_1, l_1]_{\mathbb{T}} \times [a_2, l_2]_{\mathbb{T}} \dots \times [a_n, l_n]_{\mathbb{T}}$ , where  $0 \leq a_i < l_i \leq \infty$ .
- (A2.)  $\mathbf{a} < \mathbf{b}$  if componentwise  $a_i < b_i$ ,  $i = 1, 2, \dots, n$ .
- (A3.)  $k : [\mathbf{a}, \mathbf{l}] \times [\mathbf{a}, \mathbf{l}] \rightarrow \mathbb{R}_+$  is such that

$$k(\mathbf{x}, \mathbf{y}) = \begin{cases} 1 & \text{if } \mathbf{a} \leq \mathbf{y} < \sigma(\mathbf{x}) \leq \mathbf{l}, \\ 0 & \text{otherwise,} \end{cases} \quad (2.3)$$

that is

$$k(x_1, \dots, x_n, y_1, \dots, y_n) = \begin{cases} 1 & \text{if } a_i \leq y_i < \sigma(x_i) \leq l_i, i = 1, \dots, n \\ 0 & \text{otherwise,} \end{cases} \quad (2.4)$$

- (A4.)  $\Phi(u) = u^p$ ,  $p > 1$ .

**REMARK 2.3.** Under the assumptions (A1- A4), for  $m = 1$ , Theorem 2.2 yields the inequality

$$\begin{aligned} & \left( \frac{1}{\prod_{i=1}^n (\sigma(x_i) - a_i)} \int_{a_1}^{\sigma(x_1)} \dots \int_{a_1}^{\sigma(x_1)} f(y_1, \dots, y_n) \Delta y_1 \dots \Delta y_n \right)^p \\ & \leq \frac{1}{\prod_{i=1}^n (\sigma(x_i) - a_i)} \int_{a_1}^{\sigma(x_1)} \dots \int_{a_1}^{\sigma(x_1)} f^p(y_1, \dots, y_n) \Delta y_1 \dots \Delta y_n. \end{aligned} \quad (2.5)$$

We will also need the following Lemmas for the proof of our main results in the paper.

LEMMA 2.4. Let  $\beta > 0$  and  $a, b, l \in \mathbb{T}$  be such that  $0 \leq a < b \leq l$ .

(i) If  $\beta > 1$ , then

$$\int_b^l (s-a)^{\beta-1} \Delta s \leq \frac{1}{\beta} \left[ (t-a)^\beta - (b-a)^\beta \right] \leq \int_b^l (\sigma(s)-a)^{\beta-1} \Delta s. \quad (2.6)$$

(ii) If  $\beta < 1$ , then

$$\int_b^l (s-a)^{\beta-1} \Delta s \geq \frac{1}{\beta} \left[ (l-a)^\beta - (b-a)^\beta \right] \geq \int_b^l (\sigma(s)-a)^{\beta-1} \Delta s. \quad (2.7)$$

*Proof.* For case (i), let  $\beta > 1$ . Then by applying Keller's chain [6], we find that

$$\begin{aligned} ((t-a)^\beta)^\Delta &= \beta \int_0^1 [h(\sigma(t)-a) + (1-h)(t-a)]^{\beta-1} dh \\ &\geq \beta \int_0^1 [h(t-a) + (1-h)(t-a)]^{\beta-1} dh \\ &= \beta(t-a)^{\beta-1}. \end{aligned}$$

Integrating, we obtain

$$\int_b^l (t-a)^{\beta-1} \Delta t \leq \frac{1}{\beta} \left[ l-a)^\beta - (b-a)^\beta \right]. \quad (2.8)$$

On the other hand,

$$\begin{aligned} ((t-a)^\beta)^\Delta &= \beta \int_0^1 [h(\sigma(t)-a) + (1-h)(t-a)]^{\beta-1} dh \\ &\leq \beta \int_0^1 [h(\sigma(t)-a) + (1-h)(\sigma(t)-a)]^{\beta-1} dh \\ &= \beta(\sigma(t)-a)^{\beta-1}, \end{aligned}$$

yielding

$$\frac{1}{\beta} \left[ (l-a)^\beta - (b-a)^\beta \right] \leq \int_b^l (\sigma(t)-a)^{\beta-1} \Delta t. \quad (2.9)$$

Finally, combining inequalities (2.8) and (2.9) yields the desired result.

(ii). For the case  $\beta < 1$ , the proof is similar to the proof of (i), except that the inequalities signs are reversed.  $\square$

LEMMA 2.5. Let  $n \in \mathbb{N}$ . If  $0 \leq x_i \leq y_i$ , for  $1 \leq i \leq n$ . Then

$$\prod_{i=1}^n (y_i - x_i) \leq \prod_{i=1}^n y_i - \prod_{i=1}^n x_i. \quad (2.10)$$

*Proof.* The proof is performed by induction and just noting that

$$\begin{aligned} (y_2 - x_2)(y_1 - x_1) &= y_2 y_1 - x_2 x_1 - x_2(y_1 - x_1) - x_1(y_2 - x_2) \\ &\leq y_2 y_1 - x_2 x_1. \quad \square \end{aligned}$$

### 3. Multidimensional Hardy-type inequalities for convex functions on time scales

Our first main result reads:

**THEOREM 3.1.** *Let  $0 \leq a < b < \infty$ . Let the functions  $p, \beta : [\mathbf{a}, \mathbf{b}]_{\mathbb{T}} \rightarrow \mathbb{R}$  be defined, respectively, by*

$$p(\mathbf{x}) = \begin{cases} p_o, & \mathbf{0} \leq \mathbf{x} \leq \mathbf{b}, \\ p_1, & \mathbf{x} > \mathbf{b}, \end{cases} \quad \text{and } \beta(x) = \begin{cases} \beta_o, & 0 \leq x \leq b, \\ \beta_1, & x > b. \end{cases} \quad (3.1)$$

Moreover, assume that  $p_o, p_1 \in \mathbb{R} \setminus \{0\}$  are such that  $p_o \geq 1, p_1 \geq 1$  or  $p_o \geq 1, p_1 < 0$  or  $p_o < 0, p_1 \geq 1$  or  $p_o < 0, p_1 < 0$ . If  $f : [\mathbf{a}, \mathbf{l}] \rightarrow \mathbb{R}$  is non-negative  $\Delta$ -integrable and  $f \in C_{rd}([\mathbf{a}, \mathbf{l}], \mathbb{R})$  for which

$$\int_{a_1}^{b_1} \dots \int_{a_n}^{b_n} f^{p(\mathbf{x})}(y_1, \dots, y_n) \left( \prod_{i=1}^n \frac{1}{\beta(x)} ((y_i - a_i)^{-\beta(x)}) \right) \times \left[ 1 - \prod_{i=1}^n \left( \frac{y_i - a_i}{l_i - a_i} \right)^{\beta(x)} \right] \Delta y_1 \dots \Delta y_n < \infty, \quad (3.2)$$

then

$$\begin{aligned} & \int_{a_1}^{l_1} \dots \int_{a_n}^{l_n} \left( \frac{1}{\prod_{i=1}^n (\sigma(x_i) - a_i)} \int_{a_1}^{\sigma(x_1)} \dots \int_{a_1}^{\sigma(x_1)} f(y_1, \dots, y_n) \Delta y_1 \dots \Delta y_n \right)^{p(\mathbf{x})} \\ & \times \prod_{i=1}^n (\sigma(x_i) - a_i)^{-\beta(x)} \Delta x_1 \dots \Delta x_n \\ & \leq \int_{a_1}^{b_1} \dots \int_{a_n}^{b_n} f^{p(\mathbf{x})}(y_1, \dots, y_n) \left( \prod_{i=1}^n \frac{1}{\beta(x)} ((y_i - a_i)^{-\beta(x)}) \right) \\ & \times \left[ 1 - \prod_{i=1}^n \left( \frac{y_i - a_i}{l_i - a_i} \right)^{\beta(x)} \right] \Delta y_1 \dots \Delta y_n + I_o, \end{aligned} \quad (3.3)$$

where  $I_o = 0$  if  $l \leq b$  (so that  $\beta(x) = \beta_o$  and  $p(x) = p_o$ ) and

$$\begin{aligned} I_o = & \int_{a_1}^{b_1} \dots \int_{a_n}^{b_n} f^{p_1}(y_1, \dots, y_n) \prod_{i=1}^n \left[ \frac{1}{\beta_1} ((y_i - a_i)^{-\beta_1} - (l_i - a_i)^{-\beta_1}) \right] \Delta y_1 \dots \Delta y_n \\ & - \int_{a_1}^{b_o} \dots \int_{a_n}^{b_n} f^{p_o}(y_1, \dots, y_n) \prod_{i=1}^n \left[ \frac{1}{\beta_o} ((y_i - a_i)^{-\beta_o} - (l_i - a_i)^{-\beta_o}) \right] \Delta y_1 \dots \Delta y_n. \end{aligned} \quad (3.4)$$

If  $0 < p(x) \leq 1$ , then (3.3) holds in the reverse direction.

*Proof.* Let  $b \geq l$ . By applying Jensen's inequality (see Remark 2.3), Lemma 2.1 and Lemma 2.4, we find that

$$\begin{aligned}
& \int_{a_1}^{l_1} \dots \int_{a_n}^{l_n} \left( \frac{1}{\prod_{i=1}^n (\sigma(x_i) - a_i)} \int_{a_1}^{\sigma(x_1)} \dots \int_{a_1}^{\sigma(x_1)} f(y_1, \dots, y_n) \Delta y_1 \dots \Delta y_n \right)^{p(\mathbf{x})} \\
& \quad \times \prod_{i=1}^n (\sigma(x_i) - a_i)^{-\beta(x)} \Delta x_1 \dots \Delta x_n \\
& \leq \int_{a_1}^{l_1} \dots \int_{a_n}^{l_n} \left[ \frac{1}{\prod_{i=1}^n (\sigma(x_i) - a_i)} \int_{a_1}^{\sigma(x_1)} \dots \int_{a_1}^{\sigma(x_1)} f^{p_o}(y_1, \dots, y_n) \Delta y_1 \dots \Delta y_n \right] \\
& \quad \times \prod_{i=1}^n (\sigma(x_i) - a_i)^{-\beta_o} \Delta x_1 \dots \Delta x_n \\
& \leq \int_{a_1}^{l_1} \dots \int_{a_n}^{l_n} f^{p_o}(y_1, \dots, y_n) \left[ \int_{y_1}^{l_1} \dots \int_{y_n}^{l_n} \prod_{i=1}^n (\sigma(x_i) - a_i)^{-(\beta_o+1)} \Delta x_1 \dots \Delta x_n \right] \\
& \quad \times \Delta y_1 \dots \Delta y_n \\
& \leq \int_{a_1}^{b_1} \dots \int_{a_n}^{b_n} f^{p(x)}(y_1, \dots, y_n) \left( \prod_{i=1}^n \frac{1}{\beta(x)} ((y_i - a_i)^{-\beta(x)}) \right) \\
& \quad \times \left[ 1 - \prod_{i=1}^n \left( \frac{y_i - a_i}{l_i - a_i} \right)^{\beta(x)} \right] \Delta y_1 \dots \Delta y_n.
\end{aligned}$$

Hence, (3.3) is proved for this case.

Next, let  $b \leq l$ . By applying Jensen's inequality (see Remark 2.3) and Lemma 2.1, we find that

$$\begin{aligned}
& \int_{a_1}^{l_1} \dots \int_{a_n}^{l_n} \left( \frac{1}{\prod_{i=1}^n (\sigma(x_i) - a_i)} \int_{a_1}^{\sigma(x_1)} \dots \int_{a_1}^{\sigma(x_1)} f(y_1, \dots, y_n) \Delta y_1 \dots \Delta y_n \right)^{p(\mathbf{x})} \\
& \quad \times \prod_{i=1}^n (\sigma(x_i) - a_i)^{-\beta(x_i)} \Delta x_1 \dots \Delta x_n \\
& \leq \int_{a_1}^{b_1} \dots \int_{a_n}^{b_n} \left( \frac{1}{\prod_{i=1}^n (\sigma(x_i) - a_i)} \int_{a_1}^{\sigma(x_1)} \dots \int_{a_1}^{\sigma(x_1)} f(y_1, \dots, y_n) \Delta y_1 \dots \Delta y_n \right)^{p_o} \\
& \quad \times \prod_{i=1}^n (\sigma(x_i) - a_i)^{-\beta_o} \Delta x_1 \dots \Delta x_n
\end{aligned}$$

$$\begin{aligned}
& + \int_{b_1}^{l_1} \dots \int_{b_n}^{l_n} \left( \frac{1}{\prod_{i=1}^n (\sigma(x_i) - a_i)} \int_{a_1}^{b_1} \dots \int_{a_n}^{b_n} f(y_1, \dots, y_n) \Delta y_1 \dots \Delta y_n \right)^{p_1} \\
& \times \prod_{i=1}^n (\sigma(x_i) - a_i)^{-\beta_1} \Delta x_1 \dots \Delta x_n \\
& + \int_{b_1}^{l_1} \dots \int_{b_n}^{l_n} \left( \frac{1}{\prod_{i=1}^n (\sigma(x_i) - a_i)} \int_{b_1}^{\sigma(x_1)} \dots \int_{b_n}^{\sigma(x_n)} f(y_1, \dots, y_n) \Delta y_1 \dots \Delta y_n \right)^{p_1} \\
& \times \prod_{i=1}^n (\sigma(x_i) - a_i)^{-\beta_1} \Delta x_1 \dots \Delta x_n \\
& \leqslant \int_{a_1}^{b_1} \dots \int_{a_n}^{b_n} f^{p_o}(y_1, \dots, y_n) \left( \int_{y_1}^{b_1} \dots \int_{y_n}^{b_n} \prod_{i=1}^n (\sigma(x_i) - a_i)^{-\beta_o} \Delta x_1 \dots \Delta x_n \right) \Delta y_1 \dots \Delta y_n \\
& + \int_{a_1}^{b_1} \dots \int_{a_n}^{b_n} f^{p_1}(y_1, \dots, y_n) \left( \int_{b_1}^{l_1} \dots \int_{b_n}^{l_n} \prod_{i=1}^n (\sigma(x_i) - a_i)^{-\beta_o} \Delta x_1 \dots \Delta x_n \right) \Delta y_1 \dots \Delta y_n \\
& + \int_{b_1}^{l_1} \dots \int_{b_n}^{l_n} f^{p_1}(y_1, \dots, y_n) \left( \int_{y_1}^{l_1} \dots \int_{y_n}^{l_n} \prod_{i=1}^n (\sigma(x_i) - a_i)^{-\beta_o} \Delta x_1 \dots \Delta x_n \right) \Delta y_1 \dots \Delta y_n \\
& := I. \tag{3.5}
\end{aligned}$$

By Lemma 2.4 and Lemma 2.5, we find that

$$\begin{aligned}
I & \leqslant \int_{a_1}^{b_1} \dots \int_{a_n}^{b_n} f^{p_o}(y_1, \dots, y_n) \prod_{i=1}^n \left[ \frac{1}{\beta_o} \left( (y_i - a_i)^{-\beta_o} - (b_i - a_i)^{-\beta_o} \right) \right] \Delta y_1 \dots \Delta y_n \\
& + \int_{a_1}^{b_1} \dots \int_{a_n}^{b_n} f^{p_1}(y_1, \dots, y_n) \prod_{i=1}^n \left[ \frac{1}{\beta_1} \left( (b_i - a_i)^{-\beta_1} - (l_i - a_i)^{-\beta_1} \right) \right] \Delta y_1 \dots \Delta y_n \\
& + \int_{b_1}^{l_1} \dots \int_{b_n}^{l_n} f^{p_1}(y_1, \dots, y_n) \prod_{i=1}^n \left[ \frac{1}{\beta_1} \left( (y_i - a_i)^{-\beta_1} - (l_i - a_i)^{-\beta_1} \right) \right] \Delta y_1 \dots \Delta y_n \\
& \leqslant \int_{a_1}^{b_1} \dots \int_{a_n}^{b_n} f^{p_o}(y_1, \dots, y_n) \left( \prod_{i=1}^n \frac{1}{\beta_o} \left( (y_i - a_i)^{-\beta_o} \right) \right) \left[ 1 - \prod_{i=1}^n \left( \frac{y_i - a_i}{l_i - a_i} \right)^{\beta_o} \right] \\
& \cdot \Delta y_1 \dots \Delta y_n \\
& + \int_{b_1}^{l_1} \dots \int_{b_n}^{l_n} f^{p_1}(y_1, \dots, y_n) \left( \prod_{i=1}^n \frac{1}{\beta_1} \left( (y_i - a_i)^{-\beta_1} \right) \right) \left[ 1 - \prod_{i=1}^n \left( \frac{y_i - a_i}{l_i - a_i} \right)^{\beta_1} \right] \\
& \cdot \Delta y_1 \dots \Delta y_n \\
& + \int_{a_1}^{b_1} \dots \int_{a_n}^{b_n} f^{p_1}(y_1, \dots, y_n) \prod_{i=1}^n \left[ \frac{1}{\beta_1} \left( (y_i - a_i)^{-\beta_1} - (l_i - a_i)^{-\beta_1} \right) \right] \Delta y_1 \dots \Delta y_n
\end{aligned}$$

$$\begin{aligned}
& - \int_{a_1}^{b_o} \dots \int_{a_n}^{b_n} f^{p_o}(y_1, \dots, y_n) \prod_{i=1}^n \left[ \frac{1}{\beta_o} \left( (y_i - a_i)^{-\beta_1} - (l_i - a_i)^{-\beta_o} \right) \right] \Delta y_1 \dots \Delta y_n \\
& = \int_{a_1}^{b_1} \dots \int_{a_n}^{b_n} f^{p(x)}(y_1, \dots, y_n) \left( \prod_{i=1}^n \frac{1}{\beta(x)} \left( (y_i - a_i)^{-\beta(x)} \right) \right) \\
& \quad \times \left[ 1 - \prod_{i=1}^n \left( \frac{y_i - a_i}{l_i - a_i} \right)^{\beta(x)} \right] \Delta y_1 \dots \Delta y_n + I_o.
\end{aligned} \tag{3.6}$$

By combining the inequalities (3.5) with (3.6) the inequality (3.3) follows so that the proof is complete.  $\square$

The next result concerns the dual version of Theorem 3.1 when the Hardy operator

$$H : f(x_1, \dots, x_n) \longrightarrow \frac{1}{\prod_{i=1}^n (\sigma(x_i) - a_i)} \int_{a_1}^{\sigma(x_1)} \dots \int_{a_1}^{\sigma(x_1)} f(y_1, \dots, y_n) \Delta y_1 \dots \Delta y_n$$

is replaced by the dual Hardy operator

$$H^* : f(x_1, \dots, x_n) \longrightarrow \prod_{i=1}^n (\sigma(x_i) - a_i) \int_{\sigma(x_1)}^{\infty} \dots \int_{\sigma(x_n)}^{\infty} \frac{f(y_1, \dots, y_n) \Delta y_1 \dots \Delta y_n}{\prod_{i=1}^n (\sigma(y_i) - a_i) (y_i - a_i)}.$$

Our next main result concerning the dual Hardy operator  $H^*$  reads:

**THEOREM 3.2.** *Let  $0 \leq a < b < \infty$ . Let the functions  $p, \beta : [\mathbf{a}, \mathbf{b}]_{\mathbb{T}} \rightarrow \mathbb{R}$  be defined, respectively, by*

$$p(\mathbf{x}) = \begin{cases} p_o, & \mathbf{0} \leq \mathbf{x} \leq \mathbf{b}, \\ p_1, & \mathbf{x} > \mathbf{b}, \end{cases} \quad \beta(x) = \begin{cases} \beta_o, & 0 \leq x \leq b, \\ \beta_1, & x > b. \end{cases} \tag{3.7}$$

Moreover, assume that  $p_o, p_1 \in \mathbb{R} \setminus \{0\}$  are such that  $p_o \geq 1, p_1 \geq 1$  or  $p_o \geq 1, p_1 < 0$  or  $p_o < 0, p_1 \geq 1$  or  $p_o < 0, p_1 < 0$ . If  $f : [\mathbf{a}, \mathbf{l}] \rightarrow \mathbb{R}$  is non-negative  $\Delta$ -integrable and  $f \in C_{rd}([\mathbf{a}, \mathbf{l}], \mathbb{R})$  for which

$$\begin{aligned}
& \int_{l_1}^{\infty} \dots \int_{l_n}^{\infty} f^{p_1}(y_1, \dots, y_n) \left( \prod_{i=1}^n \frac{(y_i - a_i)^{\beta(y)}}{\beta(y)} \right) \left[ 1 - \prod_{i=1}^n \left( \frac{l_i - a_i}{y_i - a_i} \right)^{\beta(y)} \right] \\
& \quad \times \frac{\Delta y_1 \dots \Delta y_n}{\prod_{i=1}^n (\sigma(y_i) - a_i) (y_i - a_i)} < \infty,
\end{aligned} \tag{3.8}$$

then

$$\int_{l_1}^{\infty} \dots \int_{l_n}^{\infty} \left( \prod_{i=1}^n (\sigma(x_i) - a_i) \int_{\sigma(x_1)}^{\infty} \dots \int_{\sigma(x_n)}^{\infty} \left( \frac{f(y_1, \dots, y_n)}{\prod_{i=1}^n (\sigma(y_i) - a_i) (y_i - a_i)} \right) \Delta y_1 \dots \Delta y_n \right)^{p(x)} \Delta x_1 \dots \Delta x_n$$

$$\begin{aligned}
& \times \left( \prod_{i=1}^n (x_i - a_i) \right)^{\beta(x)-1} \frac{\Delta x_1 \dots \Delta x_n}{\prod_{i=1}^n (\sigma(x_i) - a_i)} \\
& \leq \int_{l_1}^{\infty} \dots \int_{l_n}^{\infty} f^{p_1}(y_1, \dots, y_n) \left( \prod_{i=1}^n \frac{(y_i - a_i)^{\beta(y)}}{\beta(y)} \right) \left[ 1 - \prod_{i=1}^n \left( \frac{l_i - a_i}{y_i - a_i} \right)^{\beta(y)} \right] \\
& \quad \times \frac{\Delta y_1 \dots \Delta y_n}{\prod_{i=1}^n (\sigma(y_i) - a_i)(y_i - a_i)} + I_o. \tag{3.9}
\end{aligned}$$

where  $I_o = 0$  if  $l \leq b$  (so that  $\beta(x) = \beta_o$  and  $p(x) = p_o$ ) and

$$\begin{aligned}
I_o &= \int_{l_1}^{\infty} \dots \int_{l_n}^{\infty} \frac{f^{p_1}(y_1, \dots, y_n)}{\prod_{i=1}^n (\sigma(y_i) - a_i)(y_i - a_i)} \prod_{i=1}^n \left[ \frac{1}{\beta_1} \left( (y_i - a_i)^{\beta_1} - (l_i - a_i)^{\beta_1} \right) \right] \Delta y_1 \dots \Delta y_n \\
&\quad - \int_{l_1}^{\infty} \dots \int_{l_n}^{\infty} \frac{f^{p_o}(y_1, \dots, y_n)}{\prod_{i=1}^n (\sigma(y_i) - a_i)(y_i - a_i)} \prod_{i=1}^n \left[ \frac{1}{\beta_o} \left( (y_i - a_i)^{\beta_o} - (l_i - a_i)^{\beta_o} \right) \right] \Delta y_1 \dots \Delta y_n. \tag{3.10}
\end{aligned}$$

If  $0 < p(x) \leq 1$ , then (3.9) holds in the reverse direction.

*Proof.* Let  $b \geq l$ . Applying Jensen's inequality (see Remark 2.3) and Lemma 2.1, we obtain that

$$\begin{aligned}
& \int_{l_1}^{\infty} \dots \int_{l_n}^{\infty} \left( \prod_{i=1}^n (\sigma(x_i) - a_i) \int_{\sigma(x_1)}^{\infty} \dots \int_{\sigma(x_n)}^{\infty} \left( \frac{f(y_1, \dots, y_n)}{\prod_{i=1}^n (\sigma(y_i) - a_i)(y_i - a_i)} \right) \Delta y_1 \dots \Delta y_n \right)^{p(x)} \\
& \quad \times \left( \prod_{i=1}^n (x_i - a_i) \right)^{\beta(x)-1} \frac{\Delta x_1 \dots \Delta x_n}{\prod_{i=1}^n (\sigma(x_i) - a_i)} \\
& \leq \int_{l_1}^{\infty} \dots \int_{l_n}^{\infty} \int_{\sigma(x_1)}^{\infty} \dots \int_{\sigma(x_n)}^{\infty} \left( \frac{f^{p_1}(y_1, \dots, y_n) \Delta y_1 \dots \Delta y_n}{\prod_{i=1}^n (\sigma(y_i) - a_i)(y_i - a_i)} \right) \\
& \quad \times \left( \prod_{i=1}^n (x_i - a_i) \right)^{\beta_1-1} \Delta x_1 \dots \Delta x_n \\
& \leq \int_{l_1}^{\infty} \dots \int_{l_n}^{\infty} \frac{f^{p_1}(y_1, \dots, y_n)}{\prod_{i=1}^n (\sigma(y_i) - a_i)(y_i - a_i)} \left[ \int_{l_1}^{y_1} \dots \int_{l_n}^{y_n} \left( \prod_{i=1}^n (x_i - a_i) \right)^{\beta_1-1} \Delta x_1 \dots \Delta x_n \right]
\end{aligned}$$

$$\begin{aligned}
& \times \Delta y_1 \dots \Delta y_n \\
& \leq \int_{l_1}^{\infty} \dots \int_{l_n}^{\infty} f^{p_1}(y_1, \dots, y_n) \left( \prod_{i=1}^n \frac{(y_i - a_i)^{\beta(y)}}{\beta(y)} \right) \left[ 1 - \prod_{i=1}^n \left( \frac{l_i - a_i}{y_i - a_i} \right)^{\beta(y)} \right] \\
& \quad \times \frac{\Delta y_1 \dots \Delta y_n}{\prod_{i=1}^n (\sigma(y_i) - a_i)(y_i - a_i)}. \tag{3.11}
\end{aligned}$$

Finally, let  $b \leq l$ . Also the proof of this case is completely analogous to the corresponding part of the proof of Theorem 3.1 so we leave out the details.  $\square$

## REFERENCES

- [1] R. P. AGARWAL, M. BOHNER AND N. PETERSON, *Inequalities on time scales: a survey*, Math. Inequal. Appl. 4 (2001), 535–557.
- [2] R. P. AGARWAL, D. O'REGAN AND S. H. SAKAR, *Dynamic inequalities on time scales*, Springer, Heidelberg/New York/Dordrecht/London, 2014.
- [3] R. P. AGARWAL, D. O'REGAN AND S. H. SAKAR, *Hardy type inequalities on time scales*, Springer International Publishing, Switzerland, 2016.
- [4] M. ANWAR, R. BIBI, M. BOHNER AND J. E. PEČARIĆ, *Jensen's functional on time scales for several variables*, Int. J. Anal. 2014 (2014), 14pp.
- [5] R. BIBI, M. BOHNER, J. E. PEČARIĆ AND S. VAROŠANEC, *Minkowski and Beckenbach-Dresher inequalities and functionals on time scales*, J. Math. Inequal. 7 (2013), no. 3, 299–312.
- [6] M. BOHNER AND A. PETERSON (EDS.), *Dynamic equations on time scales: An Introductin with Applications*, Birkhäuser, Boston, Massachusetts, 2001.
- [7] M. BOHNER AND A. PETERSON (EDS.), *Advances in Dynamic equations on time scales*, Birkhäuser, Boston, Massachusetts, 2003.
- [8] T. DONCHEV, A. NOSHEEN AND J. E. PEČARIĆ, *Hardy-Type inequalities on time scales via convexity in several variables*, ISRN Math. Anal. 2013 (2013), 9pp.
- [9] G. H. HARDY, *Notes on a theorem of Hilbert*, Math. Z. 6 (1920), 314–317.
- [10] G. H. HARDY, *Notes on some points in the integral calculus, LX. An inequality between integrals*, Messenger Math. 54 (1925), 150–156.
- [11] G. H. HARDY, J. E. LITTLEWOOD AND G. PÓLYA, *Inequalities*, Cambridge University Press, Cambridge, 1959.
- [12] S. HILGER, *Ein Maßkettenkalkül mit Anwendung auf Zentrumsmannigfaltigkeiten*, PhD thesis, Universität Würzburg, 1988.
- [13] S. HILGER, *Differential and difference calculus*, Nonlinear Anal., Proceedings of Second World Congress of Nonlinear Analysts, **30** (5), (1997), 2683–2694.
- [14] S. HILGER, *Analysis on measure chains-a unified approach to continuous and discrete calculus*, Results Math. **18** (1990), no. 1–2, 18–56.
- [15] A. KUFNER, L. MALIGRANDA AND L. - E. PERSSON, *The prehistory of the Hardy inequality*, Amer. Math. Monthly 113 (2006), no. 10, 715–732.
- [16] A. KUFNER, L. MALIGRANDA AND L. - E. PERSSON, *The Hardy Inequality. About its History and Some Related Results*, Vydavatelsky Servis Publishing House, Pilsen, 2007.
- [17] A. KUFNER, L. - E. PERSSON AND N. SAMKO, *Weighted Inequalities of Hardy type*, Second Edition, World Scientific Publishing Co. Inc., River Edge, New Jersey, 2017.

- [18] J. A. OGUNTUASE AND L. E. PERSSON, *Time scales Hardy-type inequalities via superquadraticity*, Ann. Funct. Anal. 5 (2014), no.2, 61–73.
- [19] P. ŘEHÁK, *Hardy inequality on time scales and its application to half-linear dynamic equations*, J. Inequal. Appl. 5 (2005), 495–507.

(Received November 12, 2018)

O. O. Fabelurin

*Department of Mathematics  
Obafemi Awolowo University  
Ile-Ife, Osun State, Nigeria  
e-mail: fabelur@yahoo.com*

J. A. Oguntuase

*Department of Mathematics  
Federal University of Agriculture  
P.M.B. 2240, Abeokuta, Ogun State, Nigeria  
e-mail: oguntuase@yahoo.com*

L.-E. Persson

*UiT, The Arctic University of Norway  
Campus Narvik, Lødve Langes Gate 2, 8514 Narvik, Norway  
e-mail: larserik6pers@gmail.com*