

Maternal age and risk of cesarean section in women with induced labor at term - a Nordic register-based study

Thomas BERGHOLT¹, Finn Egil SKJELDESTAD², Aura PYYKÖNEN³, Steen C. RASMUSSEN¹, Anna-Maija TAPPER^{4,5}, Ragnheiður I. BJARNADÓTTIR⁶⁻⁸, Alexander SMÁRASON⁷, Birna B. MÁSDÓTTIR⁸, Kari KLUNGSØYR^{9,10}, Susanne ALBRECHTSEN^{10,11}, Karin KÄLLÉN^{12,13}, Mika GISSLER^{14,15}, Ellen Christine Leth LØKKEGAARD¹

1. Clinical Research Unit, Department of Obstetrics and Gynecology, North Zealand Hospital, University of Copenhagen, Hillerød, Denmark.
2. Department of Community Medicine, Faculty of Health Sciences, UiT the Arctic University of Norway, Tromsø, Norway
3. Department of Obstetrics and Gynecology, Helsinki University Hospital, Helsinki, Finland
4. Hyvinkää Hospital, Hyvinkää, Finland
5. University of Helsinki, Helsinki, Finland
6. The Icelandic Birth Registry, Akureyri, Iceland
7. Institute of Health Science Research, University of Akureyri, Akureyri, Iceland
8. Landspítali University Hospital, Reykjavik, Iceland
9. Division of Mental and Physical Health, Norwegian Institute of Public Health, Bergen, Norway
10. Department of Global Public Health and Primary Care, University of Bergen, Bergen, Norway
11. Haukeland University Hospital, Bergen, Norway

12. Swedish National Board of Health and Welfare, Stockholm, Sweden
13. University of Lund, Lund, Sweden
14. Finnish Institute for Health and Welfare THL, Helsinki, Finland
15. Department of Neurobiology, Care Sciences and Society, Karolinska Institute, Stockholm, Sweden

Corresponding Author:

Ellen Løkkegaard

Department of Obstetrics and Gynecology, North Zealand Hospital, University of Copenhagen,
Dyrehavevej 29, 3400 Hillerød, Denmark

Email: Ellen.Christine.Leth.Loekkegaard@regionh.dk

Conflicts of interest

None

Funding

The study has been funded by the Nordic Federation of Societies of Obstetrics and Gynecology (NFOG) Fund.

ABSTRACT

Introduction: Over the last decades, induction of labor has increased in many countries along with increasing maternal age. We assessed the effects of maternal age and labor induction on cesarean section at term among nulliparous and multiparous women without previous cesarean section. **Material and methods:** We performed a retrospective national registry-based study from Denmark, Finland, Iceland, Norway and Sweden including 3 398 586 deliveries between 2000 and 2011. We investigated the impact of age on cesarean section among 196 220 nulliparous and 188 158 multiparous women whose labor was induced, had single cephalic presentation at term and no previous cesarean section. Confounders comprised country, time-period and gestational age. **Results:** In nulliparous women with induced labor the rate of cesarean section increased from 14.0% in women less than 20 years of age to 39.9% in women 40 years and older. Compared to women aged 25-29 years, the corresponding relative risk were 0.60 (95% confidence interval (CI); 0.57 to 0.64) and 1.72 (95% CI; 1.66 to 1.79). In multiparous induced women the risk of cesarean section was 3.9% in women less than 20 years rising to 9.1% in women 40 years and older. Compared to women aged 25-29 years, the relative risk were 0.86 (95% CI; 0.54 to 1.37) and 1.98 (95% CI; 1.84 to 2.12), respectively. There were minimal confounding effects of country, time-period and gestational age on risk for cesarean section. **Conclusions:** Advanced maternal age is associated with increased risk of cesarean section in women undergoing labor induction with a single cephalic presentation at term without a previous cesarean section. The absolute risk of cesarean section is 3-5 times higher across 5-year age groups in nulliparous relative to multiparous women having induced labor.

Abbreviations:

CS cesarean section

CI confidence interval

RR relative risk

Keywords

Maternal age, induction of labor, cesarean section, birth register

Key message

Induction and cesarean section rates increased by age in both nulli- and multiparous women in all Nordic countries over the years 2001 through 2011.

For Peer Review

INTRODUCTION

For many decades maternal age at childbirth has increased in Europe (1) and the USA (2). Data from Nordic Perinatal Statistics confirms the international trends of increasing maternal age in all the Nordic countries (3). Along with increasing maternal age, the overall rate of labor inductions has doubled from year 2000 to 2012 in Denmark (4) and from 2003 to 2013 in Norway (5). In Sweden, induction of post-term pregnancies increased from 28% in 1992 to 46% in 2006 (4).

A systematic review of randomized controlled trials comparing post-term spontaneous onset of labor with induction at or beyond term concluded on a significantly decreased risk of perinatal death as well as a reduction in cesarean section (CS) among women having induced labor (5). Furthermore, randomized controlled trials have found induction of labor associated with improved maternal outcomes for women with mild hypertensive disorders beyond 37 weeks of gestation (6), and with reduced risk of shoulder dystocia and related morbidity in large-for-dates fetuses compared to expectant management (7).

There is a general concern about increasing CS rates world-wide (8). Delivery by CS has implication on postoperative maternal morbidity and complications in subsequent pregnancies (9, 10). The World Health Organization has recently re-evaluated their strategy regarding the optimal CS rate. They state that CS should only be performed in pregnancies that benefit fetal and maternal health, and this may vary around the world according to obstetric competence and organization of labor care (11). Physiological changes in the pregnant population including increasing maternal age (1, 12) and body mass index (13, 14), combined with an increasing proportion of women with prior CS (15), and multiple pregnancies (16), have been associated with increasing risk for CS in subsequent pregnancies. CS rates have been reported from 3.5% in Africa to 29.2% in Latin America and the Caribbean (17), and 32.0% in the USA (18). The overall CS rate in the Nordic countries was 17.0% from 2000 to 2011, ranging from 15.7% in Norway to 19.2% in Denmark (15). National variation in CS rates in the Nordic countries was far less than reported elsewhere (19).

The aims of the present study was to assess the effects of maternal age and labor induction on risk of CS at term among nulliparous and multiparous women without previous CS in the Nordic countries over a 12-year time-period.

MATERIAL AND METHODS

The study was initiated by the Nordic Robson Research collaboration (15, 20) based on national medical register data from the five Nordic countries, merged and processed in Statistics Denmark, Copenhagen. Selected variables were harmonized across birth register in the merged database. Maternal age was defined as the difference in completed years between date of delivery and maternal date of birth, and categorized into six groups (< 20, 20–24, 25–29, 30–34, 35–39, ≥ 40 years). Parity and previous CS were defined as the highest number based on information in the register or maternal information provided at first delivery. Gestational age was with few exceptions estimated from ultrasound screening of bi-parietal diameter in second trimester and categorized into three categories (37–38, 39–41, ≥ 42 weeks). Included in the original database were women who delivered after the 22nd gestational week, or, if gestational age was missing, women who gave birth to infants with a birth weight of 500 g or more. Start of delivery was defined as spontaneous onset of birth, induction or CS. Emergency CS's encompassed all CS's not reported as elective. Mode of delivery comprised spontaneous vaginal delivery, instrumental vaginal delivery (vacuum extraction or forceps) or CS. Data on presenting part of the baby at delivery was categorized into cephalic, breech presentation, or transverse lie. We categorized three-year periods as 2000–02, 2003–05, 2006–08 and 2009–11.

We have used the Robson classification system, a method of stratifying delivering women into ten clinically relevant groups, making comparison of interventions and outcomes within groups more clinically relevant. In the Robson system groups are mutually exclusive (21). In this study, we have included Robson groups 1 to 4. Groups 1 and 2 comprise nulliparous women with a baby in single cephalic presentation after 36 weeks of gestation. Group 1 includes women in spontaneous labor (R1) and group 2 women having labor induced (R2a) or pre-labor CS (R2b). Groups 3 and 4 comprise multiparous women with a baby in single cephalic presentation after 36 weeks of gestation and no history of CS. Group 3 includes multiparous women in spontaneous labor (R3) and group 4 multiparous women having labor induced (R4a) or pre-labor CS (R4b)

Statistical analyses

1
2
3 We estimated proportions of nulliparous women having pre-labor CS ($R2b/(R1+R2)$), proportion
4 of women having induction of labor ($R2a/(R1+R2)$), and overall CS rate in nulliparous women
5 ($CS\ in\ (R1+R2a)/(R1+R2)$). Similarly, proportions of multiparous women having pre-labor CS
6 ($R4b/(R3+R4)$), induction of labor ($R4a/(R3+R4)$), and overall CS rate in multiparous women
7 with no history of CS ($CS\ in\ (R3+R4a)/(R3+R4)$) were calculated.
8
9
10

11
12 We performed multiple unconditioned logistics regressions models to estimate odds ratio of CS
13 in induced nulliparous ($R2a$) and multiparous ($R4a$) women, and in nulliparous ($R1$) and
14 multiparous ($R3$) women having spontaneous onset of labor. Maternal age was main exposure,
15 while country, time-period and gestational age were considered confounding factors. Results are
16 presented as proportions (rates) in percentages and in relative risks (RR's) with 95% confidence
17 intervals (CI). IBM SPSS Statistics Version 24 (IBM Corp., Armonk, USA) was used for the
18 statistical analyses.
19
20
21
22
23
24
25
26
27

28 **Ethical approval**

29
30 The Danish Data protection Agency administered the Danish participation in the study (reference
31 NOH-2016-006), and the Regional Committee for Medical and Health Research Ethics South-
32 East C (REK Sør-Øst 2010/3256) assessed the Norwegian participation. The Directorate of
33 Health in Iceland, THL Finnish Institute of Health and Welfare in Finland and the Swedish
34 National Board of Health and Welfare gave their authorizations for using anonymized data.
35
36
37
38
39
40
41

42 **RESULTS**

43
44 From January 2000 to December 2011 there were in total 3 398 586 deliveries in the five Nordic
45 countries. Nulliparous women with a single baby in cephalic presentation at term included 1 294
46 452 mothers ($R1+R2$). Of these 40 572 (3.1%) women had pre-labor CS ($R2b$), 196 220 (15.2%)
47 had labor induced ($R2a$) and 1 057 660 (81.7%) had spontaneous onset of labor ($R1$).
48
49 Multiparous women with a single baby in cephalic presentation at term, with no previous CS,
50 included 1 471 057 mothers ($R3+R4$). Of these 40 885 (2.8%) women had pre-labor CS ($R4b$),
51 188 158 (12.8%) had labor induced ($R4a$) and 1 242 014 (84.4%) spontaneous onset of labor
52 ($R3$).
53
54
55
56
57
58
59
60

1
2
3 Over the 12-year study period the proportion of women having labor induction increased from
4 11.8% to 20.1% in nulliparous and from 9.9% to 16.8% in multiparous women. The increases
5 among nulliparous as well as multiparous women were consistent across countries and age
6 groups.
7
8

9
10 In the multivariate unconditional logistic regression analysis, country, time-period and
11 gestational age had no confounding effect on the age-specific or overall estimates of CS in
12 neither nulliparous nor multiparous women having labor induced (data not presented).

13
14 Consequently, we only presented crude estimates and RR in Table 3. The age-specific estimates
15 of induction on CS were consistent and relatively robust in almost all analyses of time-period
16 and country (no interactions).
17
18
19

20
21 The temporal relations between proportions of women having pre-labor CS, induction of labor,
22 and overall CS rates by parity are displayed in Figure 1. The proportions of nulliparous as well
23 as multiparous women having pre-labor CS were relatively low and stable over time with
24 estimates varying from 2.0 to 3.4%. The overall CS rate in nulliparous women (CS/R1+R2)
25 increased from 12.4% in 2000 to 14.9% in 2006 and remained stable thereafter. In the
26 multiparous women (CS/R3+R4) the CS rate varied from 4.2% in 2000, a peak of 5.6% in 2007,
27 to 5.1% in 2011.
28
29
30
31
32

33
34 Crude rates of CS in induced nulliparous women at term (R2a) by age and overall (R2a) are
35 displayed in Table 1. The overall CS rate was 25.0% increasing consistently from 14% in the
36 youngest to 40% in the oldest age group. There were small variations in the crude CS rates by
37 age and in total between nulliparous women in R2a the five Nordic countries. Age-stratified
38 analysis by time-period and gestational age found consistently increasing CS rates from the
39 youngest to the oldest age group across all subsets of analyses.
40
41
42
43
44

45
46 In the same manner, number of women under risk and crude rates of CS in induced multiparous
47 women with no previous CS (Robson 4a) are shown in Table 2. The overall CS rate in the
48 Robson 4a group was 5.5%. The risk of CS increased with age in all countries. Finland had
49 lower age-specific CS rates and a lower overall crude CS compared to the other Nordic
50 countries. Moreover, the crude CS rates ranged from 2.4% (Finland) to 4.7% (Iceland) in women
51 20-24 years of age, and from 5.2% (Finland) and 10.9% (Sweden) in women 40 years or older.
52
53
54
55

56
57 Across age groups, the RR of CS in nulliparous women with induced labor (R2a) increased
58 significantly by a five-fold from 0.60 (95% CI: 0.57-0.64) in women below 20 years of age to
59
60

1
2
3 1.72 (95% CI:1.66-1.79) in women 40 years or above (Table 3). The same pattern was observed
4 in nulliparous women with spontaneous onset of labor (R1) with RRs increasing from 0.57 (95%
5 CI:0.55-0.60) in the youngest group to 2.50 (95% CI: 2.37-2.64) in women 40 years or older.
6
7

8
9 In the 432 multiparous women, younger than 20 years, with induced labor (R4a) the crude CS
10 rate was 3.9%. The CS rate increased from 3.2% in women aged 20-24 years to 9.1% in women
11 40 years or above. The RR increased from 0.70 (95% CI: 0.63-0.77) in women 20-24 years to
12 1.98 (95% CI: 1.84-2.12) in women 40 years or above.
13
14

15
16 The absolute risk of CS in both nulliparous and multiparous women with induced labor
17 (R2a/R4a) compared to the respective women with spontaneous onset of labor (R1/R3) was 1.90
18 to 2.98 times higher among nulliparous and 2.28 to 2.80 higher among nulliparous women.
19 (Table 3).
20
21
22
23
24
25

26 **DISCUSSION**

27
28
29 The main findings from this study were an increasing incidence of induction of labor and stable
30 rates of CS before onset of labor in the Nordic countries, from 2000 to 2011 in both nulliparous
31 and multiparous women, with a single baby in single cephalic presentation at term, and no
32 previous CS. In addition, we found that increasing maternal age was associated with increasing
33 RR of CS in both nulliparous and multiparous women. The RR of CS was 2-3 times higher in
34 each age group among women having induced relative spontaneous onset of labor. These
35 estimates enable Nordic obstetricians and midwives to inform pregnant women at term, with a
36 single baby in cephalic presentation and no previous CS, about their average risk of CS if they
37 are having labor induced. These
38
39
40
41
42
43
44

45
46 Furthermore, we found an association between rising induction rates to 15% in nulliparous
47 women (R1+R2) and 13% in multiparous women (R3+R4) from 2000 to 2007 and an increasing
48 risk of overall CS to 15.0% and 5.1% respectively during the same period. When the induction
49 rates increased further to 20.1% and 16.8% in 2011, the respective overall CS rates remained
50 stable at 14.9% and 5.1%. These findings indicate that the risk of CS might be unaffected above
51 a certain proportion of women having labor induced among nulliparous as well as multiparous
52 women without a previous CS. Our findings among nulliparous women is supported by the
53 findings of a large randomized controlled trial of induction of labor in 6 104 American low-risk
54
55
56
57
58
59
60

1
2
3 women between 39+0 and 39+4 days of gestation, the ARRIVE study (22). The authors found a
4 decreased RR of CS (RR 0.84 (95% CI:0.76-0.93)) in women randomized to labor induction
5 compared to women awaiting for spontaneous onset of labor. Low risk nulliparous women are
6 currently not having labor induced before term in the Nordic countries, but if they are going to
7 be in the future, the results from the ARRIVE suggests that there may be an associated reduced
8 overall CS rate.
9
10
11
12

13
14 A major strength of the study is the large sample size with national data from the five Nordic
15 countries. Limitations of the study are the observational design combined with lack of data for
16 body mass index, more detailed information of indications for induction, and the Bishop score at
17 time for induction. However, a systematic review from 2013 concluded that Bishop score is a
18 poor predictor of outcome of induced labor including risk of CS, and the authors advocated not
19 to use Bishop score to decide whether to induce labor or not (23).
20
21
22
23
24

25 Cheng and colleagues published an observational study of 442 003 low risk nulliparous women
26 with a baby in single cephalic presentation delivering between 39 and 42 weeks of gestation
27 comparing induction of labor with expectant management (23). They found a lower risk of CS
28 (OR = 0.90, (95% CI 0.88 to 0.91)) if induced at 39 weeks of gestation compared to expectant
29 management (24). In their low-risk nulliparous population the proportion aged 35 years or more,
30 was 7.3% compared to 13.4% in our population. As high maternal age is associated with CS and
31 assisted vaginal delivery (25, 26), these findings underline the importance of maternal age
32 differences in study populations when comparing results of obstetric operative intervention in
33 obstetrics across studies. In a randomized controlled trial of induction of labor relative expectant
34 management in nulliparous women 35 years or older no significant differences in CS rates were
35 found (32.2% relative 32.8%) (27). Their CS rate in the induction group is similar with the CS
36 rate of 34.3% in nulliparous women 35 to 39 years of age in the present study. In a meta-analysis
37 of randomized controlled trials of induction of labor at full term in uncomplicated singleton
38 gestations, Saccone and Berghella reported a CS rate of 25.5% among nulliparous women with
39 labor induced (28) similar to our findings (25.0%).
40
41
42
43
44
45
46
47
48
49
50

51
52 Beside the increased use of induction of labor in post-term and hypertensive pregnancies, other
53 factors as institutional culture in obstetric practice and growing patient involvement in the
54 decision-making process can influence the use of induction of labor. In a qualitative study,
55 midwives and doctors from ten Australian hospitals expressed a large variation in clinical
56 decision-making for induction of labor. (29). In addition, the authors found that hospitals having
57
58
59
60

1
2 senior midwives or midwife managers, serving as gatekeepers with the authority to question the
3 indication for induction, had lower rates of induction of labor. A Dutch interview study of
4 doctors and midwives found that medical indications as pre-eclampsia and post-term pregnancy
5 were the most important factors in the decision for induction of labor (30). Together, these
6 findings indicate the need for continuous audit of the rates and indications for induction of labor
7 and CS at each delivery unit. This is important for surveillance of institutional standards
8 regarding induction of labor and to provide institutional data for information to the pregnant
9 women.
10
11
12
13
14
15
16
17
18
19

20 CONCLUSION

21
22 The risk of CS increases with increasing maternal age after both induced and spontaneous onset
23 of labor in nulliparous and multiparous women without previous CS. In all age groups, the risk
24 of CS is considerably higher following induced labor. The findings provide a knowledge basis
25 which can help to inform women at term about their average risk of CS associated with induction
26 of labor and help these women in the decision-making process. Women are entitled to be
27 informed of benefits and risks related to the procedure, including risk of CS, when induction of
28 labor is considered.
29
30
31
32
33
34
35
36
37

38 References

- 39
40 1. European Commission. Eurostat. Mean age of women at childbirth. Available online at: [Available
41 from:
42 [http://ec.europa.eu/eurostat/tgm/table.do?tab=table&init=1&language=en&pcode=tps00017&plugin=](http://ec.europa.eu/eurostat/tgm/table.do?tab=table&init=1&language=en&pcode=tps00017&plugin=1)
43 [1](http://ec.europa.eu/eurostat/tgm/table.do?tab=table&init=1&language=en&pcode=tps00017&plugin=1) (Accessed April 12, 2019).
44
45 2. Laughon SK, Branch DW, Beaver J, Zhang J. Changes in labor patterns over 50 years. *Am J Obstet*
46 *Gynecol.* 2012;206(5):419.e1-9.
47
48 3. Heino A GM. Nordic perinatal statistics 2016. Available online at: [updated 5th March 2018.
49 Available from: http://www.julkari.fi/bitstream/handle/10024/136095/Tr05_08.pdf?sequence=1
50 (Accessed April 11, 2019)
51
52 4. Roos N, Sahlin L, Ekman-Ordeberg G, Kieler H, Stephansson O. Maternal risk factors for postterm
53 pregnancy and cesarean delivery following labor induction. *Acta Obstet Gynecol Scand.*
54 2010;89(8):1003-10.
55
56
57
58
59
60

- 1
2
3 5. Middleton P, Shepherd E, Crowther CA. Induction of labour for improving birth outcomes for
4 women at or beyond term. *Cochrane Database Syst Rev*. 2018;5:CD004945.
- 5
6 6. Koopmans CM, Bijlenga D, Groen H, et al. Induction of labour versus expectant monitoring for
7 gestational hypertension or mild pre-eclampsia after 36 weeks' gestation (HYPITAT): a multicentre,
8 open-label randomised controlled trial. *Lancet* . 2009;374(9694):979-88.
- 9
10
11 7. Boulvain M, Senat MV, Perrotin F, et al. Induction of labour versus expectant management for
12 large-for-date fetuses: a randomised controlled trial. *Lancet*. 2015;385(9987):2600-5.
- 13
14 8. WHO Statement on Caesarean Section Rates. Available online at: [Available from:
15 http://www.who.int/reproductivehealth/publications/maternal_perinatal_health/cs-statement/en/
16 (Accessed April 12, 2019).
- 17
18
19 9. Farchi S, Polo A, Franco F, Di Lallo D, Guasticchi G. Severe postpartum morbidity and mode of
20 delivery: a retrospective cohort study. *Acta Obstet Gynecol Scand*. 2010;89(12):1600-3.
- 21
22 10. Silver RM, Landon MB, Rouse DJ, et al. Maternal morbidity associated with multiple repeat
23 cesarean deliveries. *Obstet Gynecol*. 2006;107(6):1226-32.
- 24
25 11. Betran AP, Torloni MR, Zhang JJ, Gulmezoglu AM. WHO Statement on Caesarean Section Rates.
26 *BJOG* . 2016;123(5):667-70.
- 27
28 12. Herstad L, Klungsoyr K, Skjaerven R, et al. Maternal age and emergency operative deliveries at
29 term: a population-based registry study among low-risk primiparous women. *BJOG*. 2015;122(12):1642-
30 51.
- 31
32 13. Bergholt T, Lim LK, Jørgensen JS, Robson MS. Maternal body mass index in the first trimester and
33 risk of cesarean delivery in nulliparous women in spontaneous labor. *Am J Obstet Gynecol*. 2007;196:
34 163.e1-5.
- 35
36 14. Fyfe EM, Anderson NH, North RA, et al. Risk of first-stage and second-stage cesarean delivery by
37 maternal body mass index among nulliparous women in labor at term. *Obstet Gynecol*.
38 2011;117(6):1315-22.
- 39
40 15. Pyykonen A, Gissler M, Lokkegaard E, et al. Cesarean section trends in the Nordic Countries - a
41 comparative analysis with the Robson classification. *Acta Obstet Gynecol Scand*. 2017;96(5):607-16.
- 42
43 16. Ananth CV, Chauhan SP. Epidemiology of twinning in developed countries. *Sem Perinatol*.
44 2012;36(3):156-61.
- 45
46 17. Betran AP, Merialdi M, Lauer JA, et al. Rates of caesarean section: analysis of global, regional and
47 national estimates. *Paediatr Perinat Epidemiol*. 2007;21(2):98-113.
- 48
49 18. Martin JA HB, Osterman JK, Driscoll AK, Mathews MS. Births: Final Data for 2015. *Natl Vital Stat*
50 *Rep*. 2017;66:1.
- 51
52
53
54
55
56
57
58
59
60

19. Plevani C, Incerti M, Del Sorbo D, et al. Cesarean delivery rates and obstetric culture - an Italian register-based study. *Acta Obstet Gynecol Scand.* 2017;96(3):359-65.
20. Rossen J, Klungsoyr K, Albrechtsen S, et al. Can oxytocin augmentation modify the risk of epidural analgesia by maternal age in cesarean sections? *Acta Obstet Gynecol Scand.* 2018;97(7):872-9.
21. Torloni MR, Betran AP, Souza JP, et al. Classifications for cesarean section: a systematic review. *PloS One.* 2011;6(1):e14566.
22. Grobman WA, Rice MM, Reddy UM, et al. Labor Induction versus Expectant Management in Low-Risk Nulliparous Women. *N Engl J Med.* 2018;379(6):513-23.
23. Kolkman DG, Verhoeven CJ, Brinkhorst SJ, et al. The Bishop score as a predictor of labor induction success: a systematic review. *Am J Perinatol.* 2013;30(8):625-30.
24. Cheng YW, Kaimal AJ, Snowden JM, Nicholson JM, Caughey AB. Induction of labor compared to expectant management in low-risk women and associated perinatal outcomes. *Am J Obstet Gynecol.* 2012;207(6):502.e1-8.
25. Wang Y, Tanbo T, Abyholm T, Henriksen T. The impact of advanced maternal age and parity on obstetric and perinatal outcomes in singleton gestations. *Arch Gynecol Obstet.* 2011;284(1):31-7.
26. Ankarcrona V, Altman D, Wikstrom AK, Jacobsson B, Brismar Wendel S. Delivery outcome after trial of labor in nulliparous women 40 years or older-A nationwide population-based study. *Acta Obstet Gynecol Scand.* 2019;98(9):1195-203.
27. Walker KF, Bugg GJ, Macpherson M, et al. Randomized Trial of Labor Induction in Women 35 Years of Age or Older. *New Engl J Med.* 2016;374(9):813-22.
28. Saccone G, Berghella V. Induction of labor at full term in uncomplicated singleton gestations: a systematic review and metaanalysis of randomized controlled trials. *Am J Obstet Gynecol.* 2015;213(5):629-36.
29. Nippita TA, Porter M, Seeho SK, Morris JM, Roberts CL. Variation in clinical decision-making for induction of labour: a qualitative study. *BMC Pregnancy Childbirth.* 2017;17(1):317.
30. Vellekoop J, Vrouwenraets FP, van der Steeg JW, Mol BW, Roumen FJ. Indications and results of labour induction in nulliparous women: an interview among obstetricians, residents and clinical midwives. *Eur J Obstet Gynecol Reprod Biol.* 2009;146(2):156-9.

Figure and table legends

Figure 1. The proportion of women having cesarean section (CS) before onset of labor $R2b/(R1+R2)$ and $R4b/(R3+R4)$, having labor induced $R2a/(R1+R2)$ and $R4a/(R3+R4)$ and CS rates in women with spontaneous and induced labor $CS/(R1+R2a)$ and $CS/(R3+R4a)$ by parity. Nordic countries, 2000-2011.

Table 1. Total number of nulliparous women with a singleton infant in cephalic presentation at term and with induced labor (Robson group 2a), displayed with the cesarean section (CS) rate (%), by maternal age, country, time period, and gestational age. Nordic countries, 2000-2011.

Table 2. Total number of multiparous women without previous CS, with a singleton infant in cephalic presentation at term and with induced labor (Robson group 4a), displayed with the cesarean section (CS) rate (%), by maternal age, country, time period, and gestational age. Nordic countries, 2000-2011.

Table 3. Total number of women with a singleton infant in cephalic presentation at term, no previous cesarean section (CS) and with induced or spontaneous onset of labor (Robson group 1+2a and Robson group 3+4a), displayed with number (N), CS rate (%) and relative risk (RR) with 95% confidence intervals (CI), by maternal age and parity. In addition, RR with 95% CI of CS in women having induced labor compared to women with spontaneous onset of labor is presented. Nordic countries, 2000-2011.

Figure 1:

The proportion of women having CS before onset of labor $R2b/(R1+R2)$ and $R4b/(R3+R4)$, having labor induced $R2a/(R1+R2)$ and $R4a/(R3+R4)$ and CS rates in women with spontaneous and induced labor $CS/(R1+R2a)$ and $CS/(R3+R4a)$ by parity. Nordic countries, 2000-2011




Table 1.

Total number of nulliparous women with a singleton infant in cephalic presentation at term and with induced labor (Robson group 2a), displayed along with the cesarean section rate (%), by maternal age, country, time period, and gestational age. Shown for the Nordic countries, 2000-2011.

Maternal age		<20		20-24		25-29		30-34		35-39		>40		Total	
		n	CS %	n	CS %	n	CS %	n	CS %	n	CS %	n	CS %	n	CS %
	Total	7 285	(14.0)	38 932	(18.7)	68 987	(23.2.)	54 752	(28.2)	21 484	(34.3)	4 780	(39.9)	196 220	(25.0)
Country	Denmark	1 222	(13.5)	7 478	(17.9)	16 530	(23.3)	12 023	(27.4)	4 298	(32.9)	873	(38.3)	42 424	(24.5)
	Finland	2 418	(12.4)	10 593	(17.0)	15 460	(20.4)	11 097	(25.5)	4 446	(31.4)	1 034	(38.8)	45 048	(21.9)
	Iceland	280	(14.6)	1 092	(23.9)	1 277	(25.7)	576	(32.3)	248	(35.1)	70	(35.7)	3 543	(26.2)
	Norway	1 668	(14.6)	8 683	(18.6)	14 532	(22.4)	10 707	(25.7)	3 935	(32.2)	809	(37.2)	40 334	(23.4)
	Sweden	1 697	(15.9)	11 086	(20.4)	21 188	(25.5)	20 349	(31.3)	8 557	(37.5)	1 994	(42.4)	64 871	(28.3)
Time period	2000-2002	1 654	(12.9)	8 052	(16.8)	13 986	(22.6)	9 577	(26.7)	3 317	(33.3)	574	(36.6)	37 160	(23.2)
	2003-2005	1 669	(12.4)	8 753	(18.8)	16 310	(23.0)	12 889	(28.3)	4 768	(33.7)	883	(40.7)	45 272	(24.8)
	2006-2008	1 863	(15.2)	9 834	(19.6)	17 603	(24.3)	15 019	(29.8)	6 003	(35.5)	1 399	(40.2)	51 721	(26.4)
	2009-2011	2 099	(14.9)	12 293	(19.2)	21 088	(22.7)	17 267	(27.4)	7 396	(34.3)	1 924	(40.3)	62 067	(25.0)
Gestational age	37+0 to 38+6	1 477	(13.4)	7 575	(15.6)	12 492	(19.4)	8 987	(22.6)	3 312	(27.2)	665	(37.0)	34 508	(20.2)
	39+0 to 41+6	4 016	(12.8)	21 300	(17.8)	36 604	(21.8)	28 584	(26.5)	11 756	(33.2)	3 132	(38.5)	105 392	(23.7)
	> 41+6	1 792	(17.0)	10 057	(22.9)	19 891	(28.2)	17 181	(33.9)	6 416	(40.2)	983	(46.4)	56 320	(30.3)

Table 2.

Total number of multiparous women without previous CS, with a singleton infant in cephalic presentation at term and with induced labor (Robson group 4a), along-displayed with the CS rate (%), by maternal age, country, time period, and gestational age. ~~Shown for the~~ Nordic countries, 2000-2011.

Maternal age		<20		20-24		25-29		30-34		35-39		>40		Total	
		n	CS %	n	CS %	n	CS %	n	CS %	n	CS %	n	CS %	n	CS %
		432	(3.9)	13 329	(3.2)	47 629	(4.6)	71 500	(5.2)	43 931	(6.9)	11 337	(9.1)	188 158	(5.5)
Country	Denmark	78	(5.1)	2 326	(3.9)	10 267	(5.5)	16 772	(5.6)	9 495	(7.3)	2 116	(10.1)	41 054	(6.1)
	Finland	165	(1.2)	4 928	(2.2)	13 802	(2.9)	17 773	(3.3)	10 783	(4.6)	3 084	(5.2)	50 535	(3.4)
	Iceland	9	(11.1)	256	(4.7)	1 049	(3.2)	1 392	(5.6)	827	(7.0)	236	(8.5)	3 769	(5.4)
	Norway	83	(2.4)	2 555	(4.3)	9 915	(5.4)	15 179	(5.8)	9 004	(7.9)	2 080	(10.6)	38 816	(6.3)
	Sweden	97	(8.2)	3 264	(3.5)	12 596	(5.1)	20 384	(6.1)	13 822	(7.8)	3 821	(10.9)	53 984	(6.5)
Time-period	2000-2002	75	(4.0)	2 856	(3.2)	10 389	(5.1)	14 562	(5.2)	8 214	(6.5)	1 901	(9.0)	37 997	(5.5)
	2003-2005	112	(4.5)	3 089	(3.0)	11 399	(4.7)	16 753	(5.5)	10 098	(7.5)	2 454	(8.9)	43 905	(5.8)
	2006-2008	119	(4.2)	3 356	(3.6)	11 605	(4.6)	18 591	(5.4)	11 522	(7.1)	3 151	(10.0)	48 344	(5.8)
	2009-2011	126	(3.2)	4 028	(3.2)	14 236	(4.1)	21 594	(4.8)	14 097	(6.6)	3 831	(8.6)	57 912	(5.2)
Gestational age	37+0 to 38+6	117	(4.3)	3 630	(3.0)	11 779	(4.5)	16 798	(5.2)	10 278	(7.1)	2 654	(9.0)	45 256	(5.5)
	39+0 to 41+6	252	(4.4)	7 436	(3.0)	26 706	(4.2)	39 959	(4.9)	25 010	(6.3)	6 899	(8.5)	106 262	(5.2)
	> 41+6	63	(1.6)	2 263	(4.6)	9 144	(5.7)	14 743	(6.0)	8 643	(8.4)	1 784	(11.7)	36 640	(6.7)

Table 3. Total number of women with a singleton infant in cephalic presentation at term, no previous cesarean section (CS) and with induced or spontaneous onset of labor (Robson group 1+2a and Robson group 3+4a), displayed with number (N), CS rate (%) and relative risk (RR) with 95% confidence intervals (CI), by maternal age and parity. In addition, RR with 95% CI of CS in women having induced labor compared to women with spontaneous onset of labor is presented. Nordic countries, 2000-2011.

	Maternal age Years	Induced labor (R2a)					Spontaneous labor (R1)					Induced/Spontaneous	
		Total	N	CS rate %	RR	95% CI	Total	N	CS rate %	RR	95% CI	RR	95% CI
Nulliparous women	< 20	7 285	1 020	14.0	0.60	(0.57-0.64)	48 463	2 278	4.7	0.57	(0.55-0.60)	2.98	(2.78-3.19)
	20-24	38 932	7 280	18.7	0.81	(0.79-0.83)	239 336	15 317	6.4	0.78	(0.77-0.80)	2.92	(2.85-3.00)
	25-29	68 987	16 005	23.2	1.00		409 618	33 591	8.2	1.00		2.83	(2.78-2.88)
	30-34	54 752	15 440	28.2	1.22	(1.19-1.24)	271 596	29 877	11.0	1.34	(1.32-1.36)	2.56	(2.52-2.61)
	35-39	21 484	7 369	34.3	1.48	(1.44-1.51)	77 338	11 678	15.1	1.84	(1.81-1.88)	2.27	(2.21-2.32)
	> 40	4 780	1 907	39.9	1.72	(1.66-1.79)	11 309	2 375	21.0	2.50	(2.37-2.64)	1.90	(1.81-2.00)
Multiparous women	Maternal age	Induced labor (R4a)					Spontaneous labor (R3)					Induced/Spontaneous	
	Years	Total	N	CS rate %	RR	95% CI	Total	N	CS rate %	RR	95% CI	RR	95% CI
	< 20	432	17	3.9	0.86	(0.54-1.37)	3 701	52	1.4	0.88	(0.67-1.15)	2.80	(1.63-4.80)
	20-24	13 329	432	3.2	0.70	(0.63-0.77)	104 917	1 364	1.3	0.81	(0.77-0.86)	2.49	(2.24-2.77)
	25-29	47 569	2 174	4.6	1.00		357 208	5 716	1.6	1.00		2.86	(2.72-3.00)
	30-34	71 500	3 721	5.2	1.13	(1.07-1.19)	494 883	9 403	1.9	1.19	(1.15-1.23)	2.74	(2.64-2.84)
35-39	43 931	3 040	6.9	1.50	(1.42-1.58)	245 197	6 375	2.6	1.63	(1.57-1.68)	2.66	(2.55-2.78)	
> 40	11 337	1 032	9.1	1.98	(1.84-2.12)	43 575	1 743	4.0	2.50	(2.37-2.64)	2.28	(2.11-2.45)	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

For Peer Review