



Integrability via Geometry: Dispersionless Differential Equations in Three and Four Dimensions

David M. J. Calderbank¹ , Boris Kruglikov²

¹ Department of Mathematical Sciences, University of Bath, Claverton Down, Bath BA2 7AY, UK.
E-mail: D.M.J.Calderbank@bath.ac.uk

² Institute of Mathematics and Statistics, UiT the Arctic University of Norway, 90-37 Tromsø, Norway.
E-mail: Boris.Kruglikov@uit.no


Received: 18 February 2020 / Accepted: 18 September 2020
© The Author(s) 2020

Abstract: We prove that the existence of a dispersionless Lax pair with spectral parameter for a nondegenerate hyperbolic second order partial differential equation (PDE) is equivalent to the canonical conformal structure defined by the symbol being Einstein–Weyl on any solution in 3D, and self-dual on any solution in 4D. The first main ingredient in the proof is a characteristic property for dispersionless Lax pairs. The second is the projective behaviour of the Lax pair with respect to the spectral parameter. Both are established for nondegenerate determined systems of PDEs of any order. Thus our main result applies more generally to any such PDE system whose characteristic variety is a quadric hypersurface.

1. Introduction and Main Results

The integrability of dispersionless partial differential equations is well known to admit a geometric interpretation. Twistor theory [26, 29] gives a framework to visualize this for several types of integrable systems, as demonstrated by many examples [2, 10, 11, 14, 19, 30, 37].

Recently, such a relation has been established for several classes of second order equations in 3D and one class in 4D [17]. Namely the following equivalences have been established:



Hydrodynamic integrability in 2D (also written “1 + 1 dimension”) was introduced in [9] and elaborated in [35]. Integrability via hydrodynamic reductions in $d \geq 3$ dimensions was developed in [16]. This method, although constructive, is not universal, as it applies only to translation invariant equations (invariantly, this requires the existence of a d -dimensional abelian contact symmetry group). Thus the upper part of the above diagram, at least at present, does not extend to the general class of second order PDEs.

On the other hand, the two other ingredients of the diagram are universal. The main aim of this paper is to prove the bottom equivalence for large class of PDE systems, including general second order PDEs, in 3D and 4D, where “integrable background geometry” means that a canonical conformal structure on solutions of the equation is Einstein–Weyl in 3D and self-dual in 4D (these geometries are “backgrounds” for integrable gauge theories [2]).

Consider a second order PDE

$$\mathcal{E} : F(x, u, \partial u, \partial^2 u) = 0 \tag{1}$$

for a scalar function u of an independent variable x on a connected manifold M with $\dim M = d$, where $\partial u = (u_i)$ and $\partial^2 u = (u_{ij})$ denote partial derivatives of u in local coordinates $x = (x^i)$. Let M_u denote the manifold M equipped with a given scalar function u ; concretely, we may view M_u as the graph of u in $M \times \mathbb{R}$. A tensor on M_u is, by definition, a tensor on M , which may also depend, at each $x \in M$, on finitely many derivatives of u at x .

Let σ_F be the linearization of F in second derivatives, i.e.,

$$\sigma_F = \sum_{i \leq j} \frac{\partial F}{\partial u_{ij}} \partial_i \partial_j = \sum_{i, j} \sigma_{ij}(u) \partial_i \otimes \partial_j, \quad \text{where } \sigma_{ij}(u) := \frac{1 + \delta_{ij}}{2} \frac{\partial F}{\partial u_{ij}}.$$

Invariantly, σ_F defines a section of $S^2 T M_u$, hence a quadratic form on $T_x^* M_u$ for each $x \in M_u$, called the *symbol* of F . If we change the defining function F of \mathcal{E} , σ_F changes by a conformal rescaling on \mathcal{E} . Hence the conformal class of σ_F along $F = 0$ is an invariant of \mathcal{E} , as is the *characteristic variety* $\chi^{\mathcal{E}} \rightarrow M_u$, the bundle whose fibre at $x \in M_u$ is the projective variety $\chi_x^{\mathcal{E}} := \text{Char}(\mathcal{E}, u)_x = \{[\theta] \in \mathbb{P}(T_x^* M_u) \mid \sigma_F(\theta) = 0\}$.

We assume henceforth that (1) is:

- *nondegenerate*, i.e., σ_F is nondegenerate at generic points of the zero-set \mathcal{E} of F . This is equivalent to $\det(\sigma_{ij}(u)) \neq 0$ for a generic solution u .
- *hyperbolic*, i.e., M is complex and F is holomorphic, or M is real, F is smooth and the variety $\{[\theta] \in \mathbb{P}(T^* M_u \otimes \mathbb{C}) \mid \sigma_F(\theta) = 0\}$ of complex characteristics is a complexification of $\chi^{\mathcal{E}}$ for a generic (real) solution u .

The nondegeneracy of σ_F implies that its inverse

$$g_F = \sum_{i, j} g_{ij}(u) dx^i dx^j, \quad \text{where } (g_{ij}(u)) = (\sigma_{ij}(u))^{-1},$$

defines a nondegenerate symmetric bilinear form on $T_x M_u$ for any (x, u) sufficiently close to a generic point of $F = 0$. As in [17], the corresponding conformal structure c_F plays a central role in this paper. Hyperbolicity implies that along $F = 0$, c_F is uniquely determined by the bundle $\chi^{\mathcal{E}}$ of nonsingular quadric hypersurfaces because the latter is dual to the projectivized null cone of c_F .

A *dispersionless Lax pair* [38] or *dLp* for (1) can be described as rank one covering system [36] of \mathcal{E} . Roughly speaking, this means that there is a fibre bundle $\hat{\pi} : \hat{M}_u \rightarrow M_u$ with connected rank one fibres, and a PDE system on \hat{M}_u with \mathcal{E} as a differential corollary. There are various ways to formulate this precisely; in this paper we adopt as a definition that there are linearly independent vector fields \hat{X} and \hat{Y} on \hat{M}_u , whose coefficients depend on finitely many derivatives of u , such that \mathcal{E} is the Frobenius integrability condition for their span $\hat{\Pi} \subseteq T\hat{M}_u$ —this is the condition that $[\hat{X}, \hat{Y}]$ is a section of $\hat{\Pi}$, so that $\hat{\Pi}$ is tangent to a foliation of \hat{M}_u by surfaces.

The leaf space of this foliation (for a solution u of \mathcal{E}) is sometimes called the *twistor space* $\mathcal{T}w$ of the dLp in 4D (or *minitwistor space* in 3D). However, a well-behaved twistor space may only exist over suitable open subsets of M_u , so its geometry is more conveniently described on the *correspondence space* \hat{M}_u . For instance, functions on $\mathcal{T}w$ correspond to solutions of a linear PDE system for functions on \hat{M}_u that are constant on the leaves of the foliation, while hypersurfaces in $\mathcal{T}w$ may be described as solutions of a quasilinear PDE system for sections of $\hat{\pi} : \hat{M}_u \rightarrow M_u$ that are unions of such leaves. Either of these PDE systems can equivalently be called a dLp: \mathcal{E} ensures their compatibility.

A fibre coordinate $\lambda : \hat{M}_u \rightarrow \mathbb{R}$ is called a *spectral parameter* and it locally identifies \hat{M}_u with $M_u \times \mathbb{R}$. We may then write $\hat{X} = X + m \partial_\lambda$, $\hat{Y} = Y + n \partial_\lambda$ where X, Y are λ -parametric vector fields on M_u , and a section of $\hat{\pi} : \hat{M}_u \rightarrow M_u$ may be written $\lambda = q(\mathbf{x})$ for a function $q : M_u \rightarrow \mathbb{R}$. The dLp $\hat{\Pi}$ then has the geometric interpretation that \mathcal{E} is the integrability condition for the existence of many foliations of M_u by surfaces which are tangent at any $\mathbf{x} \in M_u$ to the span $\Pi = \hat{\pi}_*(\hat{\Pi})$ of X and Y at \mathbf{x} , with $\lambda = q(\mathbf{x})$.

A fundamental motivation for this paper is that in all known examples of such dLps, it has been observed (see e.g. [17]) that Π is *characteristic* for \mathcal{E} in the sense that for any solution u , and any 1-form θ on M_u with $\Pi \subseteq \ker \theta$, we have $[\theta] \in \chi^\mathcal{E}$. Thus for any solution u of \mathcal{E} , M_u admits many foliations by *characteristic surfaces*, and indeed \mathcal{E} is the integrability condition for their existence. Our first result establishes this characteristic property in considerable generality.

Theorem 1. *Let $\hat{\Pi}$ be a dLp on $\hat{\pi} : \hat{M}_u \rightarrow M_u$ for a determined PDE system \mathcal{E} of order ℓ on M_u . Then $\Pi = \hat{\pi}_*(\hat{\Pi})$ is characteristic for \mathcal{E} .*

We refer to Sects. 2 and 3, or [21, 22, 36], for discussion of more general PDE systems and their characteristic varieties: in this introduction, we focus on second order scalar PDEs. For such PDEs, the characteristic condition means that for each solution u and each $\hat{\mathbf{x}} \in \hat{M}_u$, $\Pi_{\hat{\mathbf{x}}}$ is a coisotropic 2-plane for the conformal structure c_F . By nondegeneracy of c_F , such 2-planes can only exist for $2 \leq d \leq 4$: for $d = 2$, the condition is vacuous; for $d = 3$, $\Pi_{\hat{\mathbf{x}}}$ is then tangent to the null cone of c_F (i.e., degenerate); for $d = 4$, $\Pi_{\hat{\mathbf{x}}}$ is then contained in the null cone (i.e., totally isotropic). In the real case, the characteristic condition further implies that c_F has (up to sign) signature $(2, 1)$ for $d = 3$ or $(2, 2)$ for $d = 4$. We assume this henceforth.

For both $d = 3$ and $d = 4$, the coisotropic 2-planes at each point $\mathbf{x} \in M$ form a 1-dimensional submanifold of the grassmannian $\text{Gr}_2(T_{\mathbf{x}}M)$. For $d = 3$ this submanifold is a rational curve ($\cong \mathbb{P}^1$, the projective line) canonically isomorphic to the conic $\chi^\mathcal{E} \subseteq \mathbb{P}(T_{\mathbf{x}}^*M)$. For $d = 4$, it is a disjoint union of two rational curves, corresponding to the two rulings of the quadric surface $\chi^\mathcal{E}$; the points of the two components are called α -planes and β -planes depending on whether the 2-planes are self-dual or anti-self-dual.

If Π is coisotropic and is also an immersion, we may thus identify \hat{M}_u locally with the \mathbb{P}^1 -bundle whose fibre over $x \in M_u$ consists of all coisotropic 2-planes for $d = 3$ or the α -plane component for $d = 4$. Under this identification, $\Pi \rightarrow \hat{M}_u$ becomes the tautological bundle of coisotropic 2-planes. Any Weyl connection ∇ on M_u (a torsion-free conformal connection on M depending on finitely many derivatives of u) induces a connection on $\hat{M}_u \rightarrow M_u$ and hence a horizontal lift of Π to distribution $\hat{\Pi}_\nabla \subseteq T\hat{M}_u$.

If $d = 4$, it is well-known [29] that $\hat{\Pi}_\nabla$ is independent of ∇ (i.e., *conformally invariant*), and is integrable if and only if (M_u, c_F) is *self-dual* (SD), i.e., the Weyl tensor W_{c_F} satisfies $W_{c_F} = *W_{c_F}$. The integral surfaces of $\hat{\Pi}_\nabla$ then project to α -surfaces for c_F .

If $d = 3$, it is similarly well known [5, 19] that $\hat{\Pi}_\nabla$ is integrable if and only if (M_u, c_F, ∇) is *Einstein–Weyl* (EW), i.e., the symmetrized Ricci tensor of ∇ is proportional to any metric g_F in the conformal class: $\text{Sym}(\text{Ric}^\nabla) = \Lambda g_F$, $\Lambda \in C^\infty(M_u)$. The integral surfaces of $\hat{\Pi}_\nabla$ then project to totally geodesic null surfaces for (c_F, ∇) .

A dLp $\hat{\Pi}$ for \mathcal{E} arising in this way for $d = 3, 4$ will be called *standard*. Two dispersionless Lax pairs $\hat{\Pi}, \hat{\Pi}'$ will be called \mathcal{E} -*equivalent*, if $\hat{\Pi} = \hat{\Pi}'$ on \hat{M}_u for any solution u of \mathcal{E} .

It is an open question in the theory of integrable systems how many non-equivalent coverings a given \mathcal{E} can possess. Our second result claims that coverings of dLp type are essentially unique under a certain nondegeneracy condition on $\hat{\Pi}$. This condition, given in Definition 7 of Sect. 4.5, depends only on $\Pi = \hat{\pi}_*(\hat{\Pi})$, implies that Π immerses, and holds in all examples we know of.

The result is straightforward when $d = 4$, but when $d = 3$, it shows that $\hat{\Pi}$ can be assumed *projective*: for some choice of spectral parameter λ and vector fields \hat{X}, \hat{Y} generating $\hat{\Pi}$, the coefficients of these vector fields are cubic polynomials in λ . The result is again not restricted to second order scalar PDEs: we require only that $\chi_x^\mathcal{E}$ is a nonsingular quadric hypersurface for each $x \in M_u$.

Theorem 2. *Let $\mathcal{E} : F = 0$ be a determined PDE system of order ℓ whose characteristic variety $\chi^\mathcal{E}$ is a bundle of nonsingular quadric hypersurfaces in $\mathbb{P}(T^*M_u)$. Then any nondegenerate dLp $\hat{\Pi}$ is \mathcal{E} -equivalent to a standard dLp $\hat{\Pi}_\nabla$ for some Weyl connection ∇ .*

Our third (and main) result establishes an equivalence between the dispersionless integrability of \mathcal{E} and the EW/SD property of c_F . However, to achieve this, some care is needed in the formulation of both properties. First, in the integrability of the dLp $\hat{\Pi}$, we must account for \mathcal{E} -equivalence. Thus we say that \mathcal{E} is *integrable by a dLp $\hat{\Pi}$* if for any $\hat{\Pi}'$, which is \mathcal{E} -equivalent to $\hat{\Pi}$, the Frobenius integrability condition for $\hat{\Pi}'$ is a nontrivial differential corollary of \mathcal{E} . Secondly, the EW/SD property should be a nontrivial differential corollary of \mathcal{E} . The need for nontriviality here is illustrated by PDEs of the form $\Delta u = f(x, u, \partial u)$: this is non-integrable for generic f , but its conformal structure is independent of u and is flat, so the EW/SD property holds automatically. For more general PDE systems \mathcal{E} , a differential corollary of \mathcal{E} *holds nontrivially* if it is not a consequence of a proper subsystem \mathcal{E}' of \mathcal{E} . We can now obtain the main result as follows.

Theorem 3. *Let $\mathcal{E} : F = 0$ be a determined PDE system in 3D or 4D whose characteristic variety $\chi^\mathcal{E}$ is a bundle of nonsingular quadric hypersurfaces, for instance a nondegenerate hyperbolic second order scalar PDE (1). Let c_F be the corresponding conformal structure. Then \mathcal{E} is integrable by a nondegenerate dLp if and only if*

- 3D: the Einstein–Weyl property for c_F holds nontrivially on solutions of \mathcal{E} ;
 4D: the self-duality property for c_F holds nontrivially on solutions of \mathcal{E} .

Proof. As a preliminary, note that if F has order ℓ , then c_F depends pointwise only on derivatives of u of order $\leq \ell$ (or $\leq (\ell - 1)$ if F is quasilinear) and so is defined and is nondegenerate for almost any u (not necessarily a solution). Thus $\hat{\Pi}_\nabla$ is defined for any Weyl connection ∇ over an open subset of M_u , and its integrability there is equivalent to the EW condition for (c_F, ∇) when $d = 3$ and the SD condition for c_F when $d = 4$.

Suppose first that $\hat{\Pi} \subseteq T\hat{M}_u$ is a dLp for \mathcal{E} . By Theorem 1, $\Pi = \hat{\pi}_*(\hat{\Pi})$ is characteristic, i.e., when $F = 0$, Π is coisotropic for the conformal structure c_F (and for $d = 4$ we orient M_u so that Π is a congruence of α -planes). Nondegeneracy of $\hat{\Pi}$ implies that Π immerses into $\text{Gr}_2(TM_u)$ and so we may assume that \hat{M}_u is an open subset of the bundle of coisotropic 2-planes for all solutions u , and hence also on an open neighbourhood of (x, u) where c_F is nondegenerate. Then by Theorem 2, $\hat{\Pi}$ is \mathcal{E} -equivalent to a standard dLp $\hat{\Pi}_\nabla$ over any open subset of M_u . Hence the EW/SD condition is a nontrivial differential corollary of \mathcal{E} , as required.

Conversely, suppose that the EW/SD condition is a nontrivial differential corollary of \mathcal{E} (for some Weyl connection ∇ when $d = 3$), and let $\hat{\pi}: \hat{M}_u \rightarrow M_u$ be the bundle of null 2-planes for $d = 3$, or the bundle of α -planes for $d = 4$. Then if $\hat{\Pi}$ is \mathcal{E} -equivalent to $\hat{\Pi}_\nabla$ (for any Weyl connection ∇ when $d = 4$) on an open subset of M_u , the integrability of $\hat{\Pi}$ is a differential corollary of \mathcal{E} on that open subset (since this is true for $\hat{\Pi}_\nabla$).

Finally if any such $\hat{\Pi}$ is a differential corollary of a proper subsystem \mathcal{E}' of \mathcal{E} , then the first part of the argument implies that the EW/SD property is also a consequence of \mathcal{E}' , contradicting nontriviality. \square

Remark 1. Often, in the physics literature, little distinction is made between a system \mathcal{E} and a system \mathcal{E}' obtained by differentiation or potentiation of \mathcal{E} . While some properties of the equation can change, for instance the symmetry algebra and dimension of the solution space, the characteristic variety and integrability of \mathcal{E} are unaltered. It is easy to adjust the formulation of the theorems to such variations between \mathcal{E} and \mathcal{E}' .

This theorem shows that the EW and SD equations are master equations, in 3D and 4D respectively, for determined integrable PDE systems whose characteristic variety is a bundle of nonsingular quadric hypersurfaces. It applies in particular to first order systems and higher order scalar equations whose (principal) symbol is a power of a nondegenerate quadratic form. However, the EW and SD equations are not themselves determined systems because of the gauge freedom coming from diffeomorphism invariance. Determined forms of the EW and SD equations were derived in [12], where it was shown in particular that the Manakov–Santini system [24] is equivalent to a determined form of the EW equation. Because of their importance, we will present novel derivations of these determined master equations using the methods of this paper.

Theorem 3 is useful for at least two reasons. First, the geometric characterizations of integrability are algorithmic. In 4D, the anti-self-dual part of the Weyl tensor of c_F on M_u can be computed explicitly from finitely many derivatives of u , and so we can check whether it vanishes on solutions by imposing the equation and its prolongations formally—we do not have to be able to resolve the PDE or even to prove its solvability. In 3D, the situation is complicated slightly by the choice of Weyl connection. For the classes of translation-invariant equations considered in [17], there is a universal formula for the Weyl connection, but this formula is not generally applicable (it is not contact-invariant). Nevertheless, except in degenerate situations, the choice is uniquely determined by

finitely many derivatives of c_F , and so the EW condition may again be verified by formally imposing the PDE on a tensor depending on finitely many derivatives of u . This effective integrability criterion has many applications: for instance, it was applied in [23] to obtain infinitely many new integrable equations in 4D as deformations of integrable Monge–Ampère equations of Hirota type.

Secondly, the EW/SD property provides a canonical characteristic Lax pair, which, if the PDE on u has order ℓ , depends on at most $\ell + 1$ derivatives of u (ℓ if the PDE is quasilinear), and satisfies a ‘normality’ condition off shell which is useful in computations. None of these properties were assumed a priori. For example, the standard Lax pair [24] for the Manakov–Santini system is not normal, and the normal Lax pair may be understood as a Lax pair for an equivalent PDE system presented in [12], which we also discuss.

Apart from the Manakov–Santini system (and variants), Theorem 3 encompasses many examples in 3D, such as the Lax pairs arising in the central quadric ansatz [15], for EW manifolds in diagonal coordinates [12], and for the systems of two first order PDE on two unknown functions studied in [7]. In 4D, there are Lax pairs having no derivatives with respect to the spectral parameter λ , which cannot be normal, such as the hypercomplex Lax pair of Dunajski and Joyce (see [2, 12]) and Lax pairs for Monge–Ampère equations of Hirota type [6] as well as systems of Chasles type [8]. However, normal Lax pairs are always available, and provide a canonical choice in 4D, while in 3D they are given by a choice of Weyl connection.

We begin the body of the paper in Sect. 2 by presenting a rigorous definition of what should be called a (nondegenerate) dispersionless Lax pair, motivated by examples. The search for such formalism in general has a long history: see [3, 25] for discussion in the dispersive context. A fundamental role is played by the λ -dependent family $\Pi = \hat{\pi}_*(\hat{\Pi})$ of rank 2 subbundles of TM_u , which we call a *2-plane congruence*. We also explain the normality condition mentioned above, observing that in 4D it determines $\hat{\Pi}$ from Π .

In Sect. 3, we prove Theorem 1. Here we treat the symbol and characteristic variety of general PDE systems. For both this, and the proof of Theorem 2, we require some jet theory, which we have generally suppressed in the rest of the paper, cf. Remark 2. Having proven Theorem 1, as an addendum, we show in Sect. 3.3 that a Lax pair which is characteristic for a quadric is nondegenerate, and give a computational criterion for the existence of such a quadric for nondegenerate Lax pairs.

For PDE systems whose characteristic variety is a quadric, Theorem 1 shows that Π is essentially unique, which considerably constrains the choice of $\hat{\Pi}$, especially in 4D. In 3D, however, more work is needed to prove Theorem 2, which we develop in Sect. 4. We first discuss the standard EW/SD Lax pairs, which are not only normal, but projective. We also introduce and motivate a stronger nondegeneracy condition on the Lax pair $\hat{\Pi}$. Roughly speaking, this condition means that the equation \mathcal{E} appears nontrivially in the symbol of the integrability condition for $\hat{\Pi}$ (i.e., at highest order). From this we deduce the projective property of the Lax pair, and hence prove Theorem 2.

In Sect. 5 we discuss applications and extensions of the viewpoint we have developed. In particular, we discuss pseudopotentials and their relation to contact coverings, the twistor interpretation of this relationship, and potential generalizations of the theory.

2. Lax Pairs: Nondegeneracy and Normalization

2.1. Dispersionless pairs and 2-plane congruences. We begin with a well-known prototypical example.

Example 1. (dKP) The dispersionless Kadomtsev–Petviashvili (dKP) equation (see for example [14]) is the second order scalar PDE

$$F(x, u, \partial u, \partial^2 u) := u_{xt} + (uu_t)_t - u_{yy} = 0 \quad (2)$$

for a scalar function u on a 3-manifold $M_u \simeq M$ with coordinates (x, y, t) . (This differs from some standard conventions by the interchange $t \leftrightarrow x$ and/or $u \mapsto -u$.) The dKP equation is the compatibility condition $\psi_{xy} = \psi_{yx}$ of the first order linear system

$$\psi_x - (\lambda^2 - u) \psi_t - (u_y + \lambda u_t) \psi_\lambda = 0, \quad \psi_y - \lambda \psi_t - u_t \psi_\lambda = 0,$$

for a scalar function ψ on $\hat{M}_u = M_u \times \mathbb{R}$ with coordinates (x, y, t, λ) . It may also be described as the compatibility condition $q_{xy} = q_{yx}$ for the quasilinear system

$$q_x = (q^2 - u) q_t - q u_t - u_y, \quad q_y = q q_t - u_t$$

for a scalar function $q = q(x, y, t)$ on M_u . In more geometric terms, ψ is a function on \hat{M}_u which is invariant under the vector fields

$$\hat{X} = \partial_x - (\lambda^2 - u) \partial_t - (u_y + \lambda u_t) \partial_\lambda, \quad \hat{Y} = \partial_y - \lambda \partial_t - u_t \partial_\lambda, \quad (3)$$

while q defines a section of $\hat{\pi}: \hat{M}_u \rightarrow M_u$ such that \hat{X} and \hat{Y} are tangent to its image. The compatibility condition in either case is that \hat{X} and \hat{Y} span a distribution $\hat{\Pi} \subseteq T\hat{M}_u$ which is (Frobenius) integrable, i.e., $[\hat{X}, \hat{Y}]$ is also section of $\hat{\Pi}$. In this example, the Frobenius integrability condition holds if and only if $[\hat{X}, \hat{Y}] = 0$ if and only if (2) is satisfied.

In this paper, we take the distribution $\hat{\Pi}$ on \hat{M}_u to be the fundamental object.

Definition 1. A *dispersionless pair* of order $\leq N$ is a bundle $\hat{\pi}: \hat{M}_u \rightarrow M_u$ called the *correspondence space*, whose fibres are connected curves, together with a rank two distribution $\hat{\Pi} \subseteq T\hat{M}_u$ such that:

- for all $\hat{x} \in \hat{M}_u$, $\hat{\Pi}_{\hat{x}} \subseteq T_{\hat{x}}\hat{M}_u$ depends on u only through its partial derivatives at $\mathbf{x} = \hat{\pi}(\hat{x}) \in M_u$ of order $\leq N$;
- $\hat{\Pi}$ is transverse to the fibres of $\hat{\pi}$, i.e., $\hat{\Pi} \cap \ker \hat{\pi}_* = 0$.

A *spectral parameter* is a local fibre coordinate $\lambda = \lambda(\hat{x}): \hat{M}_u \rightarrow \mathbb{R}$ on \hat{M}_u .

If $\hat{\Pi} = \langle \hat{X}, \hat{Y} \rangle$, we thus obtain a first order linear system

$$\hat{X}(\psi) = 0, \quad \hat{Y}(\psi) = 0 \quad (4)$$

for functions ψ on \hat{M}_u . In terms of a spectral parameter λ , a section of $\hat{\pi}$ has image $\lambda = q(\mathbf{x})$ for a function $q: M_u \rightarrow \mathbb{R}$, and the corresponding first order quasilinear system is given by

$$\hat{X}(\lambda - q(\mathbf{x}))|_{\lambda=q(\mathbf{x})} = 0, \quad \hat{Y}(\lambda - q(\mathbf{x}))|_{\lambda=q(\mathbf{x})} = 0. \quad (5)$$

The system (4) is compatible if and only if (5) is compatible if and only if the distribution $\hat{\Pi}$ is integrable. Then solutions of (4) and (5) describe respectively functions and hypersurfaces in the (local) leaf space of the foliation tangent to $\hat{\Pi}$ (the twistor or minitwistor space). The integrability condition of $\hat{\Pi}$ is a PDE on u of order $\leq N + 1$. Roughly speaking—see Definition 5—dispersionless integrable systems are PDEs arising as such integrability conditions.

We need not restrict attention to scalar PDEs. Indeed we wish to encompass the following important system due to Manakov and Santini [24].

Example 2. (MS) The Manakov–Santini (MS) system is the second order coupled system of PDEs

$$S(u) + u_t^2 = 0, \quad S(v) = 0 \quad (6)$$

for functions (u, v) of (x, y, t) , where

$$S = \partial_t \partial_x + v_t \partial_t \partial_y + (u - v_y) \partial_t^2 - \partial_y^2. \quad (7)$$

(As with the dKP equation, we have aligned our coordinate conventions for consistency within this paper. Conventions in the literature [12, 24, 28] vary, but are all equivalent to the one here by point transformations.)

As noted in [24], system (6) is the Frobenius integrability condition for the dispersionless pair $\hat{\Pi} = \langle \hat{X}, \hat{Y} \rangle$ spanned by

$$\hat{X} = \partial_x - (\lambda^2 + v_t \lambda - u + v_y) \partial_t - (u_t \lambda + u_y) \partial_\lambda, \quad \hat{Y} = \partial_y - (\lambda + v_t) \partial_t - u_t \partial_\lambda. \quad (8)$$

The corresponding quasilinear covering system, which was studied in [28] and more recently in [32], is

$$q_x = (q^2 + q v_t - u + v_y) q_t - q u_t - u_y, \quad q_y = (v_t + q) q_t - u_t.$$

When $v = 0$, the MS system reduces to the dKP equation, and (8) to (3). When $u = 0$, the dLp (8) has no derivatives with respect to the spectral parameter.

If $\hat{\Pi}$ is a dispersionless pair, then $\Pi := \hat{\pi}_*(\hat{\Pi})$ is a rank 2 subbundle of $\hat{\pi}^* T M_u$, so at each $\mathbf{x} \in M_u$, we have a 1-parameter family of 2-dimensional subspaces of $T_{\mathbf{x}} M_u$.

Definition 2. A 2-plane congruence Π over M_u is a section $\Pi: \hat{M}_u \rightarrow \hat{\pi}^* \text{Gr}_2(T M_u)$, where $\text{Gr}_2(T M_u) \rightarrow M_u$ is the bundle whose fibre over $\mathbf{x} \in M_u$ is the grassmannian of 2-dimensional vector subspaces of $T_{\mathbf{x}} M_u$.

Conversely, the passage from a 2-plane congruence Π to a dispersionless pair $\hat{\Pi}$ can be understood as a *lift* with respect to the projection $\hat{\pi}: \hat{M}_u \rightarrow M_u$. It is convenient to describe the lift condition in terms of the rank 3 distribution $\Delta = \hat{\pi}_*^{-1}(\Pi) \subseteq T \hat{M}_u$: $\hat{\Pi}$ is a lift of Π if and only if it is a rank 2 subbundle of Δ transverse to the fibres of $\hat{\pi}$. For any distributions $D_1, D_2 \subseteq T \hat{M}_u$ we denote by $[D_1, D_2]$ the distribution generated by Lie brackets of sections of D_1 and D_2 . Thus the integrability condition for $\hat{\Pi}$ is that its derived distribution $[\hat{\Pi}, \hat{\Pi}]$ is equal to $\hat{\Pi}$.

More explicitly, we choose a spectral parameter λ and let X, Y be linearly independent λ -parametric vector fields on M_u depending at each \mathbf{x} only on the partial derivatives of u at \mathbf{x} of order $\leq N$. Then $\Pi = \langle X, Y \rangle$ is a 2-plane congruence, and Δ is the span of the coordinate lifts of X, Y (still denoted X, Y , with $X(\lambda) = 0 = Y(\lambda)$) and ∂_λ . Then we write a dispersionless pair $\hat{\Pi}$ on \hat{M}_u , with $\hat{\pi}_*(\hat{\Pi}) = \Pi$ as the span $\hat{\Pi} = \langle \hat{X}, \hat{Y} \rangle$ of vector fields

$$\hat{X} = X + m \partial_\lambda, \quad \hat{Y} = Y + n \partial_\lambda \quad (9)$$

with $\hat{\pi}_*(\hat{X}) = X$ and $\hat{\pi}_*(\hat{Y}) = Y$, where m, n are functions of \mathbf{x}, u , and the spectral parameter λ . The derived distribution of $\hat{\Pi}$ is now $[\hat{\Pi}, \hat{\Pi}] = \langle \hat{X}, \hat{Y}, [\hat{X}, \hat{Y}] \rangle \subseteq T \hat{M}_u$, which generically has rank 3, and the integrability condition is that it has rank 2.

In 3D, we may introduce coordinates (x, y, t) and choose generators of Π of the form

$$X = \partial_x - \alpha \partial_t, \quad Y = \partial_y - \beta \partial_t, \quad (10)$$

where the functions α, β depend on $(x, y, t), u$ and λ . Dually, the annihilator $\text{Ann}(\Pi)$ of Π in $\hat{\pi}^*T^*M_u$ is spanned by the λ -dependent 1-form

$$\theta = dt + \alpha dx + \beta dy, \quad (11)$$

$\text{Ann}(\Delta)$ is spanned by the pullback of θ to \hat{M}_u (which we still denote by θ), while $\text{Ann}(\hat{\Pi})$ is spanned by θ and the 1-form

$$\eta = d\lambda - m dx - n dy \quad (12)$$

on \hat{M}_u . Hence $\hat{\Pi}$ is the radical of the 2-form $\theta \wedge \eta$.

In 4D, we similarly may assume that we have coordinates (x, y, z, t) and generators

$$X = \partial_x - \alpha \partial_z - \beta \partial_t, \quad Y = \partial_y - \gamma \partial_z - \delta \partial_t, \quad (13)$$

where $\alpha, \beta, \gamma, \delta$ depend on $(x, y, z, t), u$ and λ . Thus $\text{Ann}(\Pi)$ is spanned by

$$\zeta = dz + \alpha dx + \gamma dy, \quad \theta = dt + \beta dx + \delta dy, \quad (14)$$

$\text{Ann}(\Delta)$ by their pullbacks, and $\text{Ann}(\hat{\Pi}) = \langle \zeta, \theta, \eta \rangle$ with η given by (12). In both 3D and 4D, with \hat{X} and \hat{Y} given by (9), $\hat{\Pi}$ is integrable if and only if $[\hat{X}, \hat{Y}] = 0$.

2.2. Normality and nondegeneracy. In order for $\hat{\Pi}$ to be a dispersionless Lax pair for an equation $\mathcal{E} : F = 0$, we require that the integrability condition $[\hat{\Pi}, \hat{\Pi}] = \hat{\Pi}$ holds modulo \mathcal{E} , i.e., when $F = 0$ or, to use physics terminology, *on shell*.

Definition 3. We say that the dispersionless pair $\hat{\Pi} \subseteq T\hat{M}_u$ is *normal* if $[\hat{\Pi}, \hat{\Pi}] \subseteq \Delta$ off shell, i.e., without assuming $F = 0$. In other words, $\hat{\pi}_*([\hat{\Pi}, \hat{\Pi}]) = \Pi$.

If $\hat{\Pi} = \langle \hat{X}, \hat{Y} \rangle$ with \hat{X} and \hat{Y} defined by (9), (10) and (13), then $\hat{\Pi}$ is normal if and only if $[\hat{X}, \hat{Y}]$ is a multiple of ∂_λ . In this case the integrability condition reduces to the vanishing of the ∂_λ -component $\hat{X}(n) - \hat{Y}(m)$ of the vector field $[\hat{X}, \hat{Y}]$ (identically in λ).

When $d = 4$, a generic 2-plane congruence Π has a unique normal lift. Indeed, generically, Δ is nonholonomic with $[\Delta, \Delta] = T\hat{M}_u$, i.e., it has the growth vector $(3, 5)$, and following Cartan [4, §11], there is a unique rank 2 subbundle $\hat{\Pi} \subseteq \Delta$ with $[\hat{\Pi}, \hat{\Pi}] = \Delta$. Such rank 2 distribution $\hat{\Pi}$ either has the growth vector $(2, 3, 5)$ or is integrable. The former case corresponds to Cartan's celebrated Pfaffian system [4] (for nonintegrable systems or off shell), the latter case corresponds to a dispersionless Lax pair (on shell).

The genericity condition we need here is as follows (and we formulate a similar condition for $d = 3$ which we will use later).

Definition 4. A 2-plane congruence Π is called *nondegenerate* if

$$\begin{aligned} \theta \wedge \theta_\lambda \wedge \theta_{\lambda\lambda} &\neq 0, & \text{where } \text{Ann}(\Pi) &= \langle \theta \rangle & \text{for } d &= 3; \\ \theta \wedge \zeta \wedge \theta_\lambda \wedge \zeta_\lambda &\neq 0, & \text{where } \text{Ann}(\Pi) &= \langle \theta, \zeta \rangle & \text{for } d &= 4. \end{aligned} \quad (15)$$

These conditions depend only on Π , not on the choices of θ or ζ : when $d = 4$ non-degeneracy means equivalently $\varpi_\lambda \wedge \varpi_\lambda \neq 0$ where $\varpi = \theta \wedge \zeta$, or dually that $X \wedge Y \wedge X_\lambda \wedge Y_\lambda \neq 0$, where $\Pi = \langle X, Y \rangle$. If we choose θ and ζ as in (11) and (14), then the nondegeneracy conditions may be written explicitly as:

$$\alpha_\lambda \beta_{\lambda\lambda} - \alpha_{\lambda\lambda} \beta_\lambda \neq 0 \quad \text{for } d = 3; \quad (16)$$

$$\alpha_\lambda \delta_\lambda - \beta_\lambda \gamma_\lambda \neq 0 \quad \text{for } d = 4. \quad (17)$$

Lemma 1. *For $d = 4$, any nondegenerate 2-plane congruence Π has a unique normal lift.*

Proof. If \hat{X} and \hat{Y} are given by (9) and (13), $dx([\hat{X}, \hat{Y}]) = 0 = dy([\hat{X}, \hat{Y}])$ identically, while $dz([\hat{X}, \hat{Y}]) = dt([\hat{X}, \hat{Y}]) = 0$ form two linear equations on m, n :

$$\begin{bmatrix} \delta_\lambda & -\beta_\lambda \\ -\gamma_\lambda & \alpha_\lambda \end{bmatrix} \begin{bmatrix} m \\ n \end{bmatrix} = \begin{bmatrix} \alpha\delta_z + \beta\delta_t - \gamma\beta_z - \delta\beta_t + \beta_y - \delta_x \\ \gamma\alpha_z + \delta\alpha_t - \alpha\gamma_z - \beta\gamma_t + \gamma_x - \alpha_y \end{bmatrix};$$

these have a unique solution by the nondegeneracy condition (17). \square

Example 3. (SDM) We illustrate this with the master equation for SD structures obtained in [12, Theorem 2]. Consider a 2-plane congruence Π spanned by (13) with $\alpha_\lambda = 0 = \delta_\lambda$ and $\beta_\lambda = 1 = -\gamma_\lambda$. This is totally isotropic for the conformal class of the metric

$$g = \theta_\lambda \zeta - \zeta_\lambda \theta = dx(dz + \alpha dx + \gamma dy) + dy(dt + \beta dx + \delta dy),$$

which is independent of λ . In particular, there is a foliation by the totally isotropic level surfaces of (x, y) . Any SD metric can be written in this form, with the isotropic surface foliation being anti-self-dual [12, 31]. The unique normal lift of Π is given by (9) with

$$m = \gamma_x - \alpha_y + \delta\alpha_t - \alpha\gamma_z + \gamma\alpha_z - \beta\gamma_t, \quad n = \delta_x - \beta_y + \delta\beta_t - \alpha\delta_z + \gamma\beta_z - \beta\delta_t.$$

Now the λ^2 term of the integrability condition $\hat{X}(n) - \hat{Y}(m) = 0$ is $(\alpha_z + \gamma_t)_z + (\beta_z + \delta_t)_t = 0$, so $\alpha_z + \gamma_t = s_t$ and $\beta_z + \delta_t = -s_z$ for some function s . However, we may use the translation freedom in λ to set $s = 0$, so that $\alpha = u_t$, $\gamma = -(\lambda + u_z)$, $\beta = \lambda - v_t$, $\delta = v_z$ for functions (u, v) of (x, y, z, t) . Thus we obtain a normal dispersionless pair $\hat{\Pi} = \langle \hat{X}, \hat{Y} \rangle$ with

$$\hat{X} = \partial_x - u_t \partial_z - (\lambda - v_t) \partial_t - Q(u) \partial_\lambda, \quad \hat{Y} = \partial_y + (\lambda + u_z) \partial_z - v_z \partial_t + Q(v) \partial_\lambda,$$

where $Q = \partial_x \partial_z + \partial_y \partial_t - u_t \partial_z^2 + (u_z + v_t) \partial_z \partial_t - v_z \partial_t^2$.

The corresponding quasilinear system (5) is

$$q_x - u_t q_z - (q - v_t) q_t = -Q(u), \quad q_y + (q + u_z) q_z - v_z q_t = Q(v), \quad (18)$$

and the integrability condition reduces to $X(Q(v)) + Y(Q(u)) = 0$, i.e.,

$$\partial_z(Q(u)) = \partial_t(Q(v)), \quad (\partial_x - u_t \partial_z + v_t \partial_t) Q(v) + (\partial_y + u_z \partial_z - v_z \partial_t) Q(u) = 0. \quad (19)$$

Up to some minor coordinate changes, this is the SD master equation (SDM) of [12].

2.3. Integrability, dispersionless Lax pairs and normalization. When $d = 3$, we do not obtain a unique normal lift.

Example 4. (MS) The dispersionless pair (8) for the Manakov–Santini system (6) satisfies

$$[\hat{X}, \hat{Y}] = -G \partial_t - F \partial_\lambda$$

with $F = S(u) + u_t^2$, $G = S(v)$, and so is not normal. However, if we set $\hat{X}' = \hat{X} - G \partial_\lambda$ then $\hat{X}' = \hat{X}$ on shell (when $F = G = 0$), while

$$[\hat{X}', \hat{Y}] = [\hat{X}, \hat{Y}] - G [\partial_\lambda, \hat{Y}] + Y(G) \partial_\lambda = -(F - G_y + (\lambda + v_t)G_t) \partial_\lambda$$

so $\hat{\Pi}' := \langle \hat{X}', \hat{Y} \rangle$ is normal, and is integrable if and only if

$$G_t = 0, \quad G_y = F,$$

i.e., $G = \psi(x, y)$ and $F = \psi_y$. However, this system is not substantively different from the Manakov–Santini system itself, because we can make a point transformation $u \mapsto u - \phi_y(x, y)$, $v \mapsto v - \phi(x, y)$ and if $\phi_{yy} = \psi$, we obtain $F = 0$, $G = 0$.

This example illustrates two important issues that we want to incorporate into the definition of a dispersionless Lax pair $\hat{\Pi}$ for an equation \mathcal{E} : first $\hat{\Pi}$ is only determined modulo \mathcal{E} , and secondly it can be too restrictive in examples to require that the integrability conditions for a dispersionless pair are equivalent to \mathcal{E} .

Definition 5. Let $\mathcal{E} : F = 0$ be a PDE system on u and $\hat{\Pi} \subseteq T\hat{M}_u$ a dispersionless pair.

- A dispersionless pair $\hat{\Pi}' \subseteq T\hat{M}_u$ is \mathcal{E} -equivalent to $\hat{\Pi}$ if $\hat{\Pi} = \hat{\Pi}'$ when $F(u) = 0$.
- $\hat{\Pi}$ is a *dispersionless Lax pair (dLp)* for \mathcal{E} if for any $\hat{\Pi}'$ \mathcal{E} -equivalent to $\hat{\Pi}$, the integrability condition $[\hat{\Pi}', \hat{\Pi}'] = \hat{\Pi}'$ is a nontrivial differential corollary of \mathcal{E} .

To make precise the notion of a differential corollary, we introduce some jet formalism, for which we refer to [21, 22, 36] for further details. A scalar PDE of order ℓ on a manifold M may be defined as an equation of the form

$$F(j^\ell u) = 0 \tag{20}$$

where $F \in C^\infty(J^\ell M)$ is a function on the bundle $\pi_\ell : J^\ell M \rightarrow M$ of ℓ -jets of functions u on M , and $j^\ell u : M \rightarrow J^\ell M$ is the ℓ -jet of u , i.e., in coordinates $j^\ell u = (x, u, \partial u, \dots, \partial^\ell u)$.

In order to discuss objects (such as dLps) depending on an arbitrary finite jet of u , we use the infinite jet bundle $\pi_\infty : J^\infty M \rightarrow M$ which is the union (inverse limit) of $J^k M$ over all k . A function $f : J^\infty M \rightarrow \mathbb{R}$ is smooth if it is the pullback of a function on $J^k M$ for some $k \in \mathbb{N}$, in which case we say f has *order* $\leq k$. A choice of coordinates x^i on M leads to coordinates (x^i, u_α) on $J^\infty M$, where $1 \leq i \leq d$ and α runs over all symmetric multi-indices in d entries. Then $f \in C^\infty(J^\infty M)$ has order $\leq k$ iff it is a function of x^i and u_α for all i and $\alpha = (i_1, \dots, i_j)$ with $|\alpha| = j \leq k$.

The bundle $J^\infty M$ has a canonical flat connection, the *Cartan distribution*, for which the horizontal lift of a vector field X on M is the *total derivative* D_X characterized by $(D_X f) \circ j^\infty u = X(f \circ j^\infty u)$ for any smooth function f on $J^\infty M$. More generally, any section X of $\pi_\infty^* TM$ has a lift to a vector field D_X on $J^\infty M$, given in local coordinates by $D_X = \sum_i a_i D_i$, where $X = \sum_i a_i \partial_i$ and $D_i = \partial_i + \sum_\alpha u_{i\alpha} \partial_{u_\alpha}$.

Higher order operators \square in total derivatives (also known as \mathcal{C} -differential operators) are generated as compositions of the derivations D_X with coefficients being smooth functions on $J^\infty M$. In local coordinates, $\square = \sum a_\alpha D_\alpha$, where $a_\alpha \in C^\infty(J^\infty M)$ and $D_\alpha = D_{i_1} \cdots D_{i_j}$ for a multi-index $\alpha = (i_1, \dots, i_j)$ with entries in $\{1, 2, \dots, d\}$.

Let \mathcal{I}_F be the ideal in $C^\infty(J^\infty M)$ generated by the pullback of $F \in C^\infty(J^\ell M)$ and its total derivatives of arbitrary order. Then the zero-set $\mathcal{E}_\infty \subseteq J^\infty M$ of \mathcal{I}_F is the space of formal solutions of (20): u is a solution of (20) iff $j^\infty u$ is a section of \mathcal{E}_∞ .

These notions extend straightforwardly to PDE systems by replacing $J^\infty M$ with the bundle $\pi_\infty: J^\infty(M, \mathcal{V}) \rightarrow M$ of jets of sections of a fibre bundle $\mathcal{V} \rightarrow M$, and F by a function of order $\leq \ell$ on $J^\infty(M, \mathcal{V})$ with values in a vector bundle $\mathcal{W} \rightarrow M$. The ideal \mathcal{I}_F in $C^\infty(J^\infty(M, \mathcal{V}))$ is now generated by the components of F and their total derivatives of arbitrary order.

In this formalism, a *differential corollary* of $\mathcal{E}: F = 0$ is a subset of \mathcal{I}_F (or, more invariantly, the ideal $\mathcal{I} \subseteq \mathcal{I}_F$ generated by this subset and its total derivatives of arbitrary order). It is *nontrivial* provided it is not a subset of $\mathcal{I}_{F'}$ for any F' whose zero-set in $J^\ell(M, \mathcal{V})$ contains the zero-set of F as a proper (closed) subset. For example, the ideal generated by u_{xy} , for a scalar function $u(x, y, t)$, is not nontrivial as a differential corollary of the system $F(j^1 u) := (u_x, u_y) = 0$, because it is a differential corollary of the equation $F'(j^1 u) := u_x = 0$ properly containing the zero-set of F . On the other hand, the equation $u_{xy} = 0$ is a nontrivial differential corollary of the equation $u_x = 0$.

Consequently, in Definition 5, the integrability condition for a dLp $\hat{\Pi}$ for $\mathcal{E}: F = 0$ need not generate \mathcal{I}_F : indeed, the freedom to replace a dLp by an \mathcal{E} -equivalent one may change the ideal $\mathcal{I} \subseteq \mathcal{I}_F$ that its integrability conditions generate.

Remark 2. In most of the paper we make minimal use of the jet formalism by using the philosophy [21, 36] that a differential equation $\mathcal{E}_\infty \subseteq J^\infty M$ is a generalized manifold whose “points” are solutions u , identified with $M_u = (j^\infty u)(M) \subseteq \mathcal{E}_\infty$ that is diffeomorphic to M via π_∞ . We are justified in working “pointwise” provided there are enough “points” (i.e., for generic $u_\infty \in \mathcal{E}_\infty$ there is a solution u with $u_\infty \in M_u$), and there are existence theorems for hyperbolic PDEs (or rather, ultrahyperbolic PDEs in signature (2, 2)) which assert this in some generality. Nevertheless, we would rather not rely upon such analytical results here, and all our results can be formalized using jets, even if we do not do so explicitly.

The following normalization result now suffices to establish Theorem 2 when $d = 4$.

Proposition 1. *Let $\hat{\Pi}$ be a dLp such that $\Pi = \hat{\pi}_*(\hat{\Pi})$ is nondegenerate. Then $\hat{\Pi}$ is \mathcal{E} -equivalent to a normal dLp. Such a Lax pair for $d = 4$ is unique.*

Proof. When $d = 4$ the Lax pair condition (on shell) implies

$$dz \circ \pi_*[\hat{X}, \hat{Y}] = \square_1 F, \quad dt \circ \pi_*[\hat{X}, \hat{Y}] = \square_2 F$$

for some operators \square_1, \square_2 in total derivatives. Let us modify $\tilde{X} = \hat{X} + A(F)\partial_\lambda$, $\tilde{Y} = \hat{Y} + B(F)\partial_\lambda$, where A, B are operators in total derivatives to be determined (they also depend on λ). The new commutation equation is

$$\begin{aligned} dz \circ \pi_*[\tilde{X}, \tilde{Y}] &= (\square_1 + \alpha_\lambda B - \gamma_\lambda A)F \\ dt \circ \pi_*[\tilde{X}, \tilde{Y}] &= (\square_2 + \beta_\lambda B - \delta_\lambda A)F. \end{aligned}$$

Vanishing of these, equivalent to normality, can be achieved by a unique choice of the operators in total derivatives A, B due to nondegeneracy condition (17).

When $d = 3$, the Lax pair condition (on shell) implies similarly

$$dt \circ \pi_*[\hat{X}, \hat{Y}] = \hat{Y}(\alpha) - \hat{X}(\beta) = \square F$$

for some operator \square in total derivatives. The modification $\tilde{X} = \hat{X} + A(F)\partial_\lambda$, $\tilde{Y} = \hat{Y} + B(F)\partial_\lambda$ gives the new commutation relations

$$dt \circ \pi_*[\tilde{X}, \tilde{Y}] = (\square + \alpha_\lambda B - \beta_\lambda A)F.$$

The equation $\beta_\lambda A - \alpha_\lambda B = \square$ admits the solution $A = \alpha_{\lambda\lambda}\square/(\beta_\lambda\alpha_{\lambda\lambda} - \alpha_\lambda\beta_{\lambda\lambda})$ and $B = \beta_{\lambda\lambda}\square/(\beta_\lambda\alpha_{\lambda\lambda} - \alpha_\lambda\beta_{\lambda\lambda})$ by (16), unique up to the freedom $(A, B) \mapsto (A, B) + (\alpha_\lambda, \beta_\lambda)L$. \square

3. The Characteristic Condition for Dispersionless Lax Pairs

3.1. Symbols and the characteristic condition. In order to prove Theorem 1 in full generality, we need the notions of symbol and characteristic variety for a general PDE system. For this we use the jet formalism. Recall from the previous section that a smooth function F on $J^\infty M$ has order $\leq \ell$ if it is a pullback from $J^\ell M$, and that $J^\infty M$ has a canonical connection, the Cartan distribution. The vertical part of the 1-form $dF \in \Omega^1(J^\infty M)$ may be viewed in coordinates as a polynomial on $\pi_\infty^* T^*M$ given by

$$\sum_{j=0}^{\ell} F_{(j)} \quad \text{where} \quad F_{(j)} = \sum_{|\alpha|=j} (\partial_{u_\alpha} F) \partial_\alpha \quad \text{is a section of} \quad \pi_\infty^* S^j T^*M.$$

The top degree term $\sigma_F = F_{(\ell)}$, called the (order ℓ) *symbol* of F , is independent of coordinates. We assume it is nonvanishing: if it vanishes, F has order $\leq \ell - 1$ and σ_F has lower degree.

This generalizes to a PDE system of order ℓ , i.e., a function F of order $\leq \ell$ on $J^\infty(M, \mathcal{V})$, for some fibre bundle \mathcal{V} , with values in a vector bundle $\mathcal{W} \rightarrow M$. The symbol σ_F of F is then a homogeneous degree ℓ polynomial on $\pi_\infty^* T^*M$ with values in $\text{Hom}(T\mathcal{V}, \mathcal{W})$, which we assume is not identically zero, so that the PDE system does not have order $\leq \ell - 1$. The characteristic variety of the PDE system $\mathcal{E} : F = 0$ is defined by [34]

$$\chi^{\mathcal{E}} = \{[\theta] \in \mathbb{P}(\pi_\infty^* T^*M) \mid \sigma_F(\theta) \text{ is not injective}\}.$$

If \mathcal{V} and \mathcal{W} have the same rank, then $[\theta]$ is characteristic iff $\sigma_F(\theta)$ is not surjective. We take $\text{rank}(\mathcal{V}) = \text{rank}(\mathcal{W})$ as the definition of a determined system, although a more proper definition is $\text{codim } \chi^{\mathcal{E}} = 1$.

Definition 6. We say that a 2-plane congruence Π (or a dLp $\hat{\Pi}$) is *characteristic for* \mathcal{E} if for any solution u of \mathcal{E} and any $\theta \in \text{Ann}(\Pi) \subseteq \hat{\pi}^* T^*M_u$, we have $[\theta] \in \chi^{\mathcal{E}}$.

In the jet formalism, a dispersionless pair $\hat{\Pi}$ lives on a rank 1-bundle $\hat{\pi} : \hat{M} \rightarrow J^\infty(M, \mathcal{V})$ (so that $\hat{M}_u = (j^\infty u)^* \hat{M}$) and we let $\hat{\pi}_\infty = \pi_\infty \circ \hat{\pi} : \hat{M} \rightarrow M$. A 2-plane congruence Π is then a rank 2 subbundle of $\hat{\pi}_\infty^* T^*M$, and $\hat{\Pi}$ is a lift of Π to $T\hat{M}$. In practice we use a spectral parameter λ to trivialize \hat{M} over $J^\infty(M, \mathcal{V})$. Then $T\hat{M}$ is the direct sum of the vertical bundle of $\hat{\pi}$, spanned by ∂_λ , and $\hat{\pi}^* T J^\infty(M, \mathcal{V})$. Thus if Π is spanned by $X, Y \in \hat{\pi}_\infty^* T^*M$, we may write the dispersionless pair $\hat{\Pi}$ as the span of

$\hat{X} = D_X + m \partial_\lambda$ and $\hat{Y} = D_Y + n \partial_\lambda$, where D_X and D_Y are total derivatives (depending also on λ) and m, n are functions on \hat{M} . Then

$$[\hat{X}, \hat{Y}] = ([D_X, D_Y] + m D_{\partial_\lambda Y} - n D_{\partial_\lambda X}) + (D_X n - D_Y m + m \partial_\lambda n - n \partial_\lambda m) \partial_\lambda.$$

The integrability condition $[\hat{X}, \hat{Y}] \in \Gamma(\hat{\Pi})$ reduces to $[X, Y] + m \partial_\lambda Y - n \partial_\lambda X = \nu_X X + \nu_Y Y$, for some ν_X, ν_Y , together with the vanishing of $D_X n - D_Y m + m \partial_\lambda n - n \partial_\lambda m - \nu_X m - \nu_Y n$. As in the previous section, we may choose X and Y so that $\nu_X = \nu_Y = 0$, and hence the Lax equation (split into the vertical and horizontal parts) becomes the system

$$D_X n - D_Y m + m \partial_\lambda n - n \partial_\lambda m = 0, \quad (21)$$

$$D_X Y - D_Y X + m \partial_\lambda Y - n \partial_\lambda X = 0. \quad (22)$$

We thus have a dLp for \mathcal{E} if these equations hold modulo \mathcal{I}_F i.e., all components (and hence their total derivatives of arbitrary order) belong to \mathcal{I}_F .

Lemma 2. *If $D_X q - D_Y p$ has order $\leq k$, for functions p, q of $u_\infty \in J^\infty(M, \mathcal{V})$ and sections X, Y of $\pi_\infty^* TM$, then its order k symbol is*

$$D_{X^{(k)}} q + D_X(q^{(k)}) + X \odot q^{(k-1)} - D_{Y^{(k)}} p - D_Y(p^{(k)}) - Y \odot p^{(k-1)}. \quad (23)$$

If X, Y are linearly independent, and P_1 and P_2 are symmetric k -vectors with $X \odot P_2 = Y \odot P_1$, there is a symmetric $(k-1)$ -vector S with $P_1 = X \odot S$ and $P_2 = Y \odot S$.

Proof. Equation (23) is straightforward from the definition of the total derivative and the product rule for the vertical differentiation. Extending X, Y pointwise to a basis, the second part reduces to the trivial observation that for any homogeneous polynomials $P_j = P_j(\xi_1, \dots, \xi_d)$, $j = 1, 2$, with $\xi_1 P_2 = \xi_2 P_1$, there is a homogeneous polynomial P with $P_j = \xi_j P$. \square

Lemma 3. *Let (21)–(22) have order $\leq k+1$ modulo \mathcal{I}_F , i.e., all their higher symbols vanish modulo \mathcal{I}_F . Then there is a symmetric k -tensor S_k and a symmetric TM -valued k -tensor Q_k such that, modulo \mathcal{I}_F , the order $k+1$ symbols of (21) and (22) are respectively*

$$\begin{aligned} & X \odot (n^{(k)} + D_{Q_k} n - D_Y S_k + S_k \partial_\lambda n - n \partial_\lambda S_k) \\ & - Y \odot (m^{(k)} + D_{Q_k} m - D_X S_k + S_k \partial_\lambda m - m \partial_\lambda S_k), \end{aligned} \quad (24)$$

$$\begin{aligned} & X \odot (Y^{(k)} + D_{Q_k} Y - D_Y Q_k + S_k \partial_\lambda Y - n \partial_\lambda Q_k) \\ & - Y \odot (X^{(k)} + D_{Q_k} X - D_X Q_k + S_k \partial_\lambda X - m \partial_\lambda Q_k). \end{aligned} \quad (25)$$

Proof. Suppose that X, Y, m, n depend only on the N -jet of u for some $N \in \mathbb{N}$, so that (21)–(22) have order $\leq N+1$, and it suffices to prove the lemma for $k \leq N$. We thus induct on $p = N - k$. For $p = 0$, the order $k+1 = N+1$ symbols of (21) and (22) are simply $X \odot n^{(k)} - Y \odot m^{(k)}$ and $X \odot Y^{(k)} - Y \odot X^{(k)}$ by (23), so we are done, with $S_k = 0 = Q_k$.

Now suppose that the lemma holds with $k = N - p$ for some $p \geq 0$, and suppose that (21)–(22) have order $\leq k$ modulo \mathcal{I}_F . Then (21) certainly has order $\leq k+1$ modulo \mathcal{I}_F , and so the inductive hypothesis implies its order $k+1$ symbol, which vanishes

modulo \mathcal{I}_F , is given by (24). Hence Lemma 2 produces a symmetric $(k-1)$ -tensor S_{k-1} such that, modulo \mathcal{I}_F ,

$$\begin{aligned} m^{(k)} &= X \odot S_{k-1} - D_{Q_k} m + D_X S_k - S_k \partial_\lambda m + m \partial_\lambda S_k, \\ n^{(k)} &= Y \odot S_{k-1} - D_{Q_k} n + D_Y S_k - S_k \partial_\lambda n + n \partial_\lambda S_k. \end{aligned}$$

Similarly, by (25), there is a symmetric TM -valued $(k-1)$ -tensor Q_{k-1} such that

$$\begin{aligned} X^{(k)} &= X \odot Q_{k-1} - D_{Q_k} X + D_X Q_k - S_k \partial_\lambda X + m \partial_\lambda Q_k, \\ Y^{(k)} &= Y \odot Q_{k-1} - D_{Q_k} Y + D_Y Q_k - S_k \partial_\lambda Y + n \partial_\lambda Q_k, \end{aligned}$$

modulo \mathcal{I}_F . By (23), the order k symbol of (21) is

$$\begin{aligned} D_{X^{(k)}} n + D_X(n^{(k)}) + X \odot n^{(k-1)} - D_{Y^{(k)}} m - D_Y(m^{(k)}) - Y \odot m^{(k-1)} \\ + m^{(k)} \partial_\lambda n + m \partial_\lambda(n^{(k)}) - n^{(k)} \partial_\lambda m - n \partial_\lambda(m^{(k)}). \end{aligned}$$

Hence, substituting for $X^{(k)}$, $Y^{(k)}$, $m^{(k)}$, $n^{(k)}$, we have

$$\begin{aligned} D_{X \odot Q_{k-1} - D_{Q_k} X + D_X Q_k - S_k \partial_\lambda X + m \partial_\lambda Q_k} n - D_{Y \odot Q_{k-1} - D_{Q_k} Y + D_Y Q_k - S_k \partial_\lambda Y + n \partial_\lambda Q_k} m \\ - D_Y(X \odot S_{k-1} - D_{Q_k} m + D_X S_k - S_k \partial_\lambda m + m \partial_\lambda S_k) \\ + D_X(Y \odot S_{k-1} - D_{Q_k} n + D_Y S_k - S_k \partial_\lambda n + n \partial_\lambda S_k) \\ + (X \odot S_{k-1} - D_{Q_k} m + D_X S_k - S_k \partial_\lambda m + m \partial_\lambda S_k) \partial_\lambda n \\ - (Y \odot S_{k-1} - D_{Q_k} n + D_Y S_k - S_k \partial_\lambda n + n \partial_\lambda S_k) \partial_\lambda m \\ - n \partial_\lambda(X \odot S_{k-1} - D_{Q_k} m + D_X S_k - S_k \partial_\lambda m + m \partial_\lambda S_k) + X \odot n^{(k-1)} \\ + m \partial_\lambda(Y \odot S_{k-1} - D_{Q_k} n + D_Y S_k - S_k \partial_\lambda n + n \partial_\lambda S_k) - Y \odot m^{(k-1)}. \end{aligned}$$

A lot of cancellation now occurs to leave

$$\begin{aligned} X \odot (n^{(k-1)} + D_{Q_{k-1}} n - D_Y S_{k-1} + S_{k-1} \partial_\lambda n - n \partial_\lambda S_{k-1}) \\ - Y \odot (m^{(k-1)} + D_{Q_{k-1}} m - D_X S_{k-1} + S_{k-1} \partial_\lambda m - m \partial_\lambda S_{k-1}) \\ + (D_X Y - D_Y X + m \partial_\lambda Y - n \partial_\lambda X) \odot S_{k-1} + (D_X n - D_Y m + m \partial_\lambda n - n \partial_\lambda m) \partial_\lambda S_k \\ + D_{D_X Y - D_Y X + m \partial_\lambda Y - n \partial_\lambda X} S_k - (D_{Q_k} + S_k \partial_\lambda)(D_X n - D_Y m + m \partial_\lambda n - n \partial_\lambda m) \end{aligned}$$

and the last two lines vanish modulo \mathcal{I}_F , which establishes (24) for $k' = N - (p+1) = k-1$. We turn now to the order k symbol of (22), which, by (23), is

$$\begin{aligned} D_{X^{(k)}} Y + D_X(Y^{(k)}) + X \odot Y^{(k-1)} - D_{Y^{(k)}} X - D_Y(X^{(k)}) - Y \odot X^{(k-1)} \\ + m^{(k)} \partial_\lambda Y + m \partial_\lambda Y^{(k)} - n^{(k)} \partial_\lambda X - n \partial_\lambda X^{(k)}. \end{aligned}$$

Hence, substituting for $X^{(k)}$, $Y^{(k)}$, $m^{(k)}$, $n^{(k)}$, we have, modulo \mathcal{I}_F ,

$$\begin{aligned} 0 &= D_{X \odot Q_{k-1} - D_{Q_k} X + D_X Q_k - S_k \partial_\lambda X + m \partial_\lambda Q_k} Y - D_{Y \odot Q_{k-1} - D_{Q_k} Y + D_Y Q_k - S_k \partial_\lambda Y + n \partial_\lambda Q_k} X \\ &\quad - D_Y(X \odot Q_{k-1} - D_{Q_k} X + D_X Q_k - S_k \partial_\lambda X + m \partial_\lambda Q_k) \\ &\quad + D_X(Y \odot Q_{k-1} - D_{Q_k} Y + D_Y Q_k - S_k \partial_\lambda Y + n \partial_\lambda Q_k) \\ &\quad + (X \odot S_{k-1} - D_{Q_k} m + D_X S_k - S_k \partial_\lambda m + m \partial_\lambda S_k) \partial_\lambda Y \\ &\quad - (Y \odot S_{k-1} - D_{Q_k} n + D_Y S_k - S_k \partial_\lambda n + n \partial_\lambda S_k) \partial_\lambda X \\ &\quad - n \partial_\lambda(X \odot Q_{k-1} - D_{Q_k} X + D_X Q_k - S_k \partial_\lambda X + m \partial_\lambda Q_k) + X \odot Y^{(k-1)} \\ &\quad + m \partial_\lambda(Y \odot Q_{k-1} - D_{Q_k} Y + D_Y Q_k - S_k \partial_\lambda Y + n \partial_\lambda Q_k) - Y \odot X^{(k-1)} \end{aligned}$$

$$\begin{aligned}
&= X \odot (Y_{(k-1)} + D_{Q_{k-1}}Y - D_Y Q_{k-1} + S_{k-1} \partial_\lambda Y - n \partial_\lambda Q_{k-1}) \\
&\quad - Y \odot (X_{(k-1)} + D_{Q_{k-1}}X - D_X Q_{k-1} + S_{k-1} \partial_\lambda X - m \partial_\lambda Q_{k-1}) \\
&\quad + (D_X Y - D_Y X + m \partial_\lambda Y - n \partial_\lambda X) \odot Q_{k-1} \\
&\quad + (D_X n - D_Y m + m \partial_\lambda n - n \partial_\lambda m) \partial_\lambda Q_k, \\
&\quad + D_{D_X Y - D_Y X + m \partial_\lambda Y - n \partial_\lambda X} Q_k - (D_{Q_k} + S_k \partial_\lambda)(D_X Y - D_Y X + m \partial_\lambda Y - n \partial_\lambda X)
\end{aligned}$$

and the last two lines again vanish modulo \mathcal{I}_F , so that (25) holds for $k' = N - (p+1) = k - 1$, completing the proof. \square

3.2. Proof of Theorem 1. The characteristic property depends only on the dispersionless pair $\hat{\Pi}$ up to \mathcal{E} -equivalence, so the strategy is to take $\hat{\Pi}$ to have integrability condition of minimal order within its \mathcal{E} -equivalence class.

As above, we let \mathcal{I}_F be the ideal generated by F and its total derivatives, and assume that $\hat{\Pi}$ is spanned by vector fields $D_X + m \partial_\lambda$ and $D_Y + n \partial_\lambda$ which commute on shell, where X, Y, m, n depend only on the N -jet of u for some $N \in \mathbb{N}$.

By the definition of a dLp, equations (21)–(22) have the form $\Lambda_1(F) = 0$ and $\Lambda_2(F) = 0$, where Λ_1 and Λ_2 are λ -dependent operators in total derivatives on the codomain \mathcal{W} of F , the latter operator being TM -valued. We suppose that $\Lambda_1(F)$ and $\Lambda_2(F)$ both have order $\leq k+1$ modulo \mathcal{I}_F and that k is minimal with this property among all dispersionless pairs \mathcal{E} -equivalent to $\hat{\Pi}$. Note that $k+1 \geq \ell$, the order of F .

In local coordinates we may write Λ_1 as a finite sum $\sum_\alpha b_\alpha(u_\infty, \lambda) D_\alpha$ where each $b_\alpha(u_\infty, \lambda)$ is a section of \mathcal{W}^* . Then for any $r \geq \ell$, the order r symbol of $\Lambda_1(F)$ satisfies

$$\begin{aligned}
\Lambda_1(F)_{(r)} &= \sum_\alpha b_\alpha(u_\infty, \lambda) (D_\alpha F)_{(r)} \pmod{\mathcal{I}_F} \\
&= \sum_{j=0}^{\ell} \sum_{|\alpha|=r-j} \partial_\alpha \odot b_\alpha(u_\infty, \lambda) F_{(j)} \pmod{\mathcal{I}_F}
\end{aligned}$$

(using the product rule for vertical differentiation). Let $r = \max\{|\alpha| : b_\alpha \notin \mathcal{I}_F\} + \ell$. Then $\Lambda_1(F)_{(r)} = \sum_{|\alpha|=r-\ell} \partial_\alpha \odot b_\alpha(u_\infty, \lambda) \sigma_F \pmod{\mathcal{I}_F}$. Since $\chi^\mathcal{E}$ is (fibrewise) a proper closed variety, $\sigma_F(\theta)$ is surjective for generic θ , and by definition of r , there exists such a θ with $\sum_{|\alpha|=r-\ell} \partial_\alpha(\theta) b_\alpha(u_\infty, \lambda) \notin \mathcal{I}_F$. Hence $\Lambda_1(F)_{(r)}(\theta) \notin \mathcal{I}_F$, and so we must have $r \leq k+1$ because $\Lambda_1(F)$ has order $\leq k+1$ modulo \mathcal{I}_F .

Thus for all $|\alpha| \geq k - \ell + 2$, we have $b_\alpha \in \mathcal{I}_F$ by definition of r , and hence $\Lambda_1(F)_{(k+1)} = L_1 \odot \sigma_F \pmod{\mathcal{I}_F}$, where L_1 is a \mathcal{W}^* -valued symmetric $(k - \ell + 1)$ -vector depending on (u_∞, λ) . An analogous argument shows $\Lambda_2(F)_{(k+1)} = L_2 \odot \sigma_F \pmod{\mathcal{I}_F}$ for a $TM \otimes \mathcal{W}^*$ -valued symmetric $(k - \ell + 1)$ -vector L_2 depending on (u_∞, λ) .

By Lemma 3, the order $k+1$ symbols of $\Lambda_1(F), \Lambda_2(F)$ have the forms (24)–(25) modulo \mathcal{I}_F . Hence, on any solution u we have that for all $\theta \in \text{Ann}(\Pi)$

$$L_1(\theta) \circ \sigma_F(\theta) = 0, \quad L_2(\theta) \circ \sigma_F(\theta) = 0 \quad (26)$$

(there is only one independent θ at each point for $d = 3$ and a pair of such for $d = 4$).

Suppose for contradiction that $\sigma_F(\theta)$ is surjective for some $\theta \in \text{Ann}(\Pi)$ —hence almost every such θ , since surjectivity is an open condition. Then (26) implies on shell that $L_1(\theta) = 0$ and $L_2(\theta) = 0$ identically (as degree $k+1 - \ell$ polynomials) in $\theta \in \text{Ann}(\Pi)$. If $k+1 = \ell$ this immediately yields that $(L_1, L_2) = 0$ modulo \mathcal{I}_F , a contradiction. For

$k \geq \ell$ we have instead $L_1 = X \odot T_1 - Y \odot U_1$ and $L_2 = X \odot T_2 - Y \odot U_2$ modulo \mathcal{I}_F for some symmetric $(k - \ell)$ -vectors T_1, U_1 with values in \mathcal{W}^* and T_2, U_2 with values in $TM \otimes \mathcal{W}^*$.

We now let $\tau_1, \nu_1, \tau_2, \nu_2$ be order $k - \ell$ operators in total derivatives such that $\tau_1 F$ has order k symbol $T_1 \odot \sigma_F$ modulo \mathcal{I}_F and so on: concretely, in local coordinates, if $T_1 = \sum_{|\alpha|=k-\ell} t_\alpha(u_\infty, \lambda) \partial_\alpha$, we may take $\tau_1 = \sum_{|\alpha|=k-\ell} t_\alpha(u_\infty, \lambda) D_\alpha$. We then modify the dispersionless pair by $m \mapsto m - \nu_1(F), n \mapsto n - \tau_1(F), X \mapsto X - \nu_2(F), Y \mapsto Y - \tau_2(F)$.

This modification is \mathcal{E} -equivalent to $\hat{\Pi}$, but the new order $k + 1$ symbols of (21)–(22) vanish modulo \mathcal{I}_F , so they have order $\leq k$ modulo \mathcal{I}_F . This contradicts minimality of k .

Thus $\sigma_F(\theta)$ is not surjective for any $\theta \in \text{Ann}(\Pi)$. Since the PDE system is determined this implies that θ is characteristic, and we are done. \square

3.3. Dispersionless pairs characteristic for a quadric. If $\hat{\Pi}$ is a dLp for an equation \mathcal{E} whose characteristic variety $\chi^\mathcal{E}$ is a quadric, then Π is coisotropic for this quadric by Theorem 1. In this section we investigate the extent to which Π recovers this quadric. We begin with a uniqueness criterion, and then discuss existence. We make essential use of the nondegeneracy conditions (16)–(17), which imply in particular that at each $\mathbf{x} \in M_u$, the image of $\Pi_{\mathbf{x}}: \lambda \mapsto \Pi_{(\mathbf{x}, \lambda)}$ does not lie in any proper projective linear subspace of $\text{Gr}_2(T_{\mathbf{x}}M_u) \subseteq \mathbb{P}(\wedge^2 T_{\mathbf{x}}M_u)$.

Proposition 2. *If a 2-plane congruence Π is coisotropic for c_F , then for any $\mathbf{x} \in M_u$ and $\lambda \in \hat{\pi}^{-1}(\mathbf{x})$ at which $\Pi_{\mathbf{x}}$ is an immersion, it is nondegenerate at \mathbf{x} . Conversely, at any point \mathbf{x} where Π is nondegenerate, there is at most one (quadratic) conformal structure c_F on $T_{\mathbf{x}}M_u$ with $\Pi_{(\mathbf{x}, \lambda)}$ coisotropic for all λ , and it must be nondegenerate and hyperbolic.*

Proof. Suppose first that $d = 3$, so that $\text{Gr}_2(T_{\mathbf{x}}M_u) \cong \mathbb{P}(T_{\mathbf{x}}^*M_u)$ is a projective plane, and $\text{Ann}(\Pi_{\mathbf{x}})$ is a curve in this plane. If Π is coisotropic, then $\text{Ann}(\Pi_{\mathbf{x}})$ lies on the nonsingular conic $\{[\theta] : \sigma_F(\theta) = 0\}$ and so if $\Pi_{\mathbf{x}}$ is immersed, its derivatives of order ≤ 2 in λ span $\mathbb{P}(T_{\mathbf{x}}^*M_u)$, hence it is nondegenerate. Conversely, two distinct nonsingular conics meet in at most four points, so $\text{Ann}(\Pi_{\mathbf{x}})$ lies on at most one nonsingular conic (which is nonempty, hence hyperbolic), and if $\text{Ann}(\Pi_{\mathbf{x}})$ lies on a singular conic, it lies on a line, hence $\Pi_{\mathbf{x}}$ is degenerate.

Suppose instead that $d = 4$, so that (the Plücker embedding of) $\text{Gr}_2(T_{\mathbf{x}}M_u)$ is the Klein quadric in $\mathbb{P}(\wedge^2 T_{\mathbf{x}}M_u)$, and $\wedge^2 \Pi_{\mathbf{x}}$ is a curve in this quadric. If Π is coisotropic, then $\wedge^2 \Pi_{\mathbf{x}}$ lies in a nondegenerate plane section of this quadric, which is a conic: the corresponding lines in $\mathbb{P}(T_{\mathbf{x}}^*M_u)$ belong to one of the rulings of the quadric surface $\{[\theta] : \sigma_F(\theta) = 0\}$ in $\mathbb{P}(T_{\mathbf{x}}^*M_u)$. In particular, if $\Pi_{\mathbf{x}}$ is immersed, its tangent does not lie in the quadric, hence it is nondegenerate. Conversely, two distinct nonsingular quadric surfaces meet in a degree four curve (containing at most four lines), so if $\Pi_{\mathbf{x}}$ is nonconstant, it lies on at most one nonsingular quadric surface (which is hyperbolic because it contains lines), and if $\Pi_{\mathbf{x}}$ has image in a singular quadric surface, then the lines pass through a point or lie in a plane, hence $\wedge^2 \Pi_{\mathbf{x}}$ lies in a proper projective linear subspace of $\text{Gr}_2(T_{\mathbf{x}}M_u)$, hence $\Pi_{\mathbf{x}}$ is degenerate. \square

Proposition 3. *Suppose $d = 3$ and the nondegeneracy condition (16) holds. Then there is a unique conformal structure c for which the 2-plane congruence $\Pi = \langle X, Y \rangle$ is null for all λ if and only if the Monge invariant $I(\alpha, \beta) = 0$. This invariant has order 5 in the*

entries and it distinguishes conics in the projective plane. In the local parametrization with $\beta = \lambda$, this condition is the following (we denote $\alpha' = \alpha_\lambda$ etc.):

$$I(\alpha, \lambda) = 9(\alpha'')^2 \alpha^{(5)} - 45\alpha'' \alpha''' \alpha^{(4)} + 40(\alpha''')^2 = 0.$$

Suppose $d = 4$ and the nondegeneracy condition (17) holds. Then there is a unique conformal structure c for which the 2-plane congruence $\Pi = \langle X, Y \rangle$ is (co)isotropic for all λ if and only if the following system of differential equations of order 3 holds, which we write in a partially integrated second order form so (again $\alpha' = \alpha_\lambda$ etc.)

$$v'w'' - v''w' = k_{vw}|\alpha'\delta' - \beta'\gamma'|^{3/2} \text{ for } v, w \in \{\alpha, \beta, \gamma, \delta\},$$

where k_{vw} are λ -independent and satisfy the ‘‘cocycle conditions’’ $k_{vw} + k_{wv} = 0$, $u'k_{vw} + v'k_{wu} + w'k_{uv} = 0$ for $u, v, w \in \{\alpha, \beta, \gamma, \delta\}$. In the normalization $\delta = \lambda$ these conditions simplify to: $(\alpha, \beta, \gamma)'' = \mathbf{v}|\alpha' - \beta'\gamma'|^{3/2}$, where \mathbf{v} is a λ -independent 3-component vector.

Proof. Let us discuss first the case $d = 3$. We are looking for a conformal structure c , represented by a pseudo-Riemannian metric g of signature $(2, 1)$, such that the planes $\Pi = \langle X, Y \rangle$ are null. Consider the Pfaffian form $\theta = dt + \alpha dx + \beta dy \in \text{Ann}(\Pi)$. The null condition is a single equation $c(\theta, \theta) = 0$. Adding to it its λ -derivatives up to order 4, we get a system of 5 equations on 6 coefficients of the metric (5 coefficients if considered up to proportionality). This system is solvable iff (16) holds. Provided this nondegeneracy condition, we can uniquely find $c = [g]$, but in order for it to be supported on M_u (and not on \hat{M}_u) the ratio of the coefficients of g must be λ -independent. This is equivalent to the condition $I(\alpha, \beta) = 0$.

Consider now the case $d = 4$. Add to the 3 equations $c(X, X) = 0$, $c(X, Y) = 0$, $c(Y, Y) = 0$ their first and second derivatives in λ . The obtained system of 9 equations on 10 coefficients of the metric (9 coefficients if considered up to proportionality) is solvable iff condition (17) holds. Provided this nondegeneracy condition, we can uniquely find $c = [g]$, but in order for it to be supported on M_u (and not on \hat{M}_u) the ratio of the coefficients of g must be λ -independent. This is equivalent to the system of equations formulated in the proposition. \square

4. Projective Dependence on the Spectral Parameter

4.1. Weyl connections and standard dLps. For any 2-plane congruence Π which is characteristic for a bundle of nonsingular quadric hypersurfaces, there is a well-known construction of lifts $\hat{\Pi}_\nabla$ of Π from Weyl connections ∇ , i.e., torsion-free connections preserving the conformal structure c defining the quadric. Such Weyl connections form an affine space modelled on the vector space of 1-forms on M_u .

Lemma 4. *Let Π be a nondegenerate 2-plane congruence on $\hat{M}_u \rightarrow M_u$, characteristic for a bundle of quadric hypersurfaces, and ∇ a Weyl connection. Then ∇ induces a connection on \hat{M}_u such that the horizontal lift $\hat{\Pi}_\nabla$ of Π is normal.*

Proof. Since ∇ is a conformal connection, it induces a connection on the bundle of coisotropic planes for c , and hence on \hat{M}_u , since Π is an immersion. The pullback of ∇ to $\hat{\pi}^*T M_u$ preserves Π and hence, since ∇ is torsion-free, the horizontal lift $\hat{\Pi}$ satisfies $\hat{\pi}_*[\hat{\Pi}, \hat{\Pi}] = \Pi$. \square

We refer to such a lift $\hat{\Pi}_\nabla$ as a *standard dLp*. For $d = 4$, any standard dLp $\hat{\Pi}_\nabla$ is the unique normal lift of $\Pi = \hat{\pi}_*(\hat{\Pi}_\nabla)$, hence independent of the choice of Weyl connection, as is well known [29]. However, for both $d = 3$ and $d = 4$, standard dLps are very special because the connection induced by ∇ on \hat{M}_u is *projective*: \hat{M}_u is locally isomorphic to a \mathbb{P}^1 -bundle over M_u and if λ is a spectral parameter induced by an affine coordinate on this projective bundle, then horizontal lifts of (λ -independent) vector fields on M_u depend quadratically on λ (because vector fields on \mathbb{P}^1 have this form in an affine chart).

Furthermore, with respect to such a projective spectral parameter λ , there is a local parametrization of vector fields spanning Π that is linear in λ , i.e., $\Pi = \langle V_1 + \lambda V_3, V_2 + \lambda V_4 \rangle$ for λ -independent vector fields V_i on M_u , so that their lifts are cubic in λ .

When $d = 4$, these properties follow from the existence of an adapted frame V_1, V_2, V_3, V_4 for M_u such that in the dual coframe $\theta_1, \theta_2, \theta_3, \theta_4$, the conformal structure is represented by $g = \theta_1\theta_4 - \theta_2\theta_3$. Then (up to a choice of orientation) $\Pi = \langle V_1 + \lambda V_3, V_2 + \lambda V_4 \rangle$, $\Delta = \langle V_1 + \lambda V_3, V_2 + \lambda V_4, \partial_\lambda \rangle$ and it is straightforward to verify that the unique normal lift of Π is

$$\hat{\Pi} = \langle V_1 + \lambda V_3 + m\partial_\lambda, V_2 + \lambda V_4 + n\partial_\lambda \rangle,$$

where the coefficients m, n are given in terms of the structure functions $c_{ij}^k = \theta_k([V_i, V_j])$ of the frame as

$$\begin{aligned} m &= -c_{12}^4 + \lambda(c_{23}^4 - c_{14}^4 + c_{12}^2) - \lambda^2(c_{23}^2 - c_{14}^2 + c_{34}^4) + \lambda^3 c_{34}^2, \\ n &= c_{12}^3 - \lambda(c_{23}^3 - c_{14}^3 + c_{12}^1) + \lambda^2(c_{23}^1 - c_{14}^1 + c_{34}^3) - \lambda^3 c_{34}^1. \end{aligned}$$

These are cubic in λ as required, and compatible with the representation $m = m_1 + \lambda m_3$ and $n = m_2 + \lambda m_4$ for coefficients m_i of ∂_λ in the lifts of V_i that are quadratic in λ .

When $d = 3$, there is similarly an adapted frame V_0, V_1, V_2 on M_u with the dual coframe $\theta_0, \theta_1, \theta_2$ such that conformal structure c_F is represented by the Lorentzian metric $g = 4\theta_0\theta_2 - \theta_1^2$ and $\Pi = \langle V_0 + \lambda V_1, V_1 + \lambda V_2 \rangle = \ker \theta(\lambda)$, where

$$\theta(\lambda) = \theta_2 - \lambda\theta_1 + \lambda^2\theta_0 \quad (27)$$

for a (projective) spectral parameter λ . We then have the following fact (cf. [14]).

Lemma 5. *Let $d = 3$ and let Π be as in Lemma 4. Then Weyl connections parametrize projective normal lifts $\hat{\Pi}$ of Π .*

Proof. By definition any projective lift given by (9), with $X = V_0 + \lambda V_1$, $Y = V_1 + \lambda V_2$ affine linear, has m, n cubic in λ , i.e., $m = \sum_{i=0}^3 m_i \lambda^i$, $n = \sum_{i=0}^3 n_i \lambda^i$. Now $\text{Ann}(\Pi)$ is spanned by the 1-form (27) where $\theta_i(V_j) = \delta_{ij}$. Hence $\theta(\hat{\pi}_*[\hat{X}, \hat{Y}]) = \theta([X, Y] - nV_1 + mV_2)$ is a quartic polynomial in λ determining 5 of the 8 coefficients of m and n . It is straightforward to check that remaining three coefficients are determined uniquely by the Weyl connection (a 1-form has three components at each point). \square

4.2. The modified Manakov–Santini master equation in 3D. As mentioned in the introduction, the integrability for a standard dLp $\hat{\Pi}_\nabla$ in 3D is well-known to be equivalent to the EW equation on (c, ∇) and has the geometric interpretation that any EW manifold locally admits (many) foliations by totally geodesic null surfaces [5, 19] (corresponding

to curves in the minitwistor space). We now use this to obtain an alternative derivation of the Manakov–Santini system [24] as a master equation in 3D, or rather a modification of this system which was previously derived in [12] by a different method.

Any totally geodesic null surface has a canonical foliation by null geodesics, so any EW manifold admits a local coordinate system (x, y, t) , where x and y are pulled back from local coordinates on the local leaf spaces of a totally geodesic null surface foliation and the induced null geodesic foliation respectively. Thus ∂_t is null and orthogonal to ∂_y and we can use the freedom in the t coordinate so that the conformal structure has a representative metric

$$g = 4(dt - b dx)dx - (dy - a dx)^2 \quad (28)$$

for some functions a and b . This has the form $4\theta_0\theta_2 - \theta_1^2$, where $\theta_0 = dx$, $\theta_1 = -dy + a dx$, $\theta_2 = dt - b dx$ is the coframe dual to $V_0 = \partial_x + a\partial_y + b\partial_t$, $V_1 = -\partial_y$ and $V_2 = \partial_t$. Thus the null 2-plane congruence $\Pi = \langle V_0 + \lambda V_1, V_1 + \lambda V_2 \rangle$ is the kernel of

$$\theta(\lambda) = (dt - b dx) + \lambda(dy - a dx) + \lambda^2 dx = dt + \lambda dy + (\lambda^2 - a\lambda - b)dx, \quad (29)$$

and is equal to W_0^\perp where

$$W_0 = V_0 + 2\lambda V_1 + \lambda^2 V_2 = \partial_x + a\partial_y + b\partial_t - 2\lambda\partial_y + \lambda^2\partial_t = \partial_x + (a - 2\lambda)\partial_y + (b + \lambda^2)\partial_t. \quad (30)$$

Since ∂_y and ∂_t are tangent to level surfaces of x , which are the null surfaces corresponding to $\lambda = \infty$, the standard dLp must have the form $\hat{\Pi}_\nabla = \langle V_0 + \lambda V_1 + m'\partial_\lambda, V_1 + \lambda V_2 + n'\partial_\lambda \rangle$ where m' and n' are quadratic in λ .

To obtain a 2-plane congruence in the form (10), we let $X = V_0 + \lambda V_1 + (a - \lambda)(V_1 + \lambda V_2)$ and $Y = -(V_1 + \lambda V_2)$ so that \hat{X} and \hat{Y} are given by (9) with $m = m' + (a - \lambda)n'$ and $n = -n'$. The Lax integrability condition $[\hat{X}, \hat{Y}] = 0$ implies n' is affine linear in λ , while m' is a quadratic in λ , where the coefficient h of λ^2 is a function of x and y . We may set h to zero using the coordinate freedom

$$x \mapsto x, \quad y \mapsto \rho(x, y), \quad t \mapsto \rho_y(x, y)^2 t, \quad \lambda \mapsto \rho_y(x, y)(\lambda - 2\rho_{yy}(x, y)t),$$

which preserves the form of $\theta(\lambda)$ (hence also g) up to rescaling by $\rho_y(x, y)^2$ and a redefinition of a and b . The Lax equation now implies that the λ coefficient of m' differs from $-a_y$ by a function of x and y which may be set to zero using the remaining coordinate freedom

$$x \mapsto x, \quad y \mapsto y, \quad t \mapsto t + \tau(x, y), \quad \lambda \mapsto \lambda - \tau_y(x, y).$$

We then find that $m' = -a_y\lambda - b_y$, $n' = a_t\lambda + b_t$, and hence

$$\begin{aligned} \hat{X} &= \partial_x + (-\lambda^2 + a\lambda + b)\partial_t - ((a_y\lambda + b_y) + (\lambda - a)(a_t\lambda + b_t))\partial_\lambda, \\ \hat{Y} &= \partial_y - \lambda\partial_t - (a_t\lambda + b_t)\partial_\lambda. \end{aligned}$$

The Lax integrability condition now reduces to the determined system

$$(a_x - aa_y + ba_t)_t = (a_y - 2aa_t)_y, \quad (b_x - ab_y + bb_t)_t = (b_y - 2ab_t)_y. \quad (31)$$

This is the form of the EW system given in [12, (11)–(12)], except that the x and t variables have been swapped in our conventions and we have used the identity $(aa_y)_t =$

$(aa_t)_y$. Substituting $a = v_t$ and $b = u - v_y$ gives the Manakov–Santini system. The modified version (31) may also be written more geometrically as

$$\Delta^g a = 0, \quad \Delta^g b + \frac{3}{2}\{a, b\}_P = 0,$$

where Δ^g is the Laplacian of the metric g in (28), and $\{a, f\}_P = a_y f_t - a_t f_y$ is the Poisson bracket with respect to the bivector field $P = \partial_y \wedge \partial_t$ tangent to the null surface foliation.

Remark 3. In [12] a translationally noninvariant version of the MS system was also derived and the question of an explicit equivalence to the standard MS system was raised. However, the translationally noninvariant version is obtained from a generic null surface foliation of the EW manifold, and the coordinate transformation to a totally geodesic null surface foliation will be transcendental in general.

4.3. Arbitrary lifts of 2-plane congruences in 3D. We showed in Proposition 1 that any dLp can be made normal. However, when $d = 3$, the normal lift of a 2-plane congruence Π is not unique. Instead, the rank 3 distribution $\Delta = \hat{\pi}_*^{-1}(\Pi) \subseteq T\hat{M}_u$ has a unique Cauchy characteristic: a rank 1 subbundle $\mathcal{C} \subseteq \Delta$ with $[\mathcal{C}, \Delta] = \Delta$. For a rank 2 subbundle $\hat{\Pi} \subseteq \Delta$ the normality condition $[\hat{\Pi}, \hat{\Pi}] \subseteq \Delta$ implies that $\mathcal{C} \subseteq \hat{\Pi}$, but one generator of $\hat{\Pi}$ remains undetermined. In the case of interest that $\Pi = \ker \theta$ is characteristic for a quadric, an easy computation shows that \mathcal{C} is spanned by the vector field

$$\hat{W} = W_0 + \sigma \partial_\lambda, \quad W_0 := V_0 + 2\lambda V_1 + \lambda^2 V_2,$$

where, using the structure functions $c_{ij}^k = \theta_k([V_i, V_j])$ of the adapted frame V_0, V_1, V_2 , we have

$$\sigma = -c_{01}^2 + \lambda(c_{01}^1 - c_{02}^2) - \lambda^2(c_{12}^2 - c_{02}^1 + c_{01}^0) + \lambda^3(c_{12}^1 - c_{02}^0) - \lambda^4 c_{12}^0. \quad (32)$$

These formulae are compatible with representation $\sigma = m_0 + 2\lambda m_1 + \lambda^2 m_2$ for the coefficients m_i of the lifts of V_i which we want to show can be chosen quadratic in λ .

Without loss of generality we may write $\hat{\Pi} = \langle \hat{W}, \hat{U} \rangle$ with

$$\hat{U} = W_1 + \psi \partial_\lambda, \quad W_1 = \frac{1}{2}(W_0)_\lambda = V_1 + \lambda V_2. \quad (33)$$

We also write $W_2 = V_2 = (W_1)_\lambda$. The nondegeneracy condition (15) on Π implies that W_0, W_1, W_2 form a (λ -dependent) frame for TM_u and indeed

$$W_0 \odot W_2 - W_1^2 = V_0 \odot V_2 - V_1^2$$

is the inverse metric to $g = 4\theta_0\theta_2 - \theta_1^2$, which is nondegenerate and independent of λ .

The Frobenius integrability condition $[\hat{\Pi}, \hat{\Pi}] = \hat{\Pi}$ is the condition that

$$[\hat{W}, \hat{U}] = [V_0, V_1] + \lambda[V_0, V_2] + \lambda^2[V_1, V_2] + \sigma V_2 - 2\psi(V_1 + \lambda V_2) + (\hat{W}(\psi) - \hat{U}(\sigma))\partial_\lambda$$

is a section of $\hat{\Pi}$. Identifying $V_1 \equiv -\lambda V_2 - \psi \partial_\lambda$, $V_0 \equiv \lambda^2 V_2 + (2\lambda\psi - \sigma)\partial_\lambda$ modulo $\hat{\Pi}$, and assuming that the lift is normal, this reduces to $\epsilon = 0$, where

$$\begin{aligned} \epsilon &:= (W_0 + q_1 + \sigma \partial_\lambda)\psi + 2\psi^2 - \hat{q}_0, \quad \hat{q}_0 = W_1 \sigma + q_0 \sigma, \\ q_1 &= c_{02}^2 - 2c_{01}^1 + \lambda(2c_{12}^2 - 3c_{02}^1 + 4c_{01}^0) - \lambda^2(4c_{12}^1 - 5c_{02}^0) + 6\lambda^3 c_{12}^0, \\ q_0 &= c_{01}^0 + \lambda c_{02}^0 + \lambda^2 c_{12}^0. \end{aligned} \quad (34)$$

Using the coefficients of the decomposition $[W_0, W_1] = \bar{c}_{01}^0 W_0 + \bar{c}_{01}^1 W_1 + \bar{c}_{01}^2 W_2$ we get

$$\sigma = -\bar{c}_{01}^2, \quad q_1 = -\bar{c}_{01}^1 - \sigma_\lambda, \quad q_0 = \bar{c}_{01}^0. \quad (35)$$

Note that $\deg_\lambda \bar{c}_{01}^0 = 2$, $\deg_\lambda \bar{c}_{01}^1 = 3$ and $\deg_\lambda \bar{c}_{01}^2 = 4$.

Example 5. (dKP) For the dKP equation (2), we have $g = dy^2 - 4dx dt + 4u dx^2$, $\theta = dt + \lambda dy + (\lambda^2 - u) dx$ and

$$\hat{W} = \partial_x - 2\lambda\partial_y + (\lambda^2 + u)\partial_t + (\lambda u_t - u_y)\partial_\lambda, \quad \hat{U} = -\partial_y + \lambda\partial_t + \psi\partial_\lambda,$$

whence $\epsilon = \psi_x - 2\lambda\psi_y + (\lambda^2 + u)\psi_t + (\lambda u_t - u_y)\psi_\lambda - u_{yy} + 2\lambda u_{yt} - \lambda^2 u_{tt} - \psi u_t + 2\psi^2$. In this case, via the change of variables $\psi = \varphi^{-1} + u_t$, the equation $\epsilon = 0$ (34) takes the linear inhomogeneous form

$$\begin{aligned} \mathcal{L}_-(\varphi) = 2 &\Leftrightarrow \mathcal{L}_+(\varphi^{-1}) = -2\varphi^{-2}, \\ \text{where } \mathcal{L}_\pm &= \partial_x - 2\lambda\partial_y + (\lambda^2 + u)\partial_t + (\lambda u_t - u_y)\partial_\lambda \pm 3u_t. \end{aligned} \quad (36)$$

If we assume ψ either local (= differential) in u or global (= algebraic) in λ , then the only solution is $\varphi^{-1} = 0$, implying the existence of a unique dLp (standard) of these types.

However, there exist solutions to (36) which are non-algebraic in λ and nonlocal in u . Indeed for any Cauchy data $u|_{t=0}$ that is non-algebraic in λ , we obtain such a solution. In this way we obtain a (characteristic but not projective or local) Lax pair that does not give rise to an EW structure. Moreover, there is no uniqueness for such Lax pairs.

In the following Sects. 4.4–4.6 we deduce the cubic behaviour of ψ in λ from the equation $\epsilon = 0$, a strengthened nondegeneracy condition, and the requirement that the dLp is local in u . This suffices to establish the projective property, and hence Theorem 2.

4.4. Scalar PDEs in 3D. We first consider the case of a scalar differential equation $\mathcal{E} : F = 0$ of order ℓ , i.e., one PDE (20) on one function u . As before, we assume that the characteristic variety $\chi^\mathcal{E}$ is a quadric, which implies that ℓ is even and the symbol $F_{(\ell)}$ of the differential operator is a power of a nondegenerate quadratic form: $\ell = 2m$, $F_{(\ell)} = Q^m$ for some $Q \in \Gamma(S^2 T M_u)$ on \mathcal{E} . (For a second order scalar PDE (1) we get $m = 1$.) Using the notation of the previous section, we have $Q = W_0 \odot W_2 - W_1^2$.

The order of the conformal structure c_F in u satisfies $k = \text{ord}(c_F) \leq \ell$, and the strict inequality is possible, for instance, when F is quasilinear (dKP is an example with $0 = k < \ell = 2$). Then the frame V_i with the dual coframe θ_i can be chosen of the same order k in u , while the structure functions c_{ij}^t and the coefficient σ in (32) have order $\leq k + 1$.

Let us suppose $\hat{\Pi}$ is a normal dLp for \mathcal{E} . We want to find an \mathcal{E} -equivalent dLp which is projective. Since $\hat{\Pi}$ is normal, we may suppose, as in the previous section, that its integrability condition is $\epsilon = 0$ with ϵ given by (34). Hence by definition of a dLp, $\epsilon = \square F$ for some operator \square in total derivatives. If ψ has order $r \geq k + 2$ then by taking the $(r + 1)$ -symbol of this equation we obtain

$$W_0 \odot \psi_{(r)} = \epsilon_{(r+1)} = \square_{(r+1-\ell)} \odot F_{(\ell)} = \square_{(r+1-2m)} \odot (W_0 \odot W_2 - W_1^2)^m,$$

and hence conclude (since $W_0 \odot W_2 - W_1^2$ is indivisible by W_0) that the symbol of ψ is divisible by that of F . Therefore we can modify ψ off shell (fixed on shell) to obtain an \mathcal{E} -equivalent dLp in which the new ψ has order $< r$. By iterating this process, we may thus assume, up to \mathcal{E} -equivalence, that ψ has order $\leq k + 1$ from the outset.

The $(k + 2)$ -symbol of $\epsilon = \square F$ now yields, using equation (34), the relation

$$W_0 \odot \psi_{(k+1)} - W_1 \odot \sigma_{(k+1)} = R \odot (W_0 \odot W_2 - W_1^2)^m \quad (37)$$

for a section $R \in \Gamma(S^{k-2m+2} T M_u)$ of the bundle of homogeneous degree $k - 2m + 2$ polynomials on $T^* M_u$, i.e., $R = \sum_{|\tau|=k-2m+2} a_\tau W_\tau$, where we let $W_\tau = W_{j_1} \odot \dots \odot W_{j_t}$ for a multi-index $\tau = (j_1 \dots j_t)$ of length $|\tau| = t$. By modification of ψ and σ off shell, we can bring this function modulo \mathcal{I}_F to the form

$$R = (-1)^{m-1} \mu W_2^{k-2m+2} \quad (38)$$

(indeed, all terms of R with W_0 factor can be absorbed into $\psi_{(k+1)}$, while those with W_1 factor into $\sigma_{(k+1)}$; this absorption is identical on shell). Formula (37) then implies that

$$\sigma_{(k+1)} = W_1 \odot R \odot Q^{m-1} + W_0 \odot T, \quad \psi_{(k+1)} = W_2 \odot R \odot Q^{m-1} + W_1 \odot T \quad (39)$$

for some $T \in \Gamma(S^k T M_u)$, and by the normalization (38), the coefficients for R and T are uniquely determined by independent components of σ and hence they are polynomial in λ . In particular, since σ is a quartic polynomial in λ , we conclude that $\mu \in C^\infty(J^{k+1} M_u)$ is a polynomial in λ with $\deg_\lambda \mu \leq 3$.

Also, T is a polynomial in λ with $\deg_\lambda T \leq 2$. Therefore, $\psi_{(k+1)}$ is a cubic polynomial in λ . Thus, there exists a function $\psi_1 = \psi_1(\partial^{k+1} u, \lambda)$ with $\deg_\lambda \psi_1 \leq 3$ such that $\psi_0 := \psi - \psi_1$ has order $\leq k$ in u . Substituting $\psi = \psi_1 + \psi_0$ into the equation $\epsilon = \square F$ we get

$$(W_0 + \tilde{q}_1 + \sigma \partial_\lambda) \psi_0 + 2\psi_0^2 = \tilde{q}_0, \quad (40)$$

where $\tilde{q}_1 = q_1 + 4\psi_1$, while \tilde{q}_0 is a expression of order $k + 1$ in u that we do not write explicitly. However, it follows from (35) that \tilde{q}_1, \tilde{q}_0 are polynomial in λ with $\deg_\lambda \tilde{q}_1 \leq 3$, $\deg_\lambda \tilde{q}_0 \leq 8$. We now want to show that ψ_0 is also polynomial in λ .

In order to do this, it is convenient to carry out computations in the nonholonomic λ -dependent frame $(W_i)_{i=0}^2$ rather than the holonomic frame $(\partial_{x^i})_{i=0}^2$ induced by local coordinates $(x^i)_{i=0}^2$ on M_u . Let (a_i^j) be the transition matrix between these frames and (b_i^j) be its inverse, i.e., $W_i = a_i^j \partial_{x^j}$ and $\partial_{x^i} = b_i^j W_j$ (summation convention). These vector fields induce vertical vector fields \mathbb{D}_{W_τ} on jets via the formulae $\mathbb{D}_{W_\tau} = b_{j_1}^{i_1} \dots b_{j_t}^{i_t} \partial_{u_{i_1 \dots i_t}}$, where $\tau = (j_1, \dots, j_t)$. If ξ is a function on the jet bundle (i.e., a differential operator) with order t symbol $\xi_{(t)}$, then $\mathbb{D}_{W_\tau} \xi$, with $|\tau| = t$, is the coefficient ξ_τ^t of $\xi_{(t)}$ in the decomposition $\xi_{(t)} = \xi_\tau^t W_\rho$ (summation over multi-indices ρ with $|\rho| = t$).

Next, since the dual co-frame to (W_0, W_1, W_2) is $(\frac{1}{2}\theta_{\lambda\lambda}, -\theta_\lambda, \theta)$, the coefficients on the right hand sides of identities (35) may be written

$$\tilde{c}_{01}^2 = \theta([W_0, W_1]), \quad \tilde{c}_{01}^1 = -\theta_\lambda([W_0, W_1]), \quad \tilde{c}_{01}^0 = \frac{1}{2}\theta_{\lambda\lambda}([W_0, W_1]).$$

This and formula (39) lead to

$$\mu = \mathbb{D}_{W_1}^{2m-1} W_2^{k-2m+2}(\sigma) = \mathbb{D}_{W_1}^{2m-2} W_2^{k-2m+3}(\psi) = \theta(\mathbb{D}_{W_1}^{2m-2} W_2^{k-2m+2}(W_0)). \quad (41)$$

To obtain the last formula note that we have $\mathbb{D}_{W_1^a W_2^b}[W_0, W_1] = -\mathbb{D}_{W_1^{a-1} W_2^b} W_0$ from the definition of the commutator. Then apply this to the identity $\sigma = -\theta([W_0, W_1])$ and the first formula for μ .

Identity (39) yields $\theta(\mathbb{D}_{W_1^r W_2^{k-r}} W_0) = 0$ unless $r = 2m - 2$. Note that by (39) we have $\sigma_{(k+1)} = \mu W_1^{2m-1} \odot W_2^{k-2m+2} \bmod W_0 \odot S^k T M_u$, and this decomposition can be refined:

$$\begin{aligned}\sigma_{(k+1)} &= \mu W_1^{2m-1} \odot W_2^{k-2m+2} + \gamma W_0 \odot W_1^{2m-2} \odot W_2^{k-2m+2} + \dots \\ \psi_{(k+1)} &= \mu W_1^{2m-2} \odot W_2^{k-2m+3} + \gamma W_1^{2m-1} \odot W_2^{k-2m+2} + \dots,\end{aligned}$$

where $\gamma = \mathbb{D}_{W_1^{2m-2} W_2^{k-2m+2}}(T) = \theta(\mathbb{D}_{W_0 W_1^{2m-3} W_2^{k-2m+2}}(W_0)) - \theta(\mathbb{D}_{W_1^{2m-2} W_2^{k-2m+2}}(W_1))$ and by dots we mean all terms with other W_τ that are irrelevant for the computation. Consequently,

$$(\sigma_\lambda)_{(k+1)} = (2m - 1)\mu W_1^{2m-2} \odot W_2^{k-2m+3} + (\mu_\lambda + 2\gamma) W_1^{2m-1} \odot W_2^{k-2m+2} + \dots$$

and since $\sigma_\lambda = (\theta[W_1, W_0])_\lambda = \theta_\lambda[W_1, W_0] + \theta[W_2, W_0] = \bar{c}_{01}^1 - \theta[W_0, W_2]$ we get

$$(\bar{c}_{01}^1)_{(k+1)} = (\sigma_\lambda)_{(k+1)} - \mu W_1^{2m-2} \odot W_2^{k-2m+3} \bmod W_0 \odot S^k T M_u.$$

Thus from (35) we get the following expression for the $(k + 1)$ -symbol

$$\begin{aligned}(\tilde{q}_1)_{(k+1)} &= -2(\sigma_\lambda)_{(k+1)} + 5\mu W_1^{2m-2} \odot W_2^{k-2m+3} + 4\gamma W_1^{2m-1} \odot W_2^{k-2m+2} + \dots \\ &= (7 - 4m)\mu W_1^{2m-2} \odot W_2^{k-2m+3} - 2\mu_\lambda W_1^{2m-1} \odot W_2^{k-2m+2} + \dots\end{aligned}$$

Taking now $(k + 1)$ -symbol of (40), we get

$$W_0 \odot (\psi_0)_{(k)} + (\tilde{q}_1)_{(k+1)} \psi_0 + \sigma_{(k+1)} (\psi_0)_\lambda = (\tilde{q}_0)_{(k+1)}.$$

Denoting $(\tilde{q}_0)_{(k+1)} = \kappa_0 W_1^{2m-2} \odot W_2^{k-2m+3} + \kappa_1 W_1^{2m-1} \odot W_2^{k-2m+2} + \dots$ and extracting the coefficients at the indicated terms (which are unchanged by \mathcal{E} -equivalence) we obtain the following system

$$(7 - 4m)\mu\psi_0 = \kappa_0, \quad \mu\partial_\lambda\psi_0 - 2\mu_\lambda\psi_0 = \kappa_1.$$

All coefficients of this linear system on ψ_0 are polynomials in λ . We assume $\mu \neq 0$ (this condition will be discussed in the next section). Then the first equation uniquely determines ψ_0 . Moreover, μ divides κ_0 as a polynomial in λ because otherwise ψ_0 is a proper rational function and then the second equation, written as $(\psi_0 \mu^{-2})_\lambda = \kappa_1 \mu^{-3}$, yields a contradiction.

Thus ψ_0 , and hence also ψ , are polynomials in λ , and $\deg_\lambda \psi \leq 5$. Since the parameter λ is manifestly projective, a projective change should not destroy the polynomial property. Using the special projective transformation $\lambda \mapsto \lambda^{-1}$ (or a similar projective transformation arbitrarily close to the identity), we conclude that in fact $\deg_\lambda \psi \leq 3$.

Moreover, the function ψ should be compatible with σ in the sense that $\psi = m_1 + \lambda m_2$, $\sigma = m_0 + 2\lambda m_1 + \lambda^2 m_2$ for some λ -quadratics m_i . This will be derived from the cubic property of ψ in the general case in Sect. 4.6.

4.5. *Nondegeneracy for scalar and vector equations.* The λ -dependent quantity μ introduced in (41) characterizes the extent to which the Lax integrability condition depends on the equation \mathcal{E} . If this condition does not involve F and its derivatives, the dLp is trivial (holds off shell). We require that the equation enters on the level of the top symbol, i.e., $(k+1)$ -jet, or equivalently that $\mu \neq 0$. We first observe that this condition is invariant under admissible transformations of \hat{M}_u as a \mathbb{P}^1 -bundle over M_u .

Proposition 4. *The scalar quantity μ is a relative differential invariant, i.e., transforms by a nonvanishing scalar multiple under admissible transformations.*

Proof. The admissible transformations of \hat{M}_u have the form $(\mathbf{x}, \lambda) \mapsto (\Phi(\mathbf{x}), \Psi(\mathbf{x}, \lambda))$, where Φ is a conformal transformation of $(M_u, [g])$ and $\Psi(\mathbf{x}, \lambda) = \frac{a(\mathbf{x})+b(\mathbf{x})\lambda}{c(\mathbf{x})+d(\mathbf{x})\lambda}$ is a parametric Möbius transformation. These preserve the algebraic behaviour of the dLp $\hat{\Pi}$, and a straightforward computation shows they scale μ by a nonvanishing scalar multiple.

Alternatively, using the framework and normalizations of Sect. 4.4, σ given by (32) is independent of the adapted frame up to scale and the leading coefficient of its symbol $\mathbb{D}_{W_1^{2m-1}W_2^{k-2m+2}}(\sigma)$ is a relative invariant, as required. \square

Let us now give the vector version, recalling the set-up. In this case $F : J^\ell(M, \mathcal{V}) \rightarrow \mathcal{W}$ is a determined (nonlinear) differential operator of order ℓ on sections \mathbf{u} of a fibre bundle \mathcal{V} over M_u with values in a rank s vector bundle \mathcal{W} . We assume, for simplicity, that \mathcal{V} is also a vector bundle of rank s so that we can identify the vertical bundle $T_u^v\mathcal{V}$ along a section \mathbf{u} with \mathcal{V} . Locally, in coordinates, F has components F^i that are scalar differential operators of order ℓ on vector-function $\mathbf{u} = (u^j)$ of \mathbf{x} , where $i, j \in \{1, \dots, s\}$.

The symbol of F at $\mathbf{u} = (u^i)$ is a map $F_{(\ell)} : S^\ell T^*M_u \otimes \mathcal{V} \rightarrow \mathcal{W}$ that we identify with an $s \times s$ matrix $\mathbf{F} = (F_i^j)$, whose coefficients are polynomials of degree ℓ on T^*M_u . Similarly, the symbol of a scalar differential operator φ can be identified with a column in components. The characteristic variety $\chi^\mathcal{E}$ of \mathcal{E} is a quadric if $\det(\mathbf{F}) = Q^m$ as before.

The setup of the previous section extends, and $\mu = \theta(\mathbb{D}_{W_1^{2m-2}W_2^{k-2m+2}}(W_0))$ is a section of the bundle \mathcal{V} over M_u for solutions u of the vector version of (20). The following statement is proved similarly to Proposition 4.

Proposition 5. *The section μ is a relative differential invariant, i.e., under admissible transformations it is mapped to another section related to μ by an automorphism of the bundle \mathcal{V} . Hence the (non-)vanishing of μ is an invariant property.*

Note that μ depends not on a lift or dLp but only on the equation $\mathcal{E} : F = 0$ itself.

Definition 7. In 3D the equation is called nondegenerate (and its dLp nontrivial) if the relative invariant μ is nonzero (identically in λ).

This condition is trivially satisfied if the conformal structure c_F has zero order in u , as happens in the dKP case. It can be proved for several classes of PDE in 3D (with $\text{ord}_u(c_F) > 0$), and we do not know of any integrable equation violating this condition.

Remark 4. In fact, the Manakov–Santini equation, which by [12] is the master equation for EW geometry, is nondegenerate in the sense of this definition. We check this for the modified version, with W_0 given by (30). Since order of the conformal structure in this formalism is $k = 0$, and also $m = 1$, we compute the symbol of W_0 by (a, b) as (∂_t, ∂_y) and applying θ given by (29) we get $\mu = (\lambda, 1) \neq 0$. Thus the MS equation is nondegenerate and we adopt this condition for our main result.

4.6. *Proof of Theorem 2.* Any nondegenerate dLp $\hat{\Pi}$ is \mathcal{E} -equivalent to a normal dLp by Proposition 1. When $d = 4$, the normal dLp has the projective property, while for $d = 3$, the main task is to show that, up to \mathcal{E} -equivalence, we may assume that ψ in (33) is cubic in λ . The proof almost directly generalizes the scalar version of Sect. 4.4, so we only indicate important differences on each step.

- (i) We begin with equation $\epsilon = \square F$ and, as before, by an off shell modification can arrange that $\text{ord } \psi \leq k+1$, where k is the order of the conformal structure c_F . Then its $(k+2)$ -symbol and (34) yield the following matrix equation

$$\begin{bmatrix} \psi^1 & \sigma^1 \\ \vdots & \vdots \\ \psi^s & \sigma^s \end{bmatrix} \odot \begin{bmatrix} W_0 \\ -W_1 \end{bmatrix} = \begin{bmatrix} F_1^1 & \cdots & F_s^1 \\ \vdots & \ddots & \vdots \\ F_1^s & \cdots & F_s^s \end{bmatrix} \odot \begin{bmatrix} R^1 \\ \cdots \\ R^s \end{bmatrix} \quad (42)$$

where ψ^i are components of the symbol $\psi_{(k+1)}$ and similarly for σ , and where R^i are the symbols of some operators in total derivatives. Multiplying this equation by the adjugate matrix $\text{adj}(\mathbf{F})$ (which satisfies $\text{adj}(\mathbf{F})\mathbf{F} = \det(\mathbf{F})I = Q^m I$) and denoting the rows of the resulting left-hand side matrix $[\tilde{\psi}^i \tilde{\sigma}^i]$ we get the equations

$$W_0 \odot \tilde{\psi}^i - W_1 \odot \tilde{\sigma}^i = R_i \odot (W_0 \odot W_2 - W_1^2)^m, \quad i \in \{1, \dots, s\},$$

from which we obtain a vector analogue of equation (39) for each component $i \in \{1, \dots, s\}$. Moreover, we can obtain normalization analogues $R_i = (-1)^{m-1} \mu_i W_2^{k-2m+2}$ of (38). This implies that $\tilde{\psi}^i$ and hence ψ^i can be chosen polynomial in λ , moreover $\deg_\lambda \psi^i \leq 3$.

- (ii) Thus there exist a decomposition $\psi = \psi_1 + \psi_0$, where ψ_1 is at most cubic in λ and has $(k+1)$ -symbol (ψ^i) at $\mathbf{u} = (u^j)$, while ψ_0 has order $\leq k$. Substituting this into the constraint $\epsilon = \square F$ we obtain a vector analogue of equation (40). Taking its $(k+1)$ -symbol and applying $\text{adj}(\mathbf{F})$ again gives

$$\tilde{q}_1^i \psi_0 + \tilde{\sigma}^i \partial_\lambda \psi_0 = \tilde{q}_0^i, \quad i \in \{1, \dots, s\}.$$

If $\mu = (\mu_1, \dots, \mu_s)$ is nonzero (identically in λ), we conclude by the same argument that ψ_0 is polynomial in λ with $\deg \psi_0 \leq 5$. Hence ψ is a polynomial in λ with $\deg \psi \leq 5$.

- (iii) If the coefficients of ψ at λ^4 or λ^5 are nonzero, then a Möbius transformation $\lambda \mapsto \frac{a(x)+b(x)\lambda}{c(x)+d(x)\lambda}$ arbitrary close to the identity maps the system of vector fields \hat{W} and $\hat{U} = V_1 + \lambda V_2 + \psi \partial_\lambda$ (after taking a proper linear combination and clearing denominators) to a system of the same form with a new ψ of higher degree. Thus we must have $\deg_\lambda \psi \leq 3$, which is a projectively invariant property.
- (iv) In addition to \hat{U} the vector field $\hat{W} - \lambda \hat{U} = V_0 + \lambda V_1 + (\sigma - \lambda \psi) \partial_\lambda$ have degree ≤ 3 in λ . Indeed, under a change of the adapted frame (V_0, V_1, V_2) and a projective change of parameter λ this field becomes of the form \hat{U} and so the claim follows from (i)–(iii).
- (v) Note that the frame V_0, V_1, V_2 is arbitrary subject to the condition that $V_0 \odot V_2 - V_1^2$ is conformal to the inverse metric. In particular, $V_1 + \lambda V_2 + \psi \partial_\lambda$ has cubic coefficient of ∂_λ for arbitrary non-null V_1 and null V_2 . Changing the pair (V_1, V_2) to the pair $(-V_1, V_2)$ and subtracting the results we get that the lift of V_1 has cubic coefficient of ∂_λ . Here V_1 is arbitrary non-null vector. Applying a similar argument to $V_0 +$

$\lambda V_1 + (\sigma - \lambda\psi)\partial_\lambda$ from (iv) we obtain that the coefficient of ∂_λ in the lift of arbitrary null vector V_0 is also at most cubic in λ .

Now the property of being cubic in λ for the coefficient of ∂_λ of the lift of a λ -independent vector is not projectively invariant, while the entire construction is projectively invariant. Therefore the coefficient of ∂_λ in the lift of arbitrary vector is actually quadratic.

Theorem 2 is now immediate. By Lemma 5 normal lifts with this projective property are bijective with Weyl connections for $d = 3$, while for $d = 4$ the normal lift is unique by Lemma 1. Thus for $d = 3$ or $d = 4$, the standard Lax pair of (M_u, c_F, ∇) or (M_u, c_F) is \mathcal{E} -equivalent to $\hat{\Pi}$. \square

5. Applications and Generalizations

5.1. Pseudopotentials. In this paper we have defined dispersionless integrable systems using a Lax pair of vector fields. In 3D, an alternative approach relies instead on pseudopotentials or nonlinear dispersionless Lax pairs, cf. [17,27,38].

Definition 8. A *pseudopotential* for a PDE $F = 0$ is a function $S: M_u \rightarrow \mathbb{R}$ whose derivative dS satisfies an overdetermined system of two equations that are compatible on shell, i.e., when $F(j^\ell u) = 0$.

Locally, in coordinates (x, y, t) , we may write these equations as $S_x = A(S_t)$ and $S_y = B(S_t)$ where A and B also depend on (x, y, t) . If they depend on (x, y, t) (only or also) through a section v of a vector bundle over M_u , and the integrability condition $\partial_y(A(S_t)) = \partial_x(B(S_t))$ is required to hold identically in S_t , we obtain a PDE system on v . Dispersionless integrable systems are often defined as those determined PDEs arising in this way.

More invariantly, the two equations determine a codimension two (hence 4-dimensional) submanifold N of the cotangent bundle T^*M_u and S is a pseudopotential with respect to these equations if dS takes values in N . The integrability condition means that N is coisotropic for the canonical symplectic form Ω on T^*M_u . Here we recall that $\Omega = d\tau$ where τ is the tautological 1-form on T^*M_u (with $\beta^*\tau = \beta$ for any 1-form β on M_u). The coisotropic condition means that the pullback of Ω to N has rank two, hence a 2-dimensional radical (or kernel).

Locally N is a fibre bundle over M_u and we may take $\lambda = S_t$ as a fibre coordinate in the above explicit formulation. Thus

$$\begin{aligned}\partial_y(A(S_t)) &= A_y + A_\lambda S_{tx} = A_y + A_\lambda B_t + A_\lambda B_\lambda S_{tt}, \\ \partial_x(B(S_t)) &= B_x + A_\lambda S_{ty} = B_x + A_t B_\lambda + A_\lambda B_\lambda S_{tt},\end{aligned}$$

and so the integrability condition is

$$A_y - B_x = \{A, B\}_P := A_t B_\lambda - A_\lambda B_t, \quad (43)$$

where $\{A, B\}_P$ is the Poisson bracket of A and B with respect to the Poisson structure $P := \partial_t \wedge \partial_\lambda$; equivalently, the vector fields $\partial_x + P(dA)$ and $\partial_y + P(dB)$ commute, where $P(dA) = A_t \partial_\lambda - A_\lambda \partial_t$ and $P(dB) = B_t \partial_\lambda - B_\lambda \partial_t$ are the hamiltonian vector fields associated to A and B by the Poisson structure P (cf. e.g. [13]).

Alternatively, if (43) holds, then the pullback of Ω to N is

$$\begin{aligned} d\lambda \wedge dt + (A_\lambda d\lambda + A_t dt) \wedge dx + (B_\lambda d\lambda + B_t dt) \wedge dy - (A_t B_\lambda - A_\lambda B_t) dx \wedge dy \\ = (d\lambda - A_t dx - B_t dy) \wedge (dt + A_\lambda dx + B_\lambda dy) \end{aligned}$$

and its radical is the dLp spanned by $\partial_x + P(dA)$ and $\partial_y + P(dB)$.

Conversely, let $\hat{\Pi}$ be a dLp on $\hat{\pi}: \hat{M}_u \rightarrow M_u$. On shell, $\hat{\Pi}$ is integrable and so \hat{M} fibres locally over a minitwistor space $\mathcal{T}w$ [19]. At least locally $\mathcal{T}w$ admits a nondegenerate (and necessarily closed) 2-form (such as $dz_1 \wedge dz_2$ in local coordinates); this then pulls back to a closed 2-form ω on \hat{M}_u with radical $\hat{\Pi}$. We may therefore write (locally, on shell) $\omega = d\alpha$ for a 1-form α on \hat{M}_u , which we may assume vanishes on the fibres of \hat{M}_u over M_u ; hence we may write $\alpha = (Id, \tilde{\alpha}) \circ \hat{\pi}_*$ for a section $(Id, \tilde{\alpha})$ of $\hat{\pi}^* T^* M_u = \{(\hat{p}, \xi) \in \hat{M}_u \times T^* M_u \mid \xi \in T_{\hat{\pi}(\hat{p})}^* M_u\}$. Then $\tilde{\alpha}: \hat{M}_u \rightarrow T^* M_u$ is an immersion whose image is coisotropic, since $\tilde{\alpha}^* \tau = \alpha$ and so $\tilde{\alpha}^* \Omega = d\alpha = \omega$ has rank two with radical $\hat{\Pi}$.

In order to do this off shell, we have to work modulo the PDE system. However, the construction of the coisotropic immersion $\tilde{\alpha}$ from a dLp requires integration, and so it may be necessary to pass to a covering system.

Example 6. (dKP) We illustrate this with the well-known example of the dKP equation (2) $(u_x + uu_t)_t = u_{yy}$ with dLp (3). We must now find a function f so that $\omega = f\eta \wedge \theta$ is closed modulo the equation, and then a 1-form α such that $d\alpha = \omega$ modulo the equation. For the first step, it happens in this case that $f = 1$ works. For the second, setting

$$\alpha = \left(\frac{1}{3}\lambda^3 - u\lambda - v\right)dx + \left(\frac{1}{2}\lambda^2 - u\right)dy + \lambda dt,$$

we have that $d\alpha = \eta \wedge \theta$ modulo the covering system $v_t = u_y$ and $v_y = u_x + uu_t$. Thus the pseudopotential system is $S_x = \frac{1}{3}S_t^3 - uS_t - v$ and $S_y = \frac{1}{2}S_t^2 - u$.

Note that the above nonlocality (usage of v) may be avoided by using the potential form $u_{xt} + u_t u_{tt} - u_{yy} = 0$ of dKP. In this case the pseudopotential S is given by the equations: $S_x = \lambda^3/3 - u_t \lambda - u_y$, $S_y = \lambda^2/2 - u_t$, $S_t = \lambda$. In both cases the parameter λ is aligned to the Lax pair in the sense that it is the projective parameter on the correspondence bundle $\hat{M}_u \rightarrow M_u$. This is no longer so with Manakov–Santini system (6).

Example 7. (MS) The MS system does admit a pseudopotential formulation; however it is neither local in u, v nor rational in λ . The system

$$\sigma(R_x - P_t) = (u_t \lambda + u_y)(Q_x - P_y), \quad \sigma(R_y - Q_t) = u_t(Q_x - P_y), \quad (44)$$

$$\sigma P_\lambda = (\lambda^2 + v_t \lambda - u + v_y)(Q_x - P_y), \quad \sigma Q_\lambda = (\lambda + v_t)(Q_x - P_y),$$

$$\sigma R_\lambda = (Q_x - P_y) \quad (45)$$


with $\sigma = u_y \lambda + uu_t - u_t v_y + u_y v_t$, is a differential covering, meaning it is compatible modulo MS. Here the last three equations (45) determine the behaviour in the spectral parameter λ , while the first two equations (44) yield a pseudopotential S via the system $S_x = P$, $S_y = Q$, $S_t = R$. Indeed, one can verify that the differential $\omega = d\alpha$ of the 1-form $\alpha = Pdx + Qdy + Rdt$ on \hat{M}_u satisfies $\hat{X} \lrcorner \omega = \hat{Y} \lrcorner \omega = 0$ modulo MS and (44)–(45), where $\hat{X} = \tilde{X}|_{\tilde{\lambda}=\lambda}$, $\hat{Y} = \tilde{Y}|_{\tilde{\lambda}=\lambda}$ in terms of formula (8) are vector fields on \hat{M}_u forming the Lax pair (with parameter λ projective).

5.2. *Twistor interpretation via contact coverings.* To relate the pseudopotential formulation more closely to the dLp formulation, we focus on the first order quasilinear system for sections of $\hat{\pi} : \hat{M}_u \rightarrow M_u$ which correspond to hypersurfaces in the twistor space. We refer to this PDE system as a *contact covering* of \mathcal{E} because the equation it defines is a codimension 2 submanifold \mathcal{Q} of $J^1\hat{\pi}$ (the bundle of 1-jets of sections of $\hat{\pi}$), which is a contact manifold.

This viewpoint gives an alternative way to understand why contact coverings are equivalent to dLps. For this, let α be a contact form on $J^1\hat{\pi}$ representing the standard contact structure and let $\alpha_{\mathcal{Q}}$ be its restriction on \mathcal{Q} . Then for $\omega_{\mathcal{Q}} = d\alpha_{\mathcal{Q}}$ we have: $\alpha_{\mathcal{Q}} \wedge \omega_{\mathcal{Q}}^{d-1} = 0$ on shell, but $\alpha_{\mathcal{Q}} \wedge \omega_{\mathcal{Q}}^{d-2} \neq 0$, which implies that $\alpha_{\mathcal{Q}}$ has a 2-dimensional radical $\hat{\Pi} \subseteq T\mathcal{Q}$: $\xi \lrcorner \alpha_{\mathcal{Q}} = 0$ and $\xi \lrcorner \omega_{\mathcal{Q}} = 0$ for all $\xi \in \hat{\Pi}$.

If \mathcal{Q} is quasilinear, as we require in the definition of a contact covering, then $\hat{\Pi}$ is projectible along the fibres $\pi_{1,0} : \mathcal{Q} \rightarrow J^0\hat{\pi} = \hat{M}_u$ and so it induces a pushforward distribution of rank 2 in $T\hat{M}_u$, which is a dLp in our formalism. This is how a nonlinear covering induces a linear one, and the inverse relation is given by a lift.

We summarize the observed relations into the following diagram, intertwining the twistor and jet concepts:



Here the dotted arrow is the restriction of the jet-projection to the contact covering \mathcal{Q} , arrows labelled by $\hat{\Pi}$ are (local) quotients by the corresponding foliations, and $\mathcal{T}w^{d-1}$ is the mini-twistor or the twistor space for $d = 3$ or $d = 4$ respectively.

The dashed arrow is well-defined locally, when a local coordinate (spectral parameter) on the fibre \hat{M}_u is chosen, but it may fail to exist globally with respect to the spectral parameter λ and locally with respect to the dependent variable u . For $d = 3$, this is precisely the theory of pseudopotentials as discussed in Sect. 5.1. In this case, the space $\mathbb{P}(T^*\mathcal{T}w)$ is the real Penrose twistor space (projecting to the Hitchin mini-twistor space with fibre \mathbb{P}^1) that embeds into the complex twistor space $\mathcal{T}w_{\mathbb{C}}$, which is the complexification of $\mathcal{T}w$ of the case $d = 4$, via a conformal Killing reduction [20]. We leave to the reader a specification of relations between different real forms (signatures of the conformal structure—related by Wick rotations in physics language).

When $d = 4$ an analogue of the theory of pseudopotentials has been developed in [33]. Geometrically, this involves making a further projection to $\mathbb{P}(T^*M_u)$ on the left hand side of the above diagram (this is why the Lax pairs of [33] are homogeneous in $\partial\psi$ for the covering function ψ). This is a 7-dimensional contact manifold, so two equations suffice to define a 5-dimensional submanifold \hat{M}_u . In this formalism the Lax pair is given by contact hamiltonian vector fields.

5.3. Extensions of the theory. First, as noted in the introduction, in 2D, the theory of dispersionless Lax pairs is vacuous, essentially because there is only one 2-plane congruence. However, if we relax the assumption that the Lax pair is transverse to the fibres of \hat{M}_u over M_u , this objection evaporates. The characteristic condition means that at points of tangency, the projection of the Lax distribution is a characteristic direction. In particular, when the characteristic variety is a quadric (two points), we expect two points of tangency, with the background given by the spinor-vortex equations [1, 2].

Secondly, it would be useful to be able to relax the requirement that the PDE system $F: J^\ell(M, \mathcal{V}) \rightarrow \mathcal{W}$ determined in the sense that $\text{rank}(\mathcal{W}) = \text{rank}(\mathcal{V})$. The theory in this paper should at least extend to (formally) overdetermined systems ($\text{rank}(\mathcal{W}) \geq \text{rank}(\mathcal{V})$) which are compatible, so that the characteristic variety is a hypersurface. We would then need to use the compatibility conditions to generalize Theorem 1.

For truly overdetermined systems, with characteristic variety of higher codimension, it would be necessary also to replace Lax pairs by Lax distributions of higher rank. Recently the characteristic property was confirmed in [18] for paraconformal structures generalizing EW structures to higher dimension, and we suggest that it applies universally.

Finally, with the latter idea, the restriction to dimensions $d = 3, 4$ can be relaxed. This would extend the framework of integrability via geometry to a wider context.

Acknowledgements. We are grateful to Jenya Ferapontov for helpful discussions. In particular, he pointed out that dispersionless Lax pairs for PDEs of Hirota type are always characteristic, suggesting that this might be true more generally, and later drew our attention to difficulties in establishing the projective property for Lax pairs in 3D. We also thank the referee for detailed comments which greatly improved the clarity of the paper.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. Burtsev, S.P., Zakharov, V.E., Mikhailov, A.V.: The inverse scattering method with a variable spectral parameter. *Teoret. Mat. Fiz.* **70**, 323–341 (1987)
2. Calderbank, D.M.J.: Integrable background geometries. *SIGMA* **10**, 34 (2014)
3. Calogero, F., Nucci, M.C.: Lax pairs galore. *J. Math. Phys.* **32**, 72–74 (1991)
4. Cartan, E.: Les systèmes de Pfaff à cinq variables et les équations aux dérivées partielles du second ordre. *Ann. Sci. Ecole Norm. Sup.* **3**(27), 109–192 (1910)
5. Cartan, E.: Sur une classe d'espaces de Weyl. *Ann. Sci. Ecole Norm. Sup.* **3**(60), 1–16 (1943)
6. Doubrov, B., Ferapontov, E.V.: On the integrability of symplectic Monge-Ampère equations. *J. Geom. Phys.* **60**, 1604–1616 (2010)
7. Doubrov, B., Ferapontov, E.V., Kruglikov, B., Novikov, V.: On the integrability in Grassmann geometries: integrable systems associated with fourfolds $\text{Gr}(3, 5)$. *Proc. LMS* **116**, 1269–1300 (2018)
8. Doubrov, B., Ferapontov, E.V., Kruglikov, B., Novikov, V.: Integrable systems in 4D associated with sixfolds in $\text{Gr}(4, 6)$. *Int. Math. Res. Notices* **21**, 6585–6613 (2019)
9. Dubrovin, B.A., Novikov, S.P.: Hydrodynamics of weakly deformed soliton lattices: differential geometry and Hamiltonian theory. *Russian Math. Surv.* **44**(6), 35–124 (1989)
10. Dunajski, M.: A class of Einstein-Weyl spaces associated to an integrable system of hydrodynamic type. *J. Geom. Phys.* **51**(1), 126–137 (2004)

11. Dunajski, M.: An interpolating dispersionless integrable system. *J. Phys. A* **41**, 315202 (2008)
12. Dunajski, M., Ferapontov, E., Kruglikov, B.: On the Einstein-Weyl and conformal self-duality equations. *J. Math. Phys.* **56**, 083501 (2015)
13. Dunajski, M., Grant, J.D.E., Strachan, I.A.B.: Multidimensional integrable systems and deformations of Lie algebra homomorphisms. *J. Math. Phys.* **48**, 093502 (2007)
14. Dunajski, M., Mason, L.J., Tod, P.: Einstein-Weyl geometry, the dKP equation and twistor theory. *J. Geom. Phys.* **37**, 63–93 (2001)
15. Ferapontov, E.V., Huard, B., Zhang, A.: On the central quadric ansatz: integrable models and Painlevé reductions. *J. Phys. A* **45**, 195–204 (2012)
16. Ferapontov, E.V., Khusnutdinova, K.R.: Hydrodynamic reductions of multi-dimensional dispersionless PDEs: the test for integrability. *J. Math. Phys.* **45**(6), 2365–2377 (2004)
17. Ferapontov, E.V., Kruglikov, B.: Dispersionless integrable systems in 3D and Einstein-Weyl geometry. *J. Differ. Geom.* **97**, 215–254 (2014)
18. Ferapontov, E.V., Kruglikov, B.: Dispersionless integrable hierarchies and $GL(2, \mathbb{R})$ geometry. *Math. Proc. Cambridge Phil. Soc.* **1–26**, (2019). <https://doi.org/10.1017/S0305004119000355>
19. Hitchin, N.J.: Complex Manifolds and Einstein's Equations, Twistor Geometry and Nonlinear Systems (Primorsko, 1980), 73–99. *Lecture Notes in Math*, vol. 970. Springer, Berlin-New York (1982)
20. Jones, P.E., Tod, K.P.: Minitwistor spaces and Einstein-Weyl spaces. *Classical Quantum Gravity* **2**(4), 565–577 (1985)
21. Krasil'shchik, J., Verbovetsky, A.: Geometry of jet spaces and integrable systems. *J. Geom. Phys.* **61**, 1633–1674 (2011)
22. Kruglikov, B., Lychagin, V.: Geometry of differential equations, *Handbook of Global Analysis*, Ed. D.Krupka, D.Saunders, Elsevier, 725–772 (2008)
23. Kruglikov, B., Morozov, O.: Integrable dispersionless PDE in 4D, their symmetry pseudogroups and deformations. *Lett. Math. Phys.* **105**, 1703–1723 (2015)
24. Manakov, S.V., Santini, P.M.: The Cauchy problem on the plane for the dispersionless Kadomtsev-Petviashvili equation. *JETP Lett.* **83**, 462–466 (2006)
25. Marvan, M.: On the spectral parameter problem. *Acta Appl. Math.* **109**, 239–255 (2010)
26. Mason, L.J., Woodhouse, N.M.J.: Integrability, self-duality, and twistor theory, *L.M.S. Monographs* **15**, Oxford University Press (1996)
27. Odesskii, A.V., Sokolov, V.V.: Integrable pseudopotentials related to generalized hypergeometric functions. *Selecta Math. (N.S.)* **16**, 145–172 (2010)
28. Pavlov, M.V., Chang, J.H., Chen, Y.T.: Integrability of the Manakov–Santini hierarchy, [arXiv:0910.2400](https://arxiv.org/abs/0910.2400) (2009)
29. Penrose, R.: Nonlinear gravitons and curved twistor theory. *Gen. Relat. Gravitation* **7**(1), 31–52 (1976)
30. Plebański, J.F.: Some solutions of complex Einstein equations. *J. Math. Phys.* **16**, 2395–2402 (1975)
31. Plebański, J.F., Robinson, I.: Left-degenerate vacuum metrics. *Phys. Rev. Lett.* **37**, 493–495 (1976)
32. Prykarpatski, A.K.: On the linearization covering technique and its application to integrable nonlinear differential systems. *SIGMA* **14**, 023 (2018)
33. Sergeyev, A.: New integrable $(3 + 1)$ -dimensional systems and contact geometry. *Lett. Math. Phys.* **108**, 359–376 (2018)
34. Spencer, D.C.: Over-determined systems of linear partial differential equations. *Bull. Am. Math. Soc.* **75**, 179–239 (1969)
35. Tsarev, S.P.: Geometry of hamiltonian systems of hydrodynamic type. Generalized hodograph method, *Izvestija AN USSR Math.* **54**(5), 1048–1068 (1990)
36. Vinogradov, A.M.: Category of nonlinear differential equations, *Global Analysis Studies and Applications I*, *Lect. Notes Math.* **1108**, Springer-Verlag, 77–102 (1984)
37. Ward, R.S.: Einstein-Weyl spaces and $SU(\infty)$ Toda fields, *Class. Quantum Grav.* **7**, no. 4 (1990)
38. Zakharov, V.E.: Dispersionless limit of integrable systems in $2 + 1$ dimensions. In: Ercolani, N.M., et al. (eds.) *Singular Limits of Dispersive Waves*, pp. 165–174. NY, Plenum Press (1994)